

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 April 2001 (12.04.2001)

PCT

(10) International Publication Number
WO 01/25273 A2

(51) International Patent Classification⁷: **C07K 14/00**

(21) International Application Number: **PCT/US00/27465**

(22) International Filing Date: **4 October 2000 (04.10.2000)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
60/157,459 4 October 1999 (04.10.1999) US

(71) Applicant (for all designated States except US): **CORIXA CORPORATION** [US/US]; Suite 200, 1124 Columbia Street, Seattle, WA 98104 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SKEIKY, Yasir, A., W.** [CA/US]; 15106 SE 47th Place, Bellevue, WA 98006 (US). **XU, Jiangchun** [US/US]; 15805 SE 43rd

Place, Bellevue, WA 98006 (US). **CHEEVER, Martin, A.** [US/US]; 6210 Southeast 22nd, Mercer Island, WA 98040 (US). **REED, Steven, G.** [US/US]; 2843 - 122nd Place NE, Bellevue, WA 98005 (US).

(74) Agents: **POTTER, Jane, E., R.**; Seed Intellectual Property Law Group PLLC, Suite 6300, 701 Fifth Avenue, Seattle, WA 98104-7092 et al. (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: **COMPOSITIONS AND METHODS FOR WT1 SPECIFIC IMMUNOTHERAPY**

EU: MGSDVRLNALLPA17PFLGGGGGCLPVSGLAQWAFVLDFAFPGASAYGSL
MC: MGSDVRLNALLPA17PFLGGGGGCLPVSGLAQWAFVLDFAFPGASAYGSL

EU: GGFAPFFA17PFLGGGGGCLPVSGLAQWAFVLDFAFPGASAYGSL
MC: GGFAPFFA17PFLGGGGGCLPVSGLAQWAFVLDFAFPGASAYGSL

EU: ACRYGPF17PFLGGGGGCLPVSGLAQWAFVLDFAFPGASAYGSL
MC: ACRYGPF17PFLGGGGGCLPVSGLAQWAFVLDFAFPGASAYGSL

EU: YGWTFSHRAACFFNESTFHEHFMGQQGSLGZCQY57PFPVYGCETFTDSCTG
MC: YGWTFSHRAACFFNESTFHEHFMGQQGSLGZCQY57PFPVYGCETFTDSCTG

EU: SQALLLRTPYSSDNLYQNTSQLECHTWIQMILGATLRGVAAGSSSTVWTE
MC: SQALLLRTPYSSDNLYQNTSQLECHTWIQMILGATLRGVAAGSSSTVWTE

EU: GQSNHSTGYESDNHETPILCGAQVLETEGTTRGQDVRVSGVAFPLVRSAS
MC: GQSNHSTGYESDNHETPILCGAQVLETEGTTRGQDVRVSGVAFPLVRSAS

EU: ETSERLPFICAVFGGGRVFFLSELQMSRNETGZEPYQCDPDCERFPR
MC: ETSERLPFICAVFGGGRVFFLSELQMSRNETGZEPYQCDPDCERFPR

EU: SDQLNLEQPFETGVEFQCHTCQPFGRSDELATETRTETGATSEKPFSCR
MC: SDQLNLEQPFETGVEFQCHTCQPFGRSDELATETRTETGATSEKPFSCR

EU: WFSQQRFPARSDLVHEHDEQRNMTLELAL
MC: WFSQQRFPARSDLVHEHDEQRNMTLELAL

(57) Abstract: Compositions and methods for the therapy of malignant diseases, such as leukemia and cancer, are disclosed. The compositions comprise one or more of a WT1 polynucleotide, a WT1 polypeptide, an antigen-presenting cell presenting a WT1 polypeptide, an antibody that specifically binds to a WT1 polypeptide; or a T cell that specifically reacts with a WT1 polypeptide. Such compositions may be used, for example, for the prevention and treatment of metastatic diseases.

WO 01/25273 A2



Published:

— *Without international search report and to be republished upon receipt of that report.*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

COMPOSITIONS AND METHODS FOR WT1 SPECIFIC IMMUNOTHERAPY

TECHNICAL FIELD

The present invention relates generally to the immunotherapy of malignant diseases such as leukemia and cancers. The invention is more specifically related to compositions for generating or enhancing an immune response to WT1, and to the use of such compositions for preventing and/or treating malignant diseases.

BACKGROUND OF THE INVENTION

Cancer and leukemia are significant health problems in the United States and throughout the world. Although advances have been made in detection and treatment of such diseases, no vaccine or other universally successful method for prevention or treatment of cancer and leukemia is currently available. Management of the diseases currently relies on a combination of early diagnosis and aggressive treatment, which may include one or more of a variety of treatments such as surgery, radiotherapy, chemotherapy and hormone therapy. The course of treatment for a particular cancer is often selected based on a variety of prognostic parameters, including an analysis of specific tumor markers. However, the use of established markers often leads to a result that is difficult to interpret, and the high mortality continues to be observed in many cancer patients.

Immunotherapies have the potential to substantially improve cancer and leukemia treatment and survival. Recent data demonstrate that leukemia can be cured by immunotherapy in the context of bone marrow transplantation (*e.g.*, donor lymphocyte infusions). Such therapies may involve the generation or enhancement of an immune response to a tumor-associated antigen (TAA). However, to date, relatively few TAAs are known and the generation of an immune response against such antigens has, with rare exceptions, not been shown to be therapeutically beneficial.

Accordingly, there is a need in the art for improved methods for leukemia and cancer prevention and therapy. The present invention fulfills these needs and further provides other related advantages.

SUMMARY OF THE INVENTION

5 Briefly stated, this invention provides compositions and methods for the diagnosis and therapy of diseases such as leukemia and cancer. In one aspect, the present invention provides polypeptides comprising an immunogenic portion of a native WT1, or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antisera and/or
10 T-cell lines or clones is not substantially diminished. Within certain embodiments, the polypeptide comprises no more than 16 consecutive amino acid residues of a native WT1 polypeptide. Within other embodiments, the polypeptide comprises an immunogenic portion of amino acid residues 1 - 174 of a native WT1 polypeptide or a variant thereof, wherein the polypeptide comprises no more than 16 consecutive amino acid residues
15 present within amino acids 175 to 449 of the native WT1 polypeptide. The immunogenic portion preferably binds to an MHC class I and/or class II molecule. Within certain embodiments, the polypeptide comprises a sequence selected from the group consisting of (a) sequences recited in any one or more of Tables II - XLVI, (b) variants of the foregoing sequences that differ in one or more substitutions, deletions, additions and/or insertions
20 such that the ability of the variant to react with antigen-specific antisera and/or T-cell lines or clones is not substantially diminished and (c) mimetics of the polypeptides recited above, such that the ability of the mimetic to react with antigen-specific antisera and/or T cell lines or clones is not substantially diminished.

Within other embodiments, the polypeptide comprises a sequence selected
25 from the group consisting of (a) ALLPAVPSL (SEQ ID NO:34), GATLKGVA (SEQ ID NO:88), CMTWNQMNL (SEQ ID NOs: 49 and 258), SCLESQPTI (SEQ ID NOs: 199 and 296), SCLESQPAI (SEQ ID NO:198), NLYQMTSQL (SEQ ID NOs: 147 and 284),

ALLPAVSSL (SEQ ID NOs: 35 and 255), RMFPNAPYL (SEQ ID NOs: 185 and 293), (b) variants of the foregoing sequences that differ in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antisera and/or T-cell lines or clones is not substantially diminished and (c) mimetics of the polypeptides recited above, such that the ability of the mimetic to react with antigen-specific antisera and/or T cell lines or clones is not substantially diminished. Mimetics may comprises amino acids in combination with one or more amino acid mimetics or may be entirely nonpeptide mimetics.

Within further aspects, the present invention provides polypeptides comprising a variant of an immunogenic portion of a WT1 protein, wherein the variant differs from the immunogenic portion due to substitutions at between 1 and 3 amino acid positions within the immunogenic portion such that the ability of the variant to react with antigen-specific antisera and/or T-cell lines or clones is enhanced relative to a native WT1 protein.

The present invention further provides WT1 polynucleotides that encode a WT1 polypeptide as described above.

Within other aspects, the present invention provides pharmaceutical compositions and vaccines. Pharmaceutical compositions may comprise a polypeptide or mimetic as described above and/or one or more of (i) a WT1 polynucleotide; (ii) an antibody or antigen-binding fragment thereof that specifically binds to a WT1 polypeptide; (iii) a T cell that specifically reacts with a WT1 polypeptide or (iv) an antigen-presenting cell that expresses a WT1 polypeptide, in combination with a pharmaceutically acceptable carrier or excipient. Vaccines comprise a polypeptide as described above and/or one or more of (i) a WT1 polynucleotide, (ii) an antigen-presenting cell that expresses a WT1 polypeptide or (iii) an anti-idiotypic antibody, and a non-specific immune response enhancer. Within certain embodiments, less than 23 consecutive amino acid residues, preferably less than 17 amino acid residues, of a native WT1 polypeptide are present within a WT1 polypeptide employed within such pharmaceutical compositions and vaccines. The

immune response enhancer may be an adjuvant. Preferably, an immune response enhancer enhances a T cell response.

The present invention further provides methods for enhancing or inducing an immune response in a patient, comprising administering to a patient a pharmaceutical composition or vaccine as described above. In certain embodiments, the patient is a human.

The present invention further provides methods for inhibiting the development of a malignant disease in a patient, comprising administering to a patient a pharmaceutical composition or vaccine as described above. Malignant diseases include, but are not limited to leukemias (*e.g.*, acute myeloid, acute lymphocytic and chronic myeloid) and cancers (*e.g.*, breast, lung, thyroid or gastrointestinal cancer or a melanoma). The patient may, but need not, be afflicted with the malignant disease, and the administration of the pharmaceutical composition or vaccine may inhibit the onset of such a disease, or may inhibit progression and/or metastasis of an existing disease.

The present invention further provides, within other aspects, methods for removing cells expressing WT1 from bone marrow and/or peripheral blood or fractions thereof, comprising contacting bone marrow, peripheral blood or a fraction of bone marrow or peripheral blood with T cells that specifically react with a WT1 polypeptide, wherein the step of contacting is performed under conditions and for a time sufficient to permit the removal of WT1 positive cells to less than 10%, preferably less than 5% and more preferably less than 1%, of the number of myeloid or lymphatic cells in the bone marrow, peripheral blood or fraction. Bone marrow, peripheral blood and fractions may be obtained from a patient afflicted with a disease associated with WT1 expression, or may be obtained from a human or non-human mammal not afflicted with such a disease.

Within related aspects, the present invention provides methods for inhibiting the development of a malignant disease in a patient, comprising administering to a patient bone marrow, peripheral blood or a fraction of bone marrow or peripheral blood prepared as described above. Such bone marrow, peripheral blood or fractions may be autologous,

or may be derived from a related or unrelated human or non-human animal (*e.g.*, syngeneic or allogeneic).

In other aspects, the present invention provides methods for stimulating (or priming) and/or expanding T cells, comprising contacting T cells with a WT1 polypeptide under conditions and for a time sufficient to permit the stimulation and/or expansion of T cells. Such T cells may be autologous, allogeneic, syngeneic or unrelated WT1-specific T cells, and may be stimulated *in vitro* or *in vivo*. Expanded T cells may, within certain embodiments, be present within bone marrow, peripheral blood or a fraction of bone marrow or peripheral blood, and may (but need not) be clonal. Within certain
5
10
embodiments, T cells may be present in a mammal during stimulation and/or expansion. WT1-specific T cells may be used, for example, within donor lymphocyte infusions.

Within related aspects, methods are provided for inhibiting the development of a malignant disease in a patient, comprising administering to a patient T cells prepared as described above. Such T cells may, within certain embodiments, be autologous,
15
syngeneic or allogeneic.

The present invention further provides, within other aspects, methods for monitoring the effectiveness of an immunization or therapy for a malignant disease associated with WT1 expression in a patient. Such methods are based on monitoring antibody, CD4+ T cell and/or CD8+ T cell responses in the patient. Within certain such
20
aspects, a method may comprise the steps of: (a) incubating a first biological sample with one or more of: (i) a WT1 polypeptide; (ii) a polynucleotide encoding a WT1 polypeptide; or (iii) an antigen presenting cell that expresses a WT1 polypeptide, wherein the first biological sample is obtained from a patient prior to a therapy or immunization, and wherein the incubation is performed under conditions and for a time sufficient to allow
25
immunocomplexes to form; (b) detecting immunocomplexes formed between the WT1 polypeptide and antibodies in the biological sample that specifically bind to the WT1 polypeptide; (c) repeating steps (a) and (b) using a second biological sample obtained from the same patient following therapy or immunization; and (d) comparing the number of

immunocomplexes detected in the first and second biological samples, and therefrom monitoring the effectiveness of the therapy or immunization in the patient.

Within certain embodiments of the above methods, the step of detecting comprises (a) incubating the immunocomplexes with a detection reagent that is capable of binding to the immunocomplexes, wherein the detection reagent comprises a reporter group, (b) removing unbound detection reagent, and (c) detecting the presence or absence of the reporter group. The detection reagent may comprise, for example, a second antibody, or antigen-binding fragment thereof, capable of binding to the antibodies that specifically bind to the WT1 polypeptide or a molecule such as Protein A. Within other embodiments, a reporter group is bound to the WT1 polypeptide, and the step of detecting comprises removing unbound WT1 polypeptide and subsequently detecting the presence or absence of the reporter group.

Within further aspects, methods for monitoring the effectiveness of an immunization or therapy for a malignant disease associated with WT1 expression in a patient may comprise the steps of: (a) incubating a first biological sample with one or more of: (i) a WT1 polypeptide; (ii) a polynucleotide encoding a WT1 polypeptide; or (iii) an antigen presenting cell that expresses a WT1 polypeptide, wherein the biological sample comprises CD4+ and/or CD8+ T cells and is obtained from a patient prior to a therapy or immunization, and wherein the incubation is performed under conditions and for a time sufficient to allow specific activation, proliferation and/or lysis of T cells; (b) detecting an amount of activation, proliferation and/or lysis of the T cells; (c) repeating steps (a) and (b) using a second biological sample comprising CD4+ and/or CD8+ T cells, wherein the second biological sample is obtained from the same patient following therapy or immunization; and (d) comparing the amount of activation, proliferation and/or lysis of T cells in the first and second biological samples, and therefrom monitoring the effectiveness of the therapy or immunization in the patient.

The present invention further provides methods for inhibiting the development of a malignant disease associated with WT1 expression in a patient,

comprising the steps of: (a) incubating CD4⁺ and/or CD8⁺ T cells isolated from a patient with one or more of: (i) a WT1 polypeptide; (ii) a polynucleotide encoding a WT1 polypeptide; or (iii) an antigen presenting cell that expresses a WT1 polypeptide, such that the T cells proliferate; and (b) administering to the patient an effective amount of the
5 proliferated T cells, and therefrom inhibiting the development of a malignant disease in the patient. Within certain embodiments, the step of incubating the T cells may be repeated one or more times.

Within other aspects, the present invention provides methods for inhibiting the development of a malignant disease associated with WT1 expression in a patient,
10 comprising the steps of: (a) incubating CD4⁺ and/or CD8⁺ T cells isolated from a patient with one or more of: (i) a WT1 polypeptide; (ii) a polynucleotide encoding a WT1 polypeptide; or (iii) an antigen presenting cell that expresses a WT1 polypeptide, such that the T cells proliferate; (b) cloning one or more cells that proliferated; and (c) administering to the patient an effective amount of the cloned T cells.

15 Within other aspects, methods are provided for determining the presence or absence of a malignant disease associated with WT1 expression in a patient, comprising the steps of: (a) incubating CD4⁺ and/or CD8⁺ T cells isolated from a patient with one or more of: (i) a WT1 polypeptide; (ii) a polynucleotide encoding a WT1 polypeptide; or (iii) an antigen presenting cell that expresses a WT1 polypeptide; and (b) detecting the presence
20 or absence of specific activation of the T cells, therefrom determining the presence or absence of a malignant disease associated with WT1 expression. Within certain embodiments, the step of detecting comprises detecting the presence or absence of proliferation of the T cells.

Within further aspects, the present invention provides methods for
25 determining the presence or absence of a malignant disease associated with WT1 expression in a patient, comprising the steps of: (a) incubating a biological sample obtained from a patient with one or more of: (i) a WT1 polypeptide; (ii) a polynucleotide encoding a WT1 polypeptide; or (iii) an antigen presenting cell that expresses a WT1

polypeptide, wherein the incubation is performed under conditions and for a time sufficient to allow immunocomplexes to form; and (b) detecting immunocomplexes formed between the WT1 polypeptide and antibodies in the biological sample that specifically bind to the WT1 polypeptide; and therefrom determining the presence or absence of a malignant disease associated with WT1 expression.

These and other aspects of the present invention will become apparent upon reference to the following detailed description and attached drawings. All references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated individually.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts a comparison of the mouse (MO) and human (HU) WT1 protein sequences (SEQ ID NOS: 320 and 319 respectively).

Figure 2 is a Western blot illustrating the detection of WT1 specific antibodies in patients with hematological malignancy (AML). Lane 1 shows molecular weight markers; lane 2 shows a positive control (WT1 positive human leukemia cell line immunoprecipitated with a WT1 specific antibody); lane 3 shows a negative control (WT1 positive cell line immunoprecipitated with mouse sera); and lane 4 shows a WT1 positive cell line immunoprecipitated with sera of a patient with AML. For lanes 2-4, the immunoprecipitate was separated by gel electrophoresis and probed with a WT1 specific antibody.

Figure 3 is a Western blot illustrating the detection of a WT1 specific antibody response in B6 mice immunized with TRAMP-C, a WT1 positive tumor cell line. Lanes 1, 3 and 5 show molecular weight markers, and lanes 2, 4 and 6 show a WT1 specific positive control (N180, Santa Cruz Biotechnology, polypeptide spanning 180 amino acids of the N-terminal region of the WT1 protein, migrating on the Western blot at 52 kD). The primary antibody used was WT180 in lane 2, sera of non-immunized B6 mice in lane 4 and sera of the immunized B6 mice in lane 6.

Figure 4 is a Western blot illustrating the detection of WT1 specific antibodies in mice immunized with representative WT1 peptides. Lanes 1, 3 and 5 show molecular weight markers and lanes 2, 4 and 6 show a WT1 specific positive control (N180, Santa Cruz Biotechnology, polypeptide spanning 180 amino acids of the N-terminal region of the WT1 protein, migrating on the Western blot at 52 kD). The primary antibody used was WT180 in lane 2, sera of non-immunized B6 mice in lane 4 and sera of the immunized B6 mice in lane 6.

Figures 5A to 5C are graphs illustrating the stimulation of proliferative T cell responses in mice immunized with representative WT1 peptides. Thymidine incorporation assays were performed using one T cell line and two different clones, as indicated, and results were expressed as cpm. Controls indicated on the x axis were no antigen (No Ag) and B6/media; antigens used were p6-22 human (p1), p117-139 (p2) or p244-262 human (p3).

Figure 6A and 6B are histograms illustrating the stimulation of proliferative T cell responses in mice immunized with representative WT1 peptides. Three weeks after the third immunization, spleen cells of mice that had been inoculated with Vaccine A or Vaccine B were cultured with medium alone (medium) or spleen cells and medium (B6/no antigen), B6 spleen cells pulsed with the peptides p6-22 (p6), p117-139 (p117), p244-262 (p244) (Vaccine A; Figure 6A) or p287-301 (p287), p299-313 (p299), p421-435 (p421) (Vaccine B; Figure 6B) and spleen cells pulsed with an irrelevant control peptide (irrelevant peptide) at 25ug/ml and were assayed after 96hr for proliferation by (³H) thymidine incorporation. Bars represent the stimulation index (SI), which is calculated as the mean of the experimental wells divided by the mean of the control (B6 spleen cells with no antigen).

Figures 7A-7D are histograms illustrating the generation of proliferative T-cell lines and clones specific for p117-139 and p6-22. Following *in vivo* immunization, the initial three *in vitro* stimulations (IVS) were carried out using all three peptides of Vaccine A or B, respectively. Subsequent IVS were carried out as single peptide stimulations using

only the two relevant peptides p117-139 and p6-22. Clones were derived from both the p6-22 and p117-139 specific T cell lines, as indicated. T cells were cultured with medium alone (medium) or spleen cells and medium (B6/no antigen), B6 spleen cells pulsed with the peptides p6-22 (p6), p117-139 (p117) or an irrelevant control peptide (irrelevant peptide) at 25ug/ml and were assayed after 96hr for proliferation by (³H) thymidine incorporation. Bars represent the stimulation index (SI), which is calculated as the mean of the experimental wells divided by the mean of the control (B6 spleen cells with no antigen).

Figures 8A and 8B present the results of TSITES Analysis of human WT1 (SEQ ID NO:319) for peptides that have the potential to elicit Th responses. Regions indicated by "A" are AMPHI midpoints of blocks, "R" indicates residues matching the Rothbard/Taylor motif, "D" indicates residues matching the IAd motif, and 'd' indicates residues matching the IEd motif.

Figures 9A and 9B are graphs illustrating the elicitation of WT1 peptide-specific CTL in mice immunized with WT1 peptides. Figure 9A illustrates the lysis of target cells by allogeneic cell lines and Figure 9B shows the lysis of peptide coated cell lines. In each case, the % lysis (as determined by standard chromium release assays) is shown at three indicated effector:target ratios. Results are provided for lymphoma cells (LSTRA and E10), as well as E10 + p235-243 (E10+P235). E10 cells are also referred to herein as EL-4 cells.

Figures 10A-10D are graphs illustrating the elicitation of WT1 specific CTL, which kill WT1 positive tumor cell lines but do not kill WT1 negative cell lines, following vaccination of B6 mice with WT1 peptide P117. Figure 10A illustrates that T-cells of non-immunized B6 mice do not kill WT1 positive tumor cell lines. Figure 10B illustrates the lysis of the target cells by allogeneic cell lines. Figures 10C and 10D demonstrate the lysis of WT1 positive tumor cell lines, as compared to WT1 negative cell lines in two different experiments. In addition, Figures 10C and 10D show the lysis of peptide-coated cell lines (WT1 negative cell line E10 coated with the relevant WT1 peptide

P117) In each case, the % lysis (as determined by standard chromium release assays) is shown at three indicated effector:target ratios. Results are provided for lymphoma cells (E10), prostate cancer cells (TRAMP-C), a transformed fibroblast cell line (BLK-SV40), as well as E10+p117.

5 Figures 11A and 11B are histograms illustrating the ability of representative peptide P117-139 specific CTL to lyse WT1 positive tumor cells. Three weeks after the third immunization, spleen cells of mice that had been inoculated with the peptides p235-243 or p117-139 were stimulated *in vitro* with the relevant peptide and tested for ability to lyse targets incubated with WT1 peptides as well as WT1 positive and negative tumor cells.

10 The bars represent the mean % specific lysis in chromium release assays performed in triplicate with an E:T ratio of 25:1. Figure 11A shows the cytotoxic activity of the p235-243 specific T cell line against the WT1 negative cell line EL-4 (EL-4, WT1 negative); EL-4 pulsed with the relevant (used for immunization as well as for restimulation) peptide p235-243 (EL-4+p235); EL-4 pulsed with the irrelevant peptides p117-139 (EL-4+p117),

15 p126-134 (EL-4+p126) or p130-138 (EL-4+p130) and the WT1 positive tumor cells BLK-SV40 (BLK-SV40, WT1 positive) and TRAMP-C (TRAMP-C, WT1 positive), as indicated. Figure 11B shows cytotoxic activity of the p117-139 specific T cell line against EL-4; EL-4 pulsed with the relevant peptide P117-139 (EL-4+p117) and EL-4 pulsed with the irrelevant peptides p123-131 (EL-4+p123), or p128-136 (EL-4+p128); BLK-SV40 and

20 TRAMP-C, as indicated.

 Figures 12A and 12B are histograms illustrating the specificity of lysis of WT1 positive tumor cells, as demonstrated by cold target inhibition. The bars represent the mean % specific lysis in chromium release assays performed in triplicate with an E:T ratio of 25:1. Figure 12A shows the cytotoxic activity of the p117-139 specific T cell line

25 against the WT1 negative cell line EL-4 (EL-4, WT1 negative); the WT1 positive tumor cell line TRAMP-C (TRAMP-C, WT1 positive); TRAMP-C cells incubated with a ten-fold excess (compared to the hot target) of EL-4 cells pulsed with the relevant peptide p117-139 (TRAMP-C + p117 cold target) without ⁵¹Cr labeling and TRAMP-C cells incubated with

EL-4 pulsed with an irrelevant peptide without ^{51}Cr labeling (TRAMP-C + irrelevant cold target), as indicated. Figure 12B shows the cytotoxic activity of the p117-139 specific T cell line against the WT1 negative cell line EL-4 (EL-4, WT1 negative); the WT1 positive tumor cell line BLK-SV40 (BLK-SV40, WT1 positive); BLK-SV40 cells incubated with the relevant cold target (BLK-SV40 + p117 cold target) and BLK-SV40 cells incubated with the irrelevant cold target (BLK-SV40 + irrelevant cold target), as indicated.

Figures 13A-13C are histograms depicting an evaluation of the 9mer CTL epitope within p117-139. The p117-139 tumor specific CTL line was tested against peptides within aa117-139 containing or lacking an appropriate H-2^b class I binding motif and following restimulation with p126-134 or p130-138. The bars represent the mean % specific lysis in chromium release assays performed in triplicate with an E:T ratio of 25:1. Figure 13A shows the cytotoxic activity of the p117-139 specific T cell line against the WT1 negative cell line EL-4 (EL-4, WT1 negative) and EL-4 cells pulsed with the peptides p117-139 (EL-4 + p117), p119-127 (EL-4 + p119), p120-128 (EL-4 + p120), p123-131 (EL-4 + p123), p126-134 (EL-4 + p126), p128-136 (EL-4 + p128), and p130-138 (EL-4 + p130). Figure 13B shows the cytotoxic activity of the CTL line after restimulation with p126-134 against the WT1 negative cell line EL-4, EL-4 cells pulsed with p117-139 (EL-4 + p117), p126-134 (EL-4 + p126) and the WT1 positive tumor cell line TRAMP-C. Figure 13C shows the cytotoxic activity of the CTL line after restimulation with p130-138 against EL-4, EL-4 cells pulsed with p117-139 (EL-4 + p117), p130-138 (EL-4 + p130) and the WT1 positive tumor cell line TRAMP-C.

DETAILED DESCRIPTION OF THE INVENTION

As noted above, the present invention is generally directed to compositions and methods for the immunotherapy and diagnosis of malignant diseases. The compositions described herein may include WT1 polypeptides, WT1 polynucleotides, antigen-presenting cells (APC, *e.g.*, dendritic cells) that express a WT1 polypeptide, agents such as antibodies that bind to a WT1 polypeptide and/or immune system cells (*e.g.*, T

cells) specific for WT1. WT1 Polypeptides of the present invention generally comprise at least a portion of a Wilms Tumor gene product (WT1) or a variant thereof. Nucleic acid sequences of the subject invention generally comprise a DNA or RNA sequence that encodes all or a portion of such a polypeptide, or that is complementary to such a sequence.

5 Antibodies are generally immune system proteins, or antigen-binding fragments thereof, that are capable of binding to a portion of a WT1 polypeptide. T cells that may be employed within such compositions are generally T cells (*e.g.*, CD4⁺ and/or CD8⁺) that are specific for a WT1 polypeptide. Certain methods described herein further employ antigen-presenting cells that express a WT1 polypeptide as provided herein.

10 The present invention is based on the discovery that an immune response raised against a Wilms Tumor (WT) gene product (*e.g.*, WT1) can provide prophylactic and/or therapeutic benefit for patients afflicted with malignant diseases characterized by increased WT1 gene expression. Such diseases include, but are not limited to, leukemias (*e.g.*, acute myeloid leukemia (AML), chronic myeloid leukemia (CML), acute
15 lymphocytic leukemia (ALL) and childhood ALL), as well as many cancers such as lung, breast, thyroid and gastrointestinal cancers and melanomas. The WT1 gene was originally identified and isolated on the basis of a cytogenetic deletion at chromosome 11p13 in patients with Wilms' tumor (*see* Call et al., U.S. Patent No. 5,350,840). The gene consists of 10 exons and encodes a zinc finger transcription factor, and sequences of mouse and
20 human WT1 proteins are provided in Figure 1 and SEQ ID NOs: 319 and 320.

WT1 POLYPEPTIDES

Within the context of the present invention, a WT1 polypeptide is a polypeptide that comprises at least an immunogenic portion of a native WT1 (*i.e.*, a WT1
25 protein expressed by an organism that is not genetically modified), or a variant thereof, as described herein. A WT1 polypeptide may be of any length, provided that it comprises at least an immunogenic portion of a native protein or a variant thereof. In other words, a WT1 polypeptide may be an oligopeptide (*i.e.*, consisting of a relatively small number of

amino acid residues, such as 8-10 residues, joined by peptide bonds), a full length WT1 protein (*e.g.*, present within a human or non-human animal, such as a mouse) or a polypeptide of intermediate size. Within certain embodiments, the use of WT1 polypeptides that contain a small number of consecutive amino acid residues of a native WT1 polypeptide is preferred. Such polypeptides are preferred for certain uses in which the generation of a T cell response is desired. For example, such a WT1 polypeptide may contain less than 23, preferably no more than 18, and more preferably no more than 15 consecutive amino acid residues, of a native WT1 polypeptide. Polypeptides comprising nine consecutive amino acid residues of a native WT1 polypeptide are generally suitable for such purposes. Additional sequences derived from the native protein and/or heterologous sequences may be present within any WT1 polypeptide, and such sequences may (but need not) possess further immunogenic or antigenic properties. Polypeptides as provided herein may further be associated (covalently or noncovalently) with other polypeptide or non-polypeptide compounds.

An "immunogenic portion," as used herein is a portion of a polypeptide that is recognized (*i.e.*, specifically bound) by a B-cell and/or T-cell surface antigen receptor. Certain preferred immunogenic portions bind to an MHC class I or class II molecule. As used herein, an immunogenic portion is said to "bind to" an MHC class I or class II molecule if such binding is detectable using any assay known in the art. For example, the ability of a polypeptide to bind to MHC class I may be evaluated indirectly by monitoring the ability to promote incorporation of ¹²⁵I labeled β 2-microglobulin (β 2m) into MHC class I/ β 2m/peptide heterotrimeric complexes (*see* Parker et al., *J. Immunol.* 152:163, 1994). Alternatively, functional peptide competition assays that are known in the art may be employed. Certain immunogenic portions have one or more of the sequences recited within one or more of Tables II - XIV. Representative immunogenic portions include, but are not limited to, RDLNALLPAVPSLG GGG (human WT1 residues 6-22; SEQ ID NO:1), PSQASSGQARMFPNAPYLPSCLE (human and mouse WT1 residues 117-139; SEQ ID NOs: 2 and 3 respectively), GATLKGVAAGSSSSVKWTE (human WT1 residues 244-

262; SEQ ID NO:4), GATLKGVAA (human WT1 residues 244-252; SEQ ID NO:88), CMTWNQMNL (human and mouse WT1 residues 235-243; SEQ ID NOs: 49 and 258 respectively), SCLESQPTI (mouse WT1 residues 136-144; SEQ ID NO:296), SCLESQPAI (human WT1 residues 136-144; SEQ ID NO:198), NLYQMTSQL (human and mouse WT1 residues 225-233; SEQ ID NOs: 147 and 284 respectively); ALLPAVSSL (mouse WT1 residues 10-18; SEQ ID NO:255); or RMFPNAPYL (human and mouse WT1 residues 126-134; SEQ ID NOs: 185 and 293 respectively). Further immunogenic portions are provided herein, and others may generally be identified using well known techniques, such as those summarized in Paul, *Fundamental Immunology*, 3rd ed., 243-247 (Raven Press, 1993) and references cited therein. Representative techniques for identifying immunogenic portions include screening polypeptides for the ability to react with antigen-specific antisera and/or T-cell lines or clones. An immunogenic portion of a native WT1 polypeptide is a portion that reacts with such antisera and/or T-cells at a level that is not substantially less than the reactivity of the full length WT1 (e.g., in an ELISA and/or T-cell reactivity assay). In other words, an immunogenic portion may react within such assays at a level that is similar to or greater than the reactivity of the full length polypeptide. Such screens may generally be performed using methods well known to those of ordinary skill in the art, such as those described in Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988.

Alternatively, immunogenic portions may be identified using computer analysis, such as the Tsites program (see Rothbard and Taylor, *EMBO J.* 7:93-100, 1988; Deavin et al., *Mol. Immunol.* 33:145-155, 1996), which searches for peptide motifs that have the potential to elicit Th responses. CTL peptides with motifs appropriate for binding to murine and human class I or class II MHC may be identified according to BIMAS (Parker et al., *J. Immunol.* 152:163, 1994) and other HLA peptide binding prediction analyses. To confirm immunogenicity, a peptide may be tested using an HLA A2 transgenic mouse model and/or an *in vitro* stimulation assay using dendritic cells, fibroblasts or peripheral blood cells.

As noted above, a composition may comprise a variant of a native WT1 protein. A polypeptide "variant," as used herein, is a polypeptide that differs from a native polypeptide in one or more substitutions, deletions, additions and/or insertions, such that the immunogenicity of the polypeptide is retained (*i.e.*, the ability of the variant to react with antigen-specific antisera and/or T-cell lines or clones is not substantially diminished relative to the native polypeptide). In other words, the ability of a variant to react with antigen-specific antisera and/or T-cell lines or clones may be enhanced or unchanged, relative to the native polypeptide, or may be diminished by less than 50%, and preferably less than 20%, relative to the native polypeptide. Such variants may generally be identified by modifying one of the above polypeptide sequences and evaluating the reactivity of the modified polypeptide with antisera and/or T-cells as described herein. It has been found, within the context of the present invention, that a relatively small number of substitutions (*e.g.*, 1 to 3) within an immunogenic portion of a WT1 polypeptide may serve to enhance the ability of the polypeptide to elicit an immune response. Suitable substitutions may generally be identified by using computer programs, as described above, and the effect confirmed based on the reactivity of the modified polypeptide with antisera and/or T-cells as described herein. Accordingly, within certain preferred embodiments, a WT1 polypeptide comprises a variant in which 1 to 3 amino acid residues within an immunogenic portion are substituted such that the ability to react with antigen-specific antisera and/or T-cell lines or clones is statistically greater than that for the unmodified polypeptide. Such substitutions are preferably located within an MHC binding site of the polypeptide, which may be identified as described above. Preferred substitutions allow increased binding to MHC class I or class II molecules.

Certain variants contain conservative substitutions. A "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydrophobic nature of the polypeptide to be substantially unchanged. Amino acid substitutions may generally be made on the basis of similarity in

polarity, charge, solubility, hydrophobicity, hydrophilicity and/or the amphipathic nature of the residues. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine and valine; glycine and alanine; asparagine and glutamine; and serine, threonine, phenylalanine and tyrosine. Other groups of amino acids that may represent conservative changes include: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his. A variant may also, or alternatively, contain nonconservative changes. Variants may also (or alternatively) be modified by, for example, the deletion or addition of amino acids that have minimal influence on the immunogenicity, secondary structure and hydrophobic nature of the polypeptide.

As noted above, WT1 polypeptides may be conjugated to a signal (or leader) sequence at the N-terminal end of the protein which co-translationally or post-translationally directs transfer of the protein. A polypeptide may also, or alternatively, be conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide (*e.g.*, poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

Within certain specific embodiments, a polypeptide may be a fusion protein that comprises multiple polypeptides as described herein, or that comprises at least one polypeptide as described herein and an unrelated sequence, such as a known tumor protein. A fusion partner may, for example, assist in providing T helper epitopes (an immunological fusion partner), preferably T helper epitopes recognized by humans, or may assist in expressing the protein (an expression enhancer) at higher yields than the native recombinant protein. Certain preferred fusion partners are both immunological and expression enhancing fusion partners. Other fusion partners may be selected so as to increase the solubility of the protein or to enable the protein to be targeted to desired

intracellular compartments. Still further fusion partners include affinity tags, which facilitate purification of the protein.

In certain embodiments, the present invention provides fusion proteins comprising a polypeptide disclosed herein together with at least one of the sequences disclosed in U.S. Patent Application No. 09/352,616, filed July 13, 1999, which is incorporated herein in its entirety. Preferably one or more of the following known prostate antigens may be employed in a fusion protein with WT1: prostate specific antigen (PSA); prostatic acid phosphatase (PAP); and prostate specific membrane antigen (PSMA). The protein sequences for PSMA, PAP and PSA are provided in SEQ ID NO: 327-329, respectively. In certain embodiments, the fusion proteins of the present invention comprise WT1, PSA, PAP and/or PSMA in combination with one or more of the following the inventive antigens: P501S (amino acid sequence provided in SEQ ID NO: 333); P703P (amino acid sequences provided in SEQ ID NO: 330-332); P704P (cDNA sequence provided in SEQ ID NO: 334); P712P (cDNA sequence provided in SEQ ID NO: 335); P775P (cDNA sequence provided in SEQ ID NO: 336); P776P (cDNA sequence provided in SEQ ID NO: 337); P790P (cDNA sequence provided in SEQ ID NO: 338). P711P (full-length cDNA sequence provided in SEQ ID NO: 339, with the corresponding amino acid sequence provided in SEQ ID NO: 340); P710P (cDNA sequence provided in SEQ ID NO: 341-345); P714P (cDNA sequence provided in SEQ ID NO: 346); P510S (cDNA sequence provided in SEQ ID NO: 347); P774P (cDNA sequence provided in SEQ ID NO: 348); P705P (cDNA sequence provided in SEQ ID NO: 349, with the corresponding amino acid sequence provided in SEQ ID NO: 350); P503S (full-length cDNA sequence provided in SEQ ID NO: 351, with the corresponding amino acid sequence provided in SEQ ID NO: 352); P713P (cDNA sequence provided in SEQ ID NO: 353); P780P (cDNA sequence provided in SEQ ID NO: 354); P788P (cDNA sequence provided in SEQ ID NO: 355);

The cDNA sequence of a fusion protein for WT1 and PSA is provided in SEQ ID NO: 356 with the corresponding amino acid provided in SEQ ID NO: 357. In preferred embodiments, the inventive fusion proteins comprise one of the following

combinations of antigens, or permutations: WT1 and PSA; WT1 and P703P; WT1 and P501S; WT1/P703P and P501S; WT1/PSA and P703P; WT1/PSA and P501S; WT1 and PAP; WT1/PAP and P703P; WT1/PAP and P501S; WT1 and PSMA; WT1/PSMA and P703P; WT1/PSMA and P501S; WT1/PSA/PAP and P703P; WT1/PSA/PAP and P501S; WT1/PAP/PSMA and P703P; WT1/PAP/PSMA and P501S; WT1/PSA/PSMA and P703P; WT1/PSA/PSMA and P501S; WT1/PSA/PAP and PSMA; WT1/PSA/PAP/PSMA and P703P; WT1/PSA/PAP/PSMA and P501S. One of skill in the art will appreciate that the order of polypeptides within a fusion protein can be altered without substantially changing the therapeutic, prophylactic or diagnostic properties of the fusion protein.

The fusion proteins described above are more immunogenic and will be effective in a greater number of prostate cancer patients than any of the individual components alone. The use of multiple antigens in the form of a fusion protein also lessens the likelihood of immunologic escape.

Fusion proteins may generally be prepared using standard techniques, including chemical conjugation. Preferably, a fusion protein is expressed as a recombinant protein, allowing the production of increased levels, relative to a non-fused protein, in an expression system. Briefly, DNA sequences encoding the polypeptide components may be assembled separately, and ligated into an appropriate expression vector. The 3' end of the DNA sequence encoding one polypeptide component is ligated, with or without a peptide linker, to the 5' end of a DNA sequence encoding the second polypeptide component so that the reading frames of the sequences are in phase. This permits translation into a single fusion protein that retains the biological activity of both component polypeptides.

A peptide linker sequence may be employed to separate the first and the second polypeptide components by a distance sufficient to ensure that each polypeptide folds into its secondary and tertiary structures. Such a peptide linker sequence is incorporated into the fusion protein using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary

structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene* 40:39-46, 1985; Murphy et al., *Proc. Natl. Acad. Sci. USA* 83:8258-8262, 1986; U.S. Patent No. 4,935,233 and U.S. Patent No. 4,751,180. The linker sequence may generally be from 1 to about 50 amino acids in length. Linker sequences are not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric interference.

The ligated DNA sequences are operably linked to suitable transcriptional or translational regulatory elements. The regulatory elements responsible for expression of DNA are located only 5' to the DNA sequence encoding the first polypeptides. Similarly, stop codons required to end translation and transcription termination signals are only present 3' to the DNA sequence encoding the second polypeptide.

Fusion proteins are also provided that comprise a polypeptide of the present invention together with an unrelated immunogenic protein. Preferably the immunogenic protein is capable of eliciting a recall response. Examples of such proteins include tetanus, tuberculosis and hepatitis proteins (*see*, for example, Stoute et al. *New Engl. J. Med.*, 336:86-91, 1997).

Within preferred embodiments, an immunological fusion partner is derived from protein D, a surface protein of the gram-negative bacterium *Haemophilus influenza B* (WO 91/18926). Preferably, a protein D derivative comprises approximately the first third of the protein (*e.g.*, the first N-terminal 100-110 amino acids), and a protein D derivative may be lipidated. Within certain preferred embodiments, the first 109 residues of a Lipoprotein D fusion partner is included on the N-terminus to provide the polypeptide with additional exogenous T-cell epitopes and to increase the expression level in *E. coli* (thus

functioning as an expression enhancer). The lipid tail ensures optimal presentation of the antigen to antigen presenting cells. Other fusion partners include the non-structural protein from influenzae virus, NS1 (hemagglutinin). Typically, the N-terminal 81 amino acids are used, although different fragments that include T-helper epitopes may be used.

5 In another embodiment, the immunological fusion partner is the protein known as LYTA, or a portion thereof (preferably a C-terminal portion). LYTA is derived from *Streptococcus pneumoniae*, which synthesizes an N-acetyl-L-alanine amidase known as amidase LYTA (encoded by the *LytA* gene; *Gene* 43:265-292, 1986). LYTA is an autolysin that specifically degrades certain bonds in the peptidoglycan backbone. The C-
10 terminal domain of the LYTA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of *E. coli* C-LYTA expressing plasmids useful for expression of fusion proteins. Purification of hybrid proteins containing the C-LYTA fragment at the amino terminus has been described (*see Biotechnology* 10:795-798, 1992). Within a preferred
15 embodiment, a repeat portion of LYTA may be incorporated into a fusion protein. A repeat portion is found in the C-terminal region starting at residue 178. A particularly preferred repeat portion incorporates residues 188-305.

 WT1 polypeptides may be prepared using any of a variety of well known techniques. Recombinant polypeptides encoded by a WT1 polynucleotide as described
20 herein may be readily prepared from the polynucleotide. In general, any of a variety of expression vectors known to those of ordinary skill in the art may be employed to express recombinant WT1 polypeptides. Expression may be achieved in any appropriate host cell that has been transformed or transfected with an expression vector containing a DNA molecule that encodes a recombinant polypeptide. Suitable host cells include prokaryotes,
25 yeast and higher eukaryotic cells. Preferably, the host cells employed are *E. coli*, yeast or a mammalian cell line such as COS or CHO. Supernatants from suitable host/vector systems which secrete recombinant protein or polypeptide into culture media may be first concentrated using a commercially available filter. The concentrate may then be applied to

a suitable purification matrix such as an affinity matrix or an ion exchange resin. Finally, one or more reverse phase HPLC steps can be employed to further purify a recombinant polypeptide. Such techniques may be used to prepare native polypeptides or variants thereof. For example, polynucleotides that encode a variant of a native polypeptide may
5 generally be prepared using standard mutagenesis techniques, such as oligonucleotide-directed site-specific mutagenesis, and sections of the DNA sequence may be removed to permit preparation of truncated polypeptides.

Certain portions and other variants may also be generated by synthetic means, using techniques well known to those of ordinary skill in the art. For example,
10 polypeptides having fewer than about 500 amino acids, preferably fewer than about 100 amino acids, and more preferably fewer than about 50 amino acids, may be synthesized. Polypeptides may be synthesized using any of the commercially available solid-phase techniques, such as the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain. *See Merrifield, J. Am. Chem. Soc.*
15 *85:2149-2146, 1963.* Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Applied BioSystems, Inc. (Foster City, CA), and may be operated according to the manufacturer's instructions.

In general, polypeptides and polynucleotides as described herein are isolated. An "isolated" polypeptide or polynucleotide is one that is removed from its
20 original environment. For example, a naturally-occurring protein is isolated if it is separated from some or all of the coexisting materials in the natural system. Preferably, such polypeptides are at least about 90% pure, more preferably at least about 95% pure and most preferably at least about 99% pure. A polynucleotide is considered to be isolated if, for example, it is cloned into a vector that is not a part of the natural environment.

25 Within further aspects, the present invention provides mimetics of WT1 polypeptides. Such mimetics may comprise amino acids linked to one or more amino acid mimetics (*i.e.*, one or more amino acids within the WT1 protein may be replaced by an amino acid mimetic) or may be entirely nonpeptide mimetics. An amino acid mimetic is a

compound that is conformationally similar to an amino acid such that it can be substituted for an amino acid within a WT1 polypeptide without substantially diminishing the ability to react with antigen-specific antisera and/or T cell lines or clones. A nonpeptide mimetic is a compound that does not contain amino acids, and that has an overall conformation that is similar to a WT1 polypeptide such that the ability of the mimetic to react with WT1-specific antisera and/or T cell lines or clones is not substantially diminished relative to the ability of a WT1 polypeptide. Such mimetics may be designed based on standard techniques (*e.g.*, nuclear magnetic resonance and computational techniques) that evaluate the three dimensional structure of a peptide sequence. Mimetics may be designed where one or more of the side chain functionalities of the WT1 polypeptide are replaced by groups that do not necessarily have the same size or volume, but have similar chemical and/or physical properties which produce similar biological responses. It should be understood that, within embodiments described herein, a mimetic may be substituted for a WT1 polypeptide.

15

WT1 POLYNUCLEOTIDES

Any polynucleotide that encodes a WT1 polypeptide as described herein is a WT1 polynucleotide encompassed by the present invention. Such polynucleotides may be single-stranded (coding or antisense) or double-stranded, and may be DNA (genomic, cDNA or synthetic) or RNA molecules. Additional coding or non-coding sequences may, but need not, be present within a polynucleotide of the present invention, and a polynucleotide may, but need not, be linked to other molecules and/or support materials.

WT1 polynucleotides may encode a native WT1 protein, or may encode a variant of WT1 as described herein. Polynucleotide variants may contain one or more substitutions, additions, deletions and/or insertions such that the immunogenicity of the encoded polypeptide is not diminished, relative to a native WT1 protein. The effect on the immunogenicity of the encoded polypeptide may generally be assessed as described herein. Preferred variants contain nucleotide substitutions, deletions, insertions and/or additions at

25

no more than 20%, preferably at no more than 10%, of the nucleotide positions that encode an immunogenic portion of a native WT1 sequence. Certain variants are substantially homologous to a native gene, or a portion thereof. Such polynucleotide variants are capable of hybridizing under moderately stringent conditions to a naturally occurring DNA
5 sequence encoding a WT1 polypeptide (or a complementary sequence). Suitable moderately stringent conditions include prewashing in a solution of 5 X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50°C-65°C, 5 X SSC, overnight; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC containing 0.1% SDS). Such hybridizing DNA sequences are also within the scope of this invention.

10 It will be appreciated by those of ordinary skill in the art that, as a result of the degeneracy of the genetic code, there are many nucleotide sequences that encode a WT1 polypeptide. Some of these polynucleotides bear minimal homology to the nucleotide sequence of any native gene. Nonetheless, polynucleotides that vary due to differences in codon usage are specifically contemplated by the present invention.

15 Once an immunogenic portion of WT1 is identified, as described above, a WT1 polynucleotide may be prepared using any of a variety of techniques. For example, a WT1 polynucleotide may be amplified from cDNA prepared from cells that express WT1. Such polynucleotides may be amplified via polymerase chain reaction (PCR). For this approach, sequence-specific primers may be designed based on the sequence of the
20 immunogenic portion and may be purchased or synthesized. For example, suitable primers for PCR amplification of a human WT1 gene include: first step - P118: 1434-1414: 5' GAG AGT CAG ACT TGA AAG CAGT 3' (SEQ ID NO:5) and P135: 5' CTG AGC CTC AGC AAA TGG GC 3' (SEQ ID NO:6); second step - P136: 5' GAG CAT GCA TGG GCT CCG ACG TGC GGG 3' (SEQ ID NO:7) and P137: 5' GGG GTA CCC ACT
25 GAA CGG TCC CCG A 3' (SEQ ID NO:8). Primers for PCR amplification of a mouse WT1 gene include: first step - P138: 5' TCC GAG CCG CAC CTC ATG 3' (SEQ ID NO:9) and P139: 5' GCC TGG GAT GCT GGA CTG 3' (SEQ ID NO:10), second step -

P140: 5' GAG CAT GCG ATG GGT TCC GAC GTG CGG 3' (SEQ ID NO:11) and P141: 5' GGG GTA CCT CAA AGC GCC ACG TGG AGT TT 3' (SEQ ID NO:12).

An amplified portion may then be used to isolate a full length gene from a human genomic DNA library or from a suitable cDNA library, using well known techniques. Alternatively, a full length gene can be constructed from multiple PCR fragments. WT1 polynucleotides may also be prepared by synthesizing oligonucleotide components, and ligating components together to generate the complete polynucleotide.

WT1 polynucleotides may also be synthesized by any method known in the art, including chemical synthesis (*e.g.*, solid phase phosphoramidite chemical synthesis). Modifications in a polynucleotide sequence may also be introduced using standard mutagenesis techniques, such as oligonucleotide-directed site-specific mutagenesis (*see* Adelman et al., *DNA* 2:183, 1983). Alternatively, RNA molecules may be generated by *in vitro* or *in vivo* transcription of DNA sequences encoding a WT1 polypeptide, provided that the DNA is incorporated into a vector with a suitable RNA polymerase promoter (such as T7 or SP6). Certain portions may be used to prepare an encoded polypeptide, as described herein. In addition, or alternatively, a portion may be administered to a patient such that the encoded polypeptide is generated *in vivo* (*e.g.*, by transfecting antigen-presenting cells such as dendritic cells with a cDNA construct encoding a WT1 polypeptide, and administering the transfected cells to the patient).

Polynucleotides that encode a WT1 polypeptide may generally be used for production of the polypeptide, *in vitro* or *in vivo*. WT1 polynucleotides that are complementary to a coding sequence (*i.e.*, antisense polynucleotides) may also be used as a probe or to inhibit WT1 expression. cDNA constructs that can be transcribed into antisense RNA may also be introduced into cells of tissues to facilitate the production of antisense RNA.

Any polynucleotide may be further modified to increase stability *in vivo*. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends; the use of phosphorothioate or 2' O-methyl rather than

phosphodiesterase linkages in the backbone; and/or the inclusion of nontraditional bases such as inosine, queosine and wybutosine, as well as acetyl- methyl-, thio- and other modified forms of adenine, cytidine, guanine, thymine and uridine.

Nucleotide sequences as described herein may be joined to a variety of other
5 nucleotide sequences using established recombinant DNA techniques. For example, a polynucleotide may be cloned into any of a variety of cloning vectors, including plasmids, phagemids, lambda phage derivatives and cosmids. Vectors of particular interest include expression vectors, replication vectors, probe generation vectors and sequencing vectors. In general, a vector will contain an origin of replication functional in at least one organism,
10 convenient restriction endonuclease sites and one or more selectable markers. Other elements will depend upon the desired use, and will be apparent to those of ordinary skill in the art.

Within certain embodiments, polynucleotides may be formulated so as to permit entry into a cell of a mammal, and expression therein. Such formulations are
15 particularly useful for therapeutic purposes, as described below. Those of ordinary skill in the art will appreciate that there are many ways to achieve expression of a polynucleotide in a target cell, and any suitable method may be employed. For example, a polynucleotide may be incorporated into a viral vector such as, but not limited to, adenovirus, adeno-associated virus, retrovirus, or vaccinia or other pox virus (*e.g.*, avian pox virus).
20 Techniques for incorporating DNA into such vectors are well known to those of ordinary skill in the art. A retroviral vector may additionally transfer or incorporate a gene for a selectable marker (to aid in the identification or selection of transduced cells) and/or a targeting moiety, such as a gene that encodes a ligand for a receptor on a specific target cell, to render the vector target specific. Targeting may also be accomplished using an
25 antibody, by methods known to those of ordinary skill in the art. cDNA constructs within such a vector may be used, for example, to transfect human or animal cell lines for use in establishing WT1 positive tumor models which may be used to perform tumor protection

and adoptive immunotherapy experiments to demonstrate tumor or leukemia-growth inhibition or lysis of such cells.

Other therapeutic formulations for polynucleotides include colloidal dispersion systems, such as macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. A preferred colloidal system for use as a delivery vehicle *in vitro* and *in vivo* is a liposome (*i.e.*, an artificial membrane vesicle). The preparation and use of such systems is well known in the art.

10 ANTIBODIES AND FRAGMENTS THEREOF

The present invention further provides binding agents, such as antibodies and antigen-binding fragments thereof, that specifically bind to a WT1 polypeptide. As used herein, an agent is said to "specifically bind" to a WT1 polypeptide if it reacts at a detectable level (within, for example, an ELISA) with a WT1 polypeptide, and does not react detectably with unrelated proteins under similar conditions. As used herein, "binding" refers to a noncovalent association between two separate molecules such that a "complex" is formed. The ability to bind may be evaluated by, for example, determining a binding constant for the formation of the complex. The binding constant is the value obtained when the concentration of the complex is divided by the product of the component concentrations. In general, two compounds are said to "bind," in the context of the present invention, when the binding constant for complex formation exceeds about 10^3 L/mol. The binding constant maybe determined using methods well known in the art.

Any agent that satisfies the above requirements may be a binding agent. In a preferred embodiment, a binding agent is an antibody or an antigen-binding fragment thereof. Certain antibodies are commercially available from, for example, Santa Cruz Biotechnology (Santa Cruz, CA). Alternatively, antibodies may be prepared by any of a variety of techniques known to those of ordinary skill in the art. *See, e.g.*, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In

general, antibodies can be produced by cell culture techniques, including the generation of monoclonal antibodies as described herein, or via transfection of antibody genes into suitable bacterial or mammalian cell hosts, in order to allow for the production of recombinant antibodies. In one technique, an immunogen comprising the polypeptide is initially injected into any of a wide variety of mammals (*e.g.*, mice, rats, rabbits, sheep or goats). In this step, the polypeptides of this invention may serve as the immunogen without modification. Alternatively, particularly for relatively short polypeptides, a superior immune response may be elicited if the polypeptide is joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. The immunogen is injected into the animal host, preferably according to a predetermined schedule incorporating one or more booster immunizations, and the animals are bled periodically. Polyclonal antibodies specific for the polypeptide may then be purified from such antisera by, for example, affinity chromatography using the polypeptide coupled to a suitable solid support.

Monoclonal antibodies specific for the antigenic polypeptide of interest may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.* 6:511-519, 1976, and improvements thereto. Briefly, these methods involve the preparation of immortal cell lines capable of producing antibodies having the desired specificity (*i.e.*, reactivity with the polypeptide of interest). Such cell lines may be produced, for example, from spleen cells obtained from an animal immunized as described above. The spleen cells are then immortalized by, for example, fusion with a myeloma cell fusion partner, preferably one that is syngeneic with the immunized animal. A variety of fusion techniques may be employed. For example, the spleen cells and myeloma cells may be combined with a nonionic detergent for a few minutes and then plated at low density on a selective medium that supports the growth of hybrid cells, but not myeloma cells. A preferred selection technique uses HAT (hypoxanthine, aminopterin, thymidine) selection. After a sufficient time, usually about 1 to 2 weeks, colonies of hybrids are observed. Single colonies are selected and their culture supernatants tested for binding activity against the polypeptide. Hybridomas having high reactivity and specificity are preferred.

Monoclonal antibodies may be isolated from the supernatants of growing hybridoma colonies. In addition, various techniques may be employed to enhance the yield, such as injection of the hybridoma cell line into the peritoneal cavity of a suitable vertebrate host, such as a mouse. Monoclonal antibodies may then be harvested from the ascites fluid or the blood. Contaminants may be removed from the antibodies by conventional techniques, such as chromatography, gel filtration, precipitation, and extraction. The polypeptides of this invention may be used in the purification process in, for example, an affinity chromatography step.

Within certain embodiments, the use of antigen-binding fragments of antibodies may be preferred. Such fragments include Fab fragments, which may be prepared using standard techniques. Briefly, immunoglobulins may be purified from rabbit serum by affinity chromatography on Protein A bead columns (Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988) and digested by papain to yield Fab and Fc fragments. The Fab and Fc fragments may be separated by affinity chromatography on protein A bead columns.

Monoclonal antibodies and fragments thereof may be coupled to one or more therapeutic agents. Suitable agents in this regard include radioactive tracers and chemotherapeutic agents, which may be used, for example, to purge autologous bone marrow *in vitro*). Representative therapeutic agents include radionuclides, differentiation inducers, drugs, toxins, and derivatives thereof. Preferred radionuclides include ^{90}Y , ^{123}I , ^{125}I , ^{131}I , ^{186}Re , ^{188}Re , ^{211}At , and ^{212}Bi . Preferred drugs include methotrexate, and pyrimidine and purine analogs. Preferred differentiation inducers include phorbol esters and butyric acid. Preferred toxins include ricin, abrin, diphtheria toxin, cholera toxin, gelonin, *Pseudomonas* exotoxin, Shigella toxin, and pokeweed antiviral protein. For diagnostic purposes, coupling of radioactive agents may be used to facilitate tracing of metastases or to determine the location of WT1-positive tumors.

A therapeutic agent may be coupled (*e.g.*, covalently bonded) to a suitable monoclonal antibody either directly or indirectly (*e.g.*, via a linker group). A direct

reaction between an agent and an antibody is possible when each possesses a substituent capable of reacting with the other. For example, a nucleophilic group, such as an amino or sulfhydryl group, on one may be capable of reacting with a carbonyl-containing group, such as an anhydride or an acid halide, or with an alkyl group containing a good leaving group (*e.g.*, a halide) on the other.

Alternatively, it may be desirable to couple a therapeutic agent and an antibody via a linker group. A linker group can function as a spacer to distance an antibody from an agent in order to avoid interference with binding capabilities. A linker group can also serve to increase the chemical reactivity of a substituent on an agent or an antibody, and thus increase the coupling efficiency. An increase in chemical reactivity may also facilitate the use of agents, or functional groups on agents, which otherwise would not be possible.

It will be evident to those skilled in the art that a variety of bifunctional or polyfunctional reagents, both homo- and hetero-functional (such as those described in the catalog of the Pierce Chemical Co., Rockford, IL), may be employed as the linker group. Coupling may be effected, for example, through amino groups, carboxyl groups, sulfhydryl groups or oxidized carbohydrate residues. There are numerous references describing such methodology, *e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.

Where a therapeutic agent is more potent when free from the antibody portion of the immunoconjugates of the present invention, it may be desirable to use a linker group which is cleavable during or upon internalization into a cell. A number of different cleavable linker groups have been described. The mechanisms for the intracellular release of an agent from these linker groups include cleavage by reduction of a disulfide bond (*e.g.*, U.S. Patent No. 4,489,710, to Spitler), by irradiation of a photolabile bond (*e.g.*, U.S. Patent No. 4,625,014, to Senter et al.), by hydrolysis of derivatized amino acid side chains (*e.g.*, U.S. Patent No. 4,638,045, to Kohn et al.), by serum complement-mediated hydrolysis (*e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.), and acid-catalyzed hydrolysis (*e.g.*, U.S. Patent No. 4,569,789, to Blattler et al.).

It may be desirable to couple more than one agent to an antibody. In one embodiment, multiple molecules of an agent are coupled to one antibody molecule. In another embodiment, more than one type of agent may be coupled to one antibody. Regardless of the particular embodiment, immunoconjugates with more than one agent
5 may be prepared in a variety of ways. For example, more than one agent may be coupled directly to an antibody molecule, or linkers which provide multiple sites for attachment can be used. Alternatively, a carrier can be used. A carrier may bear the agents in a variety of ways, including covalent bonding either directly or via a linker group. Suitable carriers include proteins such as albumins (*e.g.*, U.S. Patent No. 4,507,234, to Kato et al.), peptides
10 and polysaccharides such as aminodextran (*e.g.*, U.S. Patent No. 4,699,784, to Shih et al.). A carrier may also bear an agent by noncovalent bonding or by encapsulation, such as within a liposome vesicle (*e.g.*, U.S. Patent Nos. 4,429,008 and 4,873,088). Carriers specific for radionuclide agents include radiohalogenated small molecules and chelating compounds. For example, U.S. Patent No. 4,735,792 discloses representative
15 radiohalogenated small molecules and their synthesis. A radionuclide chelate may be formed from chelating compounds that include those containing nitrogen and sulfur atoms as the donor atoms for binding the metal, or metal oxide, radionuclide. For example, U.S. Patent No. 4,673,562, to Davison et al. discloses representative chelating compounds and their synthesis.

20 A variety of routes of administration for the antibodies and immunoconjugates may be used. Typically, administration will be intravenous, intramuscular, subcutaneous or in the bed of a resected tumor. It will be evident that the precise dose of the antibody/immunoconjugate will vary depending upon the antibody used, the antigen density on the tumor, and the rate of clearance of the antibody.

25 Also provided herein are anti-idiotypic antibodies that mimic an immunogenic portion of WT1. Such antibodies may be raised against an antibody, or antigen-binding fragment thereof, that specifically binds to an immunogenic portion of WT1, using well known techniques. Anti-idiotypic antibodies that mimic an immunogenic

portion of WT1 are those antibodies that bind to an antibody, or antigen-binding fragment thereof, that specifically binds to an immunogenic portion of WT1, as described herein.

T CELLS

5 Immunotherapeutic compositions may also, or alternatively, comprise T cells specific for WT1. Such cells may generally be prepared *in vitro* or *ex vivo*, using standard procedures. For example, T cells may be present within (or isolated from) bone marrow, peripheral blood or a fraction of bone marrow or peripheral blood of a mammal, such as a patient, using a commercially available cell separation system, such as the
10 CEPRATE™ system, available from CellPro Inc., Bothell WA (see also U.S. Patent No. 5,240,856; U.S. Patent No. 5,215,926; WO 89/06280; WO 91/16116 and WO 92/07243). Alternatively, T cells may be derived from related or unrelated humans, non-human animals, cell lines or cultures.

 T cells may be stimulated with WT1 polypeptide, polynucleotide encoding a
15 WT1 polypeptide and/or an antigen presenting cell (APC) that expresses a WT1 polypeptide. Such stimulation is performed under conditions and for a time sufficient to permit the generation of T cells that are specific for the WT1 polypeptide. Preferably, a WT1 polypeptide or polynucleotide is present within a delivery vehicle, such as a microsphere, to facilitate the generation of antigen-specific T cells. Briefly, T cells, which
20 may be isolated from a patient or a related or unrelated donor by routine techniques (such as by Ficoll/Hypaque density gradient centrifugation of peripheral blood lymphocytes), are incubated with WT1 polypeptide. For example, T cells may be incubated *in vitro* for 2-9 days (typically 4 days) at 37°C with WT1 polypeptide (*e.g.*, 5 to 25 µg/ml) or cells synthesizing a comparable amount of WT1 polypeptide. It may be desirable to incubate a
25 separate aliquot of a T cell sample in the absence of WT1 polypeptide to serve as a control.

 T cells are considered to be specific for a WT1 polypeptide if the T cells kill target cells coated with a WT1 polypeptide or expressing a gene encoding such a polypeptide. T cell specificity may be evaluated using any of a variety of standard

techniques. For example, within a chromium release assay or proliferation assay, a stimulation index of more than two fold increase in lysis and/or proliferation, compared to negative controls, indicates T cell specificity. Such assays may be performed, for example, as described in Chen et al., *Cancer Res.* 54:1065-1070, 1994. Alternatively, detection of the proliferation of T cells may be accomplished by a variety of known techniques. For example, T cell proliferation can be detected by measuring an increased rate of DNA synthesis (*e.g.*, by pulse-labeling cultures of T cells with tritiated thymidine and measuring the amount of tritiated thymidine incorporated into DNA). Other ways to detect T cell proliferation include measuring increases in interleukin-2 (IL-2) production, Ca^{2+} flux, or dye uptake, such as 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-tetrazolium. Alternatively, synthesis of lymphokines (such as interferon-gamma) can be measured or the relative number of T cells that can respond to a WT1 polypeptide may be quantified. Contact with a WT1 polypeptide (200 ng/ml - 100 $\mu\text{g/ml}$, preferably 100 ng/ml - 25 $\mu\text{g/ml}$) for 3 - 7 days should result in at least a two fold increase in proliferation of the T cells and/or contact as described above for 2-3 hours should result in activation of the T cells, as measured using standard cytokine assays in which a two fold increase in the level of cytokine release (*e.g.*, TNF or IFN- γ) is indicative of T cell activation (*see* Coligan et al., *Current Protocols in Immunology*, vol. 1, Wiley Interscience (Greene 1998). WT1 specific T cells may be expanded using standard techniques. Within preferred embodiments, the T cells are derived from a patient or a related or unrelated donor and are administered to the patient following stimulation and expansion.

T cells that have been activated in response to a WT1 polypeptide, polynucleotide or WT1-expressing APC may be CD4^+ and/or CD8^+ . Specific activation of CD4^+ or CD8^+ T cells may be detected in a variety of ways. Methods for detecting specific T cell activation include detecting the proliferation of T cells, the production of cytokines (*e.g.*, lymphokines), or the generation of cytolytic activity (*i.e.*, generation of cytotoxic T cells specific for WT1). For CD4^+ T cells, a preferred method for detecting specific T cell activation is the detection of the proliferation of T cells. For CD8^+ T cells, a

preferred method for detecting specific T cell activation is the detection of the generation of cytolytic activity.

For therapeutic purposes, CD4⁺ or CD8⁺ T cells that proliferate in response to the WT1 polypeptide, polynucleotide or APC can be expanded in number either *in vitro* or *in vivo*. Proliferation of such T cells *in vitro* may be accomplished in a variety of ways. For example, the T cells can be re-exposed to WT1 polypeptide, with or without the addition of T cell growth factors, such as interleukin-2, and/or stimulator cells that synthesize a WT1 polypeptide. The addition of stimulator cells is preferred where generating CD8⁺ T cell responses. T cells can be grown to large numbers *in vitro* with retention of specificity in response to intermittent restimulation with WT1 polypeptide. Briefly, for the primary *in vitro* stimulation (IVS), large numbers of lymphocytes (*e.g.*, greater than 4×10^7) may be placed in flasks with media containing human serum. WT1 polypeptide (*e.g.*, peptide at 10 µg/ml) may be added directly, along with tetanus toxoid (*e.g.*, 5 µg/ml). The flasks may then be incubated (*e.g.*, 37°C for 7 days). For a second IVS, T cells are then harvested and placed in new flasks with $2-3 \times 10^7$ irradiated peripheral blood mononuclear cells. WT1 polypeptide (*e.g.*, 10 µg/ml) is added directly. The flasks are incubated at 37°C for 7 days. On day 2 and day 4 after the second IVS, 2-5 units of interleukin-2 (IL-2) may be added. For a third IVS, the T cells may be placed in wells and stimulated with the individual's own EBV transformed B cells coated with the peptide. IL-2 may be added on days 2 and 4 of each cycle. As soon as the cells are shown to be specific cytotoxic T cells, they may be expanded using a 10 day stimulation cycle with higher IL-2 (20 units) on days 2, 4 and 6.

Alternatively, one or more T cells that proliferate in the presence of WT1 polypeptide can be expanded in number by cloning. Methods for cloning cells are well known in the art, and include limiting dilution. Responder T cells may be purified from the peripheral blood of sensitized patients by density gradient centrifugation and sheep red cell rosetting and established in culture by stimulating with the nominal antigen in the presence of irradiated autologous filler cells. In order to generate CD4⁺ T cell lines, WT1

polypeptide is used as the antigenic stimulus and autologous peripheral blood lymphocytes (PBL) or lymphoblastoid cell lines (LCL) immortalized by infection with Epstein Barr virus are used as antigen presenting cells. In order to generate CD8⁺ T cell lines, autologous antigen-presenting cells transfected with an expression vector which produces
5 WT1 polypeptide may be used as stimulator cells. Established T cell lines may be cloned 2-4 days following antigen stimulation by plating stimulated T cells at a frequency of 0.5 cells per well in 96-well flat-bottom plates with 1×10^6 irradiated PBL or LCL cells and recombinant interleukin-2 (rIL2) (50 U/ml). Wells with established clonal growth may be identified at approximately 2-3 weeks after initial plating and restimulated with appropriate
10 antigen in the presence of autologous antigen-presenting cells, then subsequently expanded by the addition of low doses of rIL2 (10 U/ml) 2-3 days following antigen stimulation. T cell clones may be maintained in 24-well plates by periodic restimulation with antigen and rIL2 approximately every two weeks.

Within certain embodiments, allogeneic T-cells may be primed (*i.e.*,
15 sensitized to WT1) *in vivo* and/or *in vitro*. Such priming may be achieved by contacting T cells with a WT1 polypeptide, a polynucleotide encoding such a polypeptide or a cell producing such a polypeptide under conditions and for a time sufficient to permit the priming of T cells. In general, T cells are considered to be primed if, for example, contact with a WT1 polypeptide results in proliferation and/or activation of the T cells, as
20 measured by standard proliferation, chromium release and/or cytokine release assays as described herein. A stimulation index of more than two fold increase in proliferation or lysis, and more than three fold increase in the level of cytokine, compared to negative controls, indicates T-cell specificity. Cells primed *in vitro* may be employed, for example, within a bone marrow transplantation or as donor lymphocyte infusion.

25

PHARMACEUTICAL COMPOSITIONS AND VACCINES

Within certain aspects, polypeptides, polynucleotides, antibodies and/or T cells may be incorporated into pharmaceutical compositions or vaccines. Alternatively, a

pharmaceutical composition may comprise an antigen-presenting cell (*e.g.*, a dendritic cell) transfected with a WT1 polynucleotide such that the antigen presenting cell expresses a WT1 polypeptide. Pharmaceutical compositions comprise one or more such compounds or cells and a physiologically acceptable carrier or excipient. Certain vaccines may comprise one or more such compounds or cells and a non-specific immune response enhancer, such as an adjuvant or a liposome (into which the compound is incorporated). Pharmaceutical compositions and vaccines may additionally contain a delivery system, such as biodegradable microspheres which are disclosed, for example, in U.S. Patent Nos. 4,897,268 and 5,075,109. Pharmaceutical compositions and vaccines within the scope of the present invention may also contain other compounds, which may be biologically active or inactive.

Within certain embodiments, pharmaceutical compositions and vaccines are designed to elicit T cell responses specific for a WT1 polypeptide in a patient, such as a human. In general, T cell responses may be favored through the use of relatively short polypeptides (*e.g.*, comprising less than 23 consecutive amino acid residues of a native WT1 polypeptide, preferably 4-16 consecutive residues, more preferably 8-16 consecutive residues and still more preferably 8-10 consecutive residues. Alternatively, or in addition, a vaccine may comprise a non-specific immune response enhancer that preferentially enhances a T cell response. In other words, the immune response enhancer may enhance the level of a T cell response to a WT1 polypeptide by an amount that is proportionally greater than the amount by which an antibody response is enhanced. For example, when compared to a standard oil based adjuvant, such as CFA, an immune response enhancer that preferentially enhances a T cell response may enhance a proliferative T cell response by at least two fold, a lytic response by at least 10%, and/or T cell activation by at least two fold compared to WT1-negative control cell lines, while not detectably enhancing an antibody response. The amount by which a T cell or antibody response to a WT1 polypeptide is enhanced may generally be determined using any representative technique known in the art, such as the techniques provided herein.

A pharmaceutical composition or vaccine may contain DNA encoding one or more of the polypeptides as described above, such that the polypeptide is generated *in situ*. As noted above, the DNA may be present within any of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid expression systems, bacterial and viral expression systems and mammalian expression systems. Appropriate nucleic acid expression systems contain the necessary DNA, cDNA or RNA sequences for expression in the patient (such as a suitable promoter and terminating signal). Bacterial delivery systems involve the administration of a bacterium (such as *Bacillus-Calmette-Guerrin*) that expresses an immunogenic portion of the polypeptide on its cell surface. In a preferred embodiment, the DNA may be introduced using a viral expression system (e.g., vaccinia or other pox virus, retrovirus, or adenovirus), which may involve the use of a non-pathogenic (defective), replication competent virus. Techniques for incorporating DNA into such expression systems are well known to those of ordinary skill in the art. The DNA may also be "naked," as described, for example, in Ulmer et al., *Science* 259:1745-1749, 1993 and reviewed by Cohen, *Science* 259:1691-1692, 1993. The uptake of naked DNA may be increased by coating the DNA onto biodegradable beads, which are efficiently transported into the cells.

As noted above, a pharmaceutical composition or vaccine may comprise an antigen-presenting cell that expresses a WT1 polypeptide. For therapeutic purposes, as described herein, the antigen presenting cell is preferably an autologous dendritic cell. Such cells may be prepared and transfected using standard techniques, such as those described by Reeves et al., *Cancer Res.* 56:5672-5677, 1996; Tuting et al., *J. Immunol.* 160:1139-1147, 1998; and Nair et al., *Nature Biotechnol.* 16:364-369, 1998). Expression of a WT1 polypeptide on the surface of an antigen-presenting cell may be confirmed by *in vitro* stimulation and standard proliferation as well as chromium release assays, as described herein.

While any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier will vary

depending on the mode of administration. Compositions of the present invention may be formulated for any appropriate manner of administration, including for example, topical, oral, nasal, intravenous, intracranial, intraperitoneal, subcutaneous or intramuscular administration. For parenteral administration, such as subcutaneous injection, the carrier
5 preferably comprises water, saline, alcohol, a fat, a wax or a buffer. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and magnesium carbonate, may be employed. Biodegradable microspheres (*e.g.*, polylactate polyglycolate) may also be employed as carriers for the pharmaceutical compositions of
10 this invention. For certain topical applications, formulation as a cream or lotion, using well known components, is preferred.

Such compositions may also comprise buffers (*e.g.*, neutral buffered saline or phosphate buffered saline), carbohydrates (*e.g.*, glucose, mannose, sucrose or dextrans), mannitol, proteins, polypeptides or amino acids such as glycine, antioxidants, chelating
15 agents such as EDTA or glutathione, adjuvants (*e.g.*, aluminum hydroxide) and/or preservatives. Alternatively, compositions of the present invention may be formulated as a lyophilizate. Compounds may also be encapsulated within liposomes using well known technology.

Any of a variety of non-specific immune response enhancers, such as
20 adjuvants, may be employed in the vaccines of this invention. Most adjuvants contain a substance designed to protect the antigen from rapid catabolism, such as aluminum hydroxide or mineral oil, and a stimulator of immune responses, such as lipid A, *Bordetella pertussis* or *Mycobacterium tuberculosis* derived proteins. Suitable non-specific immune response enhancers include alum-based adjuvants (*e.g.*, Alhydrogel, Rehydrigel, aluminum
25 phosphate, Algamulin, aluminum hydroxide); oil based adjuvants (Freund's adjuvant (FA), Specol, RIBI, TiterMax, Montanide ISA50 or Seppic MONTANIDE ISA 720; cytokines (*e.g.*, GM-CSF or Flt3-ligand); microspheres; nonionic block copolymer-based adjuvants; dimethyl dioctadecyl ammoniumbromide (DDA) based adjuvants AS-1, AS-2

(Smith Kline Beecham); Ribi Adjuvant system based adjuvants; QS21 (Aquila); saponin based adjuvants (crude saponin, the saponin Quil A); muramyl dipeptide (MDP) based adjuvants such as SAF (Syntex adjuvant in its microfluidized form (SAF-m)); dimethyl-dioctadecyl ammonium bromide (DDA); human complement based adjuvants *m. vaccae* and derivatives; immune stimulating complex (iscom) based adjuvants; inactivated toxins; and attenuated infectious agents (such as *M. tuberculosis*).

As noted above, within certain embodiments, immune response enhancers are chosen for their ability to preferentially elicit or enhance a T cell response (*e.g.*, CD4⁺ and/or CD8⁺) to a WT1 polypeptide. Such immune response enhancers are well known in the art, and include (but are not limited to) Montanide ISA50, Seppic MONTANIDE ISA 720, cytokines (*e.g.*, GM-CSF, Flt3-ligand), microspheres, dimethyl dioctadecyl ammoniumbromide (DDA) based adjuvants, AS-1 (Smith Kline Beecham), AS-2 (Smith Kline Beecham), Ribi Adjuvant system based adjuvants, QS21 (Aquila), saponin based adjuvants (crude saponin, the saponin Quil A), Syntex adjuvant in its microfluidized form (SAF-m), MV, ddMV (Genesis), immune stimulating complex (iscom) based adjuvants and inactivated toxins.

The compositions and vaccines described herein may be administered as part of a sustained release formulation (*i.e.*, a formulation such as a capsule or sponge that effects a slow release of compound following administration). Such formulations may generally be prepared using well known technology and administered by, for example, oral, rectal or subcutaneous implantation, or by implantation at the desired target site. Sustained-release formulations may contain a polypeptide, polynucleotide, antibody or cell dispersed in a carrier matrix and/or contained within a reservoir surrounded by a rate controlling membrane. Carriers for use within such formulations are biocompatible, and may also be biodegradable; preferably the formulation provides a relatively constant level of active component release. The amount of active compound contained within a sustained release formulation depends upon the site of implantation, the rate and expected duration of release and the nature of the condition to be treated or prevented.

THERAPY OF MALIGNANT DISEASES

In further aspects of the present invention, the compositions and vaccines described herein may be used to inhibit the development of malignant diseases (*e.g.*, progressive or metastatic diseases or diseases characterized by small tumor burden such as minimal residual disease). In general, such methods may be used to prevent, delay or treat a disease associated with WT1 expression. In other words, therapeutic methods provided herein may be used to treat an existing WT1-associated disease, or may be used to prevent or delay the onset of such a disease in a patient who is free of disease or who is afflicted with a disease that is not yet associated with WT1 expression.

As used herein, a disease is "associated with WT1 expression" if diseased cells (*e.g.*, tumor cells) at some time during the course of the disease generate detectably higher levels of a WT1 polypeptide than normal cells of the same tissue. Association of WT1 expression with a malignant disease does not require that WT1 be present on a tumor. For example, overexpression of WT1 may be involved with initiation of a tumor, but the protein expression may subsequently be lost. Alternatively, a malignant disease that is not characterized by an increase in WT1 expression may, at a later time, progress to a disease that is characterized by increased WT1 expression. Accordingly, any malignant disease in which diseased cells formerly expressed, currently express or are expected to subsequently express increased levels of WT1 is considered to be "associated with WT1 expression."

Immunotherapy may be performed using any of a variety of techniques, in which compounds or cells provided herein function to remove WT1-expressing cells from a patient. Such removal may take place as a result of enhancing or inducing an immune response in a patient specific for WT1 or a cell expressing WT1. Alternatively, WT1-expressing cells may be removed *ex vivo* (*e.g.*, by treatment of autologous bone marrow, peripheral blood or a fraction of bone marrow or peripheral blood). Fractions of bone marrow or peripheral blood may be obtained using any standard technique in the art.

Within such methods, pharmaceutical compositions and vaccines may be administered to a patient. As used herein, a “patient” refers to any warm-blooded animal, preferably a human. A patient may or may not be afflicted with a malignant disease. Accordingly, the above pharmaceutical compositions and vaccines may be used to prevent the onset of a disease (*i.e.*, prophylactically) or to treat a patient afflicted with a disease (*e.g.*, to prevent or delay progression and/or metastasis of an existing disease). A patient afflicted with a disease may have a minimal residual disease (*e.g.*, a low tumor burden in a leukemia patient in complete or partial remission or a cancer patient following reduction of the tumor burden after surgery radiotherapy and/or chemotherapy). Such a patient may be immunized to inhibit a relapse (*i.e.*, prevent or delay the relapse, or decrease the severity of a relapse). Within certain preferred embodiments, the patient is afflicted with a leukemia (*e.g.*, AML, CML, ALL or childhood ALL), a myelodysplastic syndrome (MDS) or a cancer (*e.g.*, gastrointestinal, lung, thyroid or breast cancer or a melanoma), where the cancer or leukemia is WT1 positive (*i.e.*, reacts detectably with an anti-WT1 antibody, as provided herein or expresses WT1 mRNA at a level detectable by RT-PCR, as described herein) or suffers from an autoimmune disease directed against WT1-expressing cells.

The compositions provided herein may be used alone or in combination with conventional therapeutic regimens such as surgery, irradiation, chemotherapy and/or bone marrow transplantation (autologous, syngeneic, allogeneic or unrelated). As discussed in greater detail below, binding agents and T cells as provided herein may be used for purging of autologous stem cells. Such purging may be beneficial prior to, for example, bone marrow transplantation or transfusion of blood or components thereof. Binding agents, T cells, antigen presenting cells (APC) and compositions provided herein may further be used for expanding and stimulating (or priming) autologous, allogeneic, syngeneic or unrelated WT1-specific T-cells *in vitro* and/or *in vivo*. Such WT1-specific T cells may be used, for example, within donor lymphocyte infusions.

Routes and frequency of administration, as well as dosage, will vary from individual to individual, and may be readily established using standard techniques. In

general, the pharmaceutical compositions and vaccines may be administered by injection (*e.g.*, intracutaneous, intramuscular, intravenous or subcutaneous), intranasally (*e.g.*, by aspiration) or orally. In some tumors, pharmaceutical compositions or vaccines may be administered locally (by, for example, rectocoloscopy, gastroscopy, videoendoscopy, angiography or other methods known in the art). Preferably, between 1 and 10 doses may be administered over a 52 week period. Preferably, 6 doses are administered, at intervals of 1 month, and booster vaccinations may be given periodically thereafter. Alternate protocols may be appropriate for individual patients. A suitable dose is an amount of a compound that, when administered as described above, is capable of promoting an anti-tumor immune response that is at least 10-50% above the basal (*i.e.*, untreated) level. Such response can be monitored by measuring the anti-tumor antibodies in a patient or by vaccine-dependent generation of cytolytic effector cells capable of killing the patient's tumor cells *in vitro*. Such vaccines should also be capable of causing an immune response that leads to an improved clinical outcome (*e.g.*, more frequent complete or partial remissions, or longer disease-free and/or overall survival) in vaccinated patients as compared to non-vaccinated patients. In general, for pharmaceutical compositions and vaccines comprising one or more polypeptides, the amount of each polypeptide present in a dose ranges from about 100 μ g to 5 mg. Suitable dose sizes will vary with the size of the patient, but will typically range from about 0.1 mL to about 5 mL.

In general, an appropriate dosage and treatment regimen provides the active compound(s) in an amount sufficient to provide therapeutic and/or prophylactic benefit. Such a response can be monitored by establishing an improved clinical outcome (*e.g.*, more frequent complete or partial remissions, or longer disease-free and/or overall survival) in treated patients as compared to non-treated patients. Increases in preexisting immune responses to WT1 generally correlate with an improved clinical outcome. Such immune responses may generally be evaluated using standard proliferation, cytotoxicity or cytokine assays, which may be performed using samples obtained from a patient before and after treatment.

Within further aspects, methods for inhibiting the development of a malignant disease associated with WT1 expression involve the administration of autologous T cells that have been activated in response to a WT1 polypeptide or WT1-expressing APC, as described above. Such T cells may be CD4⁺ and/or CD8⁺, and may be proliferated as described above. The T cells may be administered to the individual in an amount effective to inhibit the development of a malignant disease. Typically, about 1×10^9 to 1×10^{11} T cells/M² are administered intravenously, intracavitary or in the bed of a resected tumor. It will be evident to those skilled in the art that the number of cells and the frequency of administration will be dependent upon the response of the patient.

Within certain embodiments, T cells may be stimulated prior to an autologous bone marrow transplantation. Such stimulation may take place *in vivo* or *in vitro*. For *in vitro* stimulation, bone marrow and/or peripheral blood (or a fraction of bone marrow or peripheral blood) obtained from a patient may be contacted with a WT1 polypeptide, a polynucleotide encoding a WT1 polypeptide and/or an APC that expresses a WT1 polypeptide under conditions and for a time sufficient to permit the stimulation of T cells as described above. Bone marrow, peripheral blood stem cells and/or WT1-specific T cells may then be administered to a patient using standard techniques.

Within related embodiments, T cells of a related or unrelated donor may be stimulated prior to a syngeneic or allogeneic (related or unrelated) bone marrow transplantation. Such stimulation may take place *in vivo* or *in vitro*. For *in vitro* stimulation, bone marrow and/or peripheral blood (or a fraction of bone marrow or peripheral blood) obtained from a related or unrelated donor may be contacted with a WT1 polypeptide, WT1 polynucleotide and/or APC that expresses a WT1 polypeptide under conditions and for a time sufficient to permit the stimulation of T cells as described above. Bone marrow, peripheral blood stem cells and/or WT1-specific T cells may then be administered to a patient using standard techniques.

Within other embodiments, WT1-specific T cells as described herein may be used to remove cells expressing WT1 from autologous bone marrow, peripheral blood or a

fraction of bone marrow or peripheral blood (*e.g.*, CD34⁺ enriched peripheral blood (PB) prior to administration to a patient). Such methods may be performed by contacting bone marrow or PB with such T cells under conditions and for a time sufficient to permit the reduction of WT1 expressing cells to less than 10%, preferably less than 5% and more preferably less than 1%, of the total number of myeloid or lymphatic cells in the bone marrow or peripheral blood. The extent to which such cells have been removed may be readily determined by standard methods such as, for example, qualitative and quantitative PCR analysis, morphology, immunohistochemistry and FACS analysis. Bone marrow or PB (or a fraction thereof) may then be administered to a patient using standard techniques.

DIAGNOSTIC METHODS

The present invention further provides methods for detecting a malignant disease associated with WT1 expression, and for monitoring the effectiveness of an immunization or therapy for such a disease. Such methods are based on the discovery, within the present invention, that an immune response specific for WT1 protein can be detected in patients afflicted with such diseases, and that methods which enhance such immune responses may provide a preventive or therapeutic benefit.

To determine the presence or absence of a malignant disease associated with WT1 expression, a patient may be tested for the level of T cells specific for WT1. Within certain methods, a biological sample comprising CD4⁺ and/or CD8⁺ T cells isolated from a patient is incubated with a WT1 polypeptide, a polynucleotide encoding a WT1 polypeptide and/or an APC that expresses a WT1 polypeptide, and the presence or absence of specific activation of the T cells is detected, as described herein. Suitable biological samples include, but are not limited to, isolated T cells. For example, T cells may be isolated from a patient by routine techniques (such as by Ficoll/Hypaque density gradient centrifugation of peripheral blood lymphocytes). T cells may be incubated *in vitro* for 2-9 days (typically 4 days) at 37°C with WT1 polypeptide (*e.g.*, 5 - 25 µg/ml). It may be desirable to incubate another aliquot of a T cell sample in the absence of WT1 polypeptide

to serve as a control. For CD4⁺ T cells, activation is preferably detected by evaluating proliferation of the T cells. For CD8⁺ T cells, activation is preferably detected by evaluating cytolytic activity. A level of proliferation that is at least two fold greater and/or a level of cytolytic activity that is at least 20% greater than in disease-free patients indicates the presence of a malignant disease associated with WT1 expression. Further correlation may be made, using methods well known in the art, between the level of proliferation and/or cytolytic activity and the predicted response to therapy. In particular, patients that display a higher antibody, proliferative and/or lytic response may be expected to show a greater response to therapy.

Within other methods, a biological sample obtained from a patient is tested for the level of antibody specific for WT1. The biological sample is incubated with a WT1 polypeptide, a polynucleotide encoding a WT1 polypeptide and/or an APC that expresses a WT1 polypeptide under conditions and for a time sufficient to allow immunocomplexes to form. Immunocomplexes formed between the WT1 polypeptide and antibodies in the biological sample that specifically bind to the WT1 polypeptide are then detected. A biological sample for use within such methods may be any sample obtained from a patient that would be expected to contain antibodies. Suitable biological samples include blood, sera, ascites, bone marrow, pleural effusion, and cerebrospinal fluid.

The biological sample is incubated with the WT1 polypeptide in a reaction mixture under conditions and for a time sufficient to permit immunocomplexes to form between the polypeptide and antibodies specific for WT1. For example, a biological sample and WT1 polypeptide may be incubated at 4°C for 24-48 hours.

Following the incubation, the reaction mixture is tested for the presence of immunocomplexes. Detection of immunocomplexes formed between the WT1 polypeptide and antibodies present in the biological sample may be accomplished by a variety of known techniques, such as radioimmunoassays (RIA) and enzyme linked immunosorbent assays (ELISA). Suitable assays are well known in the art and are amply described in the scientific and patent literature (*e.g.*, Harlow and Lane, *Antibodies: A Laboratory Manual*,

Cold Spring Harbor Laboratory, 1988). Assays that may be used include, but are not limited to, the double monoclonal antibody sandwich immunoassay technique of David et al. (U.S. Patent 4,376,110); monoclonal-polyclonal antibody sandwich assays (Wide et al., in Kirkham and Hunter, eds., *Radioimmunoassay Methods*, E. and S. Livingstone, 5 Edinburgh, 1970); the "western blot" method of Gordon et al. (U.S. Patent 4,452,901); immunoprecipitation of labeled ligand (Brown et al., *J. Biol. Chem.* 255:4980-4983, 1980); enzyme-linked immunosorbent assays as described by, for example, Raines and Ross (*J. Biol. Chem.* 257:5154-5160, 1982); immunocytochemical techniques, including the use of fluorochromes (Brooks et al., *Clin. Exp. Immunol.* 39: 477, 1980); and neutralization of 10 activity (Bowen-Pope et al., *Proc. Natl. Acad. Sci. USA* 81:2396-2400, 1984). Other immunoassays include, but are not limited to, those described in U.S. Patent Nos.: 3,817,827; 3,850,752; 3,901,654; 3,935,074; 3,984,533; 3,996,345; 4,034,074; and 4,098,876.

For detection purposes, WT1 polypeptide may either be labeled or 15 unlabeled. Unlabeled WT1 polypeptide may be used in agglutination assays or in combination with labeled detection reagents that bind to the immunocomplexes (*e.g.*, anti-immunoglobulin, protein G, protein A or a lectin and secondary antibodies, or antigen-binding fragments thereof, capable of binding to the antibodies that specifically bind to the WT1 polypeptide). If the WT1 polypeptide is labeled, the reporter group may be any 20 suitable reporter group known in the art, including radioisotopes, fluorescent groups, luminescent groups, enzymes, biotin and dye particles.

Within certain assays, unlabeled WT1 polypeptide is immobilized on a solid support. The solid support may be any material known to those of ordinary skill in the art to which the polypeptide may be attached. For example, the solid support may be a test 25 well in a microtiter plate or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681. The

polypeptide may be immobilized on the solid support using a variety of techniques known to those of skill in the art, which are amply described in the patent and scientific literature. In the context of the present invention, the term "immobilization" refers to both noncovalent association, such as adsorption, and covalent attachment (which may be a direct linkage between the antigen and functional groups on the support or may be a linkage by way of a cross-linking agent). Immobilization by adsorption to a well in a microtiter plate or to a membrane is preferred. In such cases, adsorption may be achieved by contacting the WT1 polypeptide, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and about 1 day. In general, contacting a well of a plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of polypeptide ranging from about 10 ng to about 10 μ g, and preferably about 100 ng to about 1 μ g, is sufficient to immobilize an adequate amount of polypeptide.

Following immobilization, the remaining protein binding sites on the support are typically blocked. Any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin, Tween 20TM (Sigma Chemical Co., St. Louis, MO), heat-inactivated normal goat serum (NGS), or BLOTTO (buffered solution of nonfat dry milk which also contains a preservative, salts, and an antifoaming agent). The support is then incubated with a biological sample suspected of containing specific antibody. The sample can be applied neat, or, more often, it can be diluted, usually in a buffered solution which contains a small amount (0.1%-5.0% by weight) of protein, such as BSA, NGS, or BLOTTO. In general, an appropriate contact time (*i.e.*, incubation time) is a period of time that is sufficient to detect the presence of antibody that specifically binds WT1 within a sample containing such an antibody. Preferably, the contact time is sufficient to achieve a level of binding that is at least about 95% of that achieved at equilibrium between bound and unbound antibody. Those of ordinary skill in the art will recognize that the time necessary to achieve equilibrium may be readily determined by assaying the level of

binding that occurs over a period of time. At room temperature, an incubation time of about 30 minutes is generally sufficient.

Unbound sample may then be removed by washing the solid support with an appropriate buffer, such as PBS containing 0.1% Tween 20™. A detection reagent that
5 binds to the immunocomplexes and that comprises a reporter group may then be added. The detection reagent is incubated with the immunocomplex for an amount of time sufficient to detect the bound antibody. An appropriate amount of time may generally be determined by assaying the level of binding that occurs over a period of time. Unbound
10 detection reagent is then removed and bound detection reagent is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the reporter group. For radioactive groups, scintillation counting or autoradiographic methods are generally appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and fluorescent groups. Biotin may be detected using
15 avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups (*e.g.*, horseradish peroxidase, beta-galactosidase, alkaline phosphatase and glucose oxidase) may generally be detected by the addition of substrate (generally for a specific period of time), followed by spectroscopic or other analysis of the reaction products. Regardless of the specific method employed, a level of
20 bound detection reagent that is at least two fold greater than background (*i.e.*, the level observed for a biological sample obtained from a disease-free individual) indicates the presence of a malignant disease associated with WT1 expression.

In general, methods for monitoring the effectiveness of an immunization or therapy involve monitoring changes in the level of antibodies or T cells specific for WT1 in the patient. Methods in which antibody levels are monitored may comprise the steps of:
25 (a) incubating a first biological sample, obtained from a patient prior to a therapy or immunization, with a WT1 polypeptide, wherein the incubation is performed under conditions and for a time sufficient to allow immunocomplexes to form; (b) detecting immunocomplexes formed between the WT1 polypeptide and antibodies in the biological

sample that specifically bind to the WT1 polypeptide; (c) repeating steps (a) and (b) using a second biological sample taken from the patient following therapy or immunization; and (d) comparing the number of immunocomplexes detected in the first and second biological samples. Alternatively, a polynucleotide encoding a WT1 polypeptide, or an APC
5 expressing a WT1 polypeptide may be employed in place of the WT1 polypeptide. Within such methods, immunocomplexes between the WT1 polypeptide encoded by the polynucleotide, or expressed by the APC, and antibodies in the biological sample are detected.

Methods in which T cell activation and/or the number of WT1 specific
10 precursors are monitored may comprise the steps of: (a) incubating a first biological sample comprising CD4+ and/or CD8+ cells (*e.g.*, bone marrow, peripheral blood or a fraction thereof), obtained from a patient prior to a therapy or immunization, with a WT1 polypeptide, wherein the incubation is performed under conditions and for a time sufficient to allow specific activation, proliferation and/or lysis of T cells; (b) detecting an amount of
15 activation, proliferation and/or lysis of the T cells; (c) repeating steps (a) and (b) using a second biological sample comprising CD4+ and/or CD8+ T cells, and taken from the same patient following therapy or immunization; and (d) comparing the amount of activation, proliferation and/or lysis of T cells in the first and second biological samples. Alternatively, a polynucleotide encoding a WT1 polypeptide, or an APC expressing a WT1
20 polypeptide may be employed in place of the WT1 polypeptide.

A biological sample for use within such methods may be any sample obtained from a patient that would be expected to contain antibodies, CD4+ T cells and/or CD8+ T cells. Suitable biological samples include blood, sera, ascites, bone marrow, pleural effusion and cerebrospinal fluid. A first biological sample may be obtained prior to
25 initiation of therapy or immunization or part way through a therapy or vaccination regime. The second biological sample should be obtained in a similar manner, but at a time following additional therapy or immunization. The second biological sample may be obtained at the completion of, or part way through, therapy or immunization, provided that

at least a portion of therapy or immunization takes place between the isolation of the first and second biological samples.

Incubation and detection steps for both samples may generally be performed as described above. A statistically significant increase in the number of immunocomplexes
5 in the second sample relative to the first sample reflects successful therapy or immunization.

The following Examples are offered by way of illustration and not by way of limitation.

EXAMPLES

Example 1Identification of an Immune Response to WT1
in Patients with Hematological Malignancies

This Example illustrates the identification of an existent immune response in patients with a hematological malignancy.

To evaluate the presence of preexisting WT1 specific antibody responses in patients, sera of patients with AML, ALL, CML and severe aplastic anemia were analyzed using Western blot analysis. Sera were tested for the ability to immunoprecipitate WT1 from the human leukemic cell line K562 (American Type Culture Collection, Manassas, VA). In each case, immunoprecipitates were separated by gel electrophoresis, transferred to membrane and probed with the anti WT-1 antibody WT180 (Santa Cruz Biotechnology, Inc., Santa Cruz, CA). This Western blot analysis identified potential WT1 specific antibodies in patients with hematological malignancy. A representative Western blot showing the results for a patient with AML is shown in Figure 2. A 52 kD protein in the immunoprecipitate generated using the patient sera was recognized by the WT1 specific antibody. The 52 kD protein migrated at the same size as the positive control.

Example 2Induction of Antibodies to WT1 in Mice Immunized with Cell Lines Expressing WT1

This Example illustrates the use of cells expressing WT1 to induce a WT1 specific antibody response *in vivo*.

Detection of existent antibodies to WT1 in patients with leukemia strongly implied that it is possible to immunize to WT1 protein to elicit immunity to WT1. To test whether immunity to WT1 can be generated by vaccination, mice were injected with TRAMP-C, a WT1 positive tumor cell line of B6 origin. Briefly, male B6 mice were

immunized with 5×10^6 TRAMP-C cells subcutaneously and boosted twice with 5×10^6 cells at three week intervals. Three weeks after the final immunization, sera were obtained and single cell suspensions of spleens were prepared in RPMI 1640 medium (GIBCO) with $25\mu\text{M}$ β -2-mercaptoethanol, 200 units of penicillin per ml, 10mM L-glutamine, and 10% fetal bovine serum.

Following immunization to TRAMP-C, a WT1 specific antibody response in the immunized animals was detectable. A representative Western blot is shown in Figure 3. These results show that immunization to WT1 protein can elicit an immune response to WT1 protein.

Example 3

Induction of Th and Antibody Responses in Mice Immunized with WT1 Peptides

This Example illustrates the ability of immunization with WT1 peptides to elicit an immune response specific for WT1.

Peptides suitable for eliciting Ab and proliferative T cell responses were identified according to the Tsites program (Rothbard and Taylor, *EMBO J.* 7:93-100, 1988; Deavin et al., *Mol. Immunol.* 33:145-155, 1996), which searches for peptide motifs that have the potential to elicit Th responses. Peptides shown in Table I were synthesized and sequenced.

Table I
WT1 Peptides

Peptide	Sequence	Comments
Mouse: p6-22	RDLNALLPAVSSLGGGG (SEQ ID NO:13)	1 mismatch relative to human WT1 sequence
Human: p6-22	RDLNALLPAVPSLGGGG (SEQ ID NO:1)	
Human/mouse: p117-139	PSQASSGQARMFPNAPYLPSCLE (SEQ ID NOS: 2 and 3)	

Mouse: p244-262	GATLKGMAAGSSSSSVKWTE (SEQ ID NO:14)	1 mismatch relative to human WT1 sequence
Human: p244-262	GATLKGVAAGSSSSSVKWTE (SEQ ID NO:4)	
Human/mouse: p287-301	RIHTHGVFRGIQDVR (SEQ ID NOs: 15 and 16)	
Mouse: p299-313	VRRVSGVAPTLVRS (SEQ ID NO:17)	1 mismatch relative to human WT1 sequence
Human/mouse: p421-435	CQKKFARSDELVRHH (SEQ ID NOs: 19 and 20)	

For immunization, peptides were grouped as follows:

- | | | |
|----|-----------------|--|
| 5 | <u>Group A:</u> | p6-22 human: 10.9mg in 1ml (10μl = 100μg)
p117-139 human/mouse: 7.6mg in 1ml (14μl = 100μg)
p244-262 human: 4.6.mg in 1ml (22μl = 100μg) |
| 10 | <u>Group B:</u> | p287-301 human/mouse: 7.2mg in 1ml (14μl = 100μg)
mouse p299-313: 6.6.mg in 1ml (15μl = 100μg)
p421-435 human/mouse: 3.3mg in 1ml (30μl = 100μg) |
| | <u>Control:</u> | (FBL peptide 100μg) + CFA/IFA |
| 15 | <u>Control:</u> | (CD45 peptide 100μg) + CFA/IFA |

Group A contained peptides present within the amino terminus portion of WT1 (exon 1) and Group B contained peptides present within the carboxy terminus, which contains a four zinc finger region with sequence homology to other DNA-binding proteins. Within group B, p287-301 and p299-313 were derived from exon 7, zinc finger 1, and p421-435 was derived from exon 10, zinc finger IV.

B6 mice were immunized with a group of WT1 peptides or with a control peptide. Peptides were dissolved in 1ml sterile water for injection, and B6 mice were immunized 3 times at time intervals of three weeks. Adjuvants used were CFA/IFA, GM-
25 CSF, and Montinide. The presence of antibodies specific for WT1 was then determined as

described in Examples 1 and 2, and proliferative T cell responses were evaluated using a standard thymidine incorporation assay, in which cells were cultured in the presence of antigen and proliferation was evaluated by measuring incorporated radioactivity (Chen et al., *Cancer Res.* 54:1065-1070, 1994). In particular, lymphocytes were cultured in 96-well
5 plates at 2×10^5 cells per well with 4×10^5 irradiated (3000 rads) syngeneic spleen cells and the designated peptide.

Immunization of mice with the group of peptides designated as Group A elicited an antibody response to WT1 (Figure 4). No antibodies were detected following immunization to Vaccine B, which is consistent with a lack of helper T cell response from
10 immunization with Vaccine B. P117-139 elicited proliferative T cell responses (Figures 5A-5C). The stimulation indices (SI) varied between 8 and 72. Other peptides (P6-22 and P299-313) also were shown to elicit proliferative T cell responses. Immunization with P6-22 resulted in a stimulation index (SI) of 2.3 and immunization with P299-313 resulted in a SI of 3.3. Positive controls included ConA stimulated T cells, as well as T cells stimulated
15 with known antigens, such as CD45 and FBL, and allogeneic T cell lines (DeBruijn et al., *Eur. J. Immunol.* 21:2963-2970, 1991).

Figures 6A and 6B show the proliferative response observed for each of the three peptides within vaccine A (Figure 6A) and vaccine B (Figure 6B). Vaccine A elicited proliferative T cell responses to the immunizing peptides p6-22 and p117-139, with
20 stimulation indices (SI) varying between 3 and 8 (bulk lines). No proliferative response to p244-262 was detected (Figure 6A).

Subsequent *in vitro* stimulations were carried out as single peptide stimulations using only p6-22 and p117-139. Stimulation of the Vaccine A specific T cell line with p117-139 resulted in proliferation to p117-139 with no response to p6-22
25 (Figure 7A). Clones derived from the line were specific for p117-139 (Figure 7B). By contrast, stimulation of the Vaccine A specific T cell line with p6-22 resulted in proliferation to p6-22 with no response to p117-139 (Figure 7C). Clones derived from the line were specific for p6-22 (Figure 7D).

These results show that vaccination with WT1 peptides can elicit antibody responses to WT1 protein and proliferative T cell responses to the immunizing peptides.

5

Example 4

Induction of CTL Responses in Mice Immunized with WT1 Peptides

This Example illustrates the ability of WT1 peptides to elicit CTL immunity.

Peptides (9-mers) with motifs appropriate for binding to class I MHC were identified using a BIMAS HLA peptide binding prediction analysis (Parker et al., *J. Immunol.* 152:163, 1994). Peptides identified within such analyses are shown in Tables II - XLIV. In each of these tables, the score reflects the theoretical binding affinity (half-time of dissociation) of the peptide to the MHC molecule indicated.

Peptides identified using the Tsites program (Rothbard and Taylor, *EMBO J.* 7:93-100, 1988; Deavin et al., *Mol. Immunol.* 33:145-155, 1996), which searches for peptide motifs that have the potential to elicit Th responses are further shown in Figures 8A and 8B, and Table XLV.

20

Table II
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA A1

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	137	CLESQPAIR (SEQ ID NO:47)	18.000
2	80	GAEPHEEQC (SEQ ID NO:87)	9.000
3	40	FAPPGASAY (SEQ ID NO:74)	5.000
4	354	QCDFKDCER (SEQ ID NO:162)	5.000
5	2	GSDVRDLNA (SEQ	3.750

		ID NO:101)	
6	152	VTFDGTPSY (SEQ ID NO:244)	2.500
7	260	WTEGQSNHS (SEQ ID NO:247)	2.250
8	409	TSEKPFSCR (SEQ ID NO:232)	1.350
9	73	KQEPSWGGA (SEQ ID NO:125)	1.350
10	386	KTCQRKFSR (SEQ ID NO:128)	1.250
11	37	VLDFAPPGA (SEQ ID NO:241)	1.000
12	325	CAYPGCNKR (SEQ ID NO:44)	1.000
13	232	QLECMTWNQ (SEQ ID NO:167)	0.900
14	272	ESDNHTTPI (SEQ ID NO:71)	0.750
15	366	RSDQLKRHQ (SEQ ID NO:193)	0.750
16	222	SSDNLYQMT (SEQ ID NO:217)	0.750
17	427	RSDELVRHH (SEQ ID NO:191)	0.750
18	394	RSDHLKTHT (SEQ ID NO:192)	0.750
19	317	TSEKRPFMC (SEQ ID NO:233)	0.675
20	213	QALLLRTPY (SEQ ID NO:160)	0.500

Table III
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA A 0201

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	126	RMFPNAPYL (SEQ	313.968

		ID NO:185)	
2	187	SLGEQQYSV (SEQ ID NO:214)	285.163
3	10	ALLPAVPSL (SEQ ID NO:34)	181.794
4	242	NLGATLKGV (SEQ ID NO:146)	159.970
5	225	NLYQMTSQL (SEQ ID NO:147)	68.360
6	292	GVFRGIQDV (SEQ ID NO:103)	51.790
7	191	QQYSVPPP (SEQ ID NO:171)	22.566
8	280	ILCGAQYRI (SEQ ID NO:116)	17.736
9	235	CMTWNQMN (SEQ ID NO:49)	15.428
10	441	NMTKLQAL (SEQ ID NO:149)	15.428
11	7	DLNALLPAV (SEQ ID NO:58)	11.998
12	227	YQMTSQLE (SEQ ID NO:251)	8.573
13	239	NQMNLGATL (SEQ ID NO:151)	8.014
14	309	TLVRSASET (SEQ ID NO:226)	7.452
15	408	KTSEKPFSC (SEQ ID NO:129)	5.743
16	340	LQMHSRKHT (SEQ ID NO:139)	4.752
17	228	QMTSQLECM (SEQ ID NO:169)	4.044
18	93	TVHFSGQFT (SEQ ID NO:235)	3.586
19	37	VLDFAAPGA (SEQ ID NO:241)	3.378
20	86	EQCLSAFTV (SEQ ID NO:69)	3.068

Table IV
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA A 0205

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	10	ALLPAVPSL (SEQ ID NO:34)	42.000
2	292	GVFRGIQDV (SEQ ID NO:103)	24.000
3	126	RMFPNAPYL (SEQ ID NO:185)	21.000
4	225	NLYQMTSQL (SEQ ID NO:147)	21.000
5	239	NQMNLGATL (SEQ ID NO:151)	16.800
6	302	RVPGVAPTL (SEQ ID NO:195)	14.000
7	441	NMTKLQLAL (SEQ ID NO:149)	7.000
8	235	CMTWNQMNL (SEQ ID NO:49)	7.000
9	187	SLGEQQYSV (SEQ ID NO:214)	6.000
10	191	QQYSVPPPV (SEQ ID NO:171)	4.800
11	340	LQMHSRKHT (SEQ ID NO:139)	4.080
12	242	NLGATLKGV (SEQ ID NO:146)	4.000
13	227	YQMTSQLEC (SEQ ID NO:251)	3.600
14	194	SVPPPVYGC (SEQ ID NO:218)	2.000
15	93	TVHFSGQFT (SEQ ID NO:235)	2.000
16	280	ILCGAQYRI (SEQ ID NO:116)	1.700
17	98	GQFTGTAGA (SEQ ID NO:99)	1.200

18	309	TLVRSASET (SEQ ID NO:226)	1.000
19	81	AEPHEEQCL (SEQ ID NO:30)	0.980
20	73	KQEPSWGGA (SEQ ID NO:125)	0.960

Table V
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA A24

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	302	RVPGVAPTL (SEQ ID NO:195)	16.800
2	218	RTPYSSDNL (SEQ ID NO:194)	12.000
3	356	DFKDCERRF (SEQ ID NO:55)	12.000
4	126	RMFPNAPYL (SEQ ID NO:185)	9.600
5	326	AYPGCNKRY (SEQ ID NO:42)	7.500
6	270	GYESDNHT (SEQ ID NO:106)T	7.500
7	239	NQMNLGATL (SEQ ID NO:151)	7.200
8	10	ALLPAVPSL (SEQ ID NO:34)	7.200
9	130	NAPYLPSCSL (SEQ ID NO:144)	7.200
10	329	GCKNRYFKL (SEQ ID NO:90)	6.600
11	417	RWPSCQKKF (SEQ ID NO:196)	6.600
12	47	AYGSLGGPA (SEQ ID NO:41)	6.000
13	180	DPMGQQGSL (SEQ ID NO:59)	6.000

14	4	DVRDLNALL (SEQ ID NO:62)	5.760
15	285	QYRIHTHGV (SEQ ID NO:175)	5.000
16	192	QYSVPPPVS (SEQ ID NO:176)	5.000
17	207	DSCTGSQAL (SEQ ID NO:61)	4.800
18	441	NMTKLQAL (SEQ ID NO:149)	4.800
19	225	NLYQMTSQL (SEQ ID NO:147)	4.000
20	235	CMTWNQMNL (SEQ ID NO:49)	4.000

Table VI
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA A3

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	436	NMHQRNMTK (SEQ ID NO:148)	40.000
2	240	QMNLGATLK (SEQ ID NO:168)	20.000
3	88	CLSAFTVHF (SEQ ID NO:48)	6.000
4	126	RMFPNAPYL (SEQ ID NO:185)	4.500
5	169	AQFPNHSFK (SEQ ID NO:36)	4.500
6	10	ALLPAVPSL (SEQ ID NO:34)	4.050
7	137	CLESQPAIR (SEQ ID NO:47)	4.000
8	225	NLYQMTSQL (SEQ ID NO:147)	3.000
9	32	AQWAPVLDF (SEQ ID NO:37)	2.700

10	280	ILCGAQYRI (SEQ ID NO:116)	2.700
11	386	KTCQRKFSR (SEQ ID NO:128)	1.800
12	235	CMTWNQMNL (SEQ ID NO:49)	1.200
13	441	NMTKLQLAL (SEQ ID NO:149)	1.200
14	152	VTFDGTPSY (SEQ ID NO:244)	1.000
15	187	SLGEQQYSV (SEQ ID NO:214)	0.900
16	383	FQCKTCQRK (SEQ ID NO:80)	0.600
17	292	GVFRGIQDV (SEQ ID NO:103)	0.450
18	194	SVPPPVYGC (SEQ ID NO:218)	0.405
19	287	RIHTHGVFR (SEQ ID NO:182)	0.400
20	263	GQSNHSTGY (SEQ ID NO:100)	0.360

Table VII
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA A68.1

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	100	FTGTAGACR (SEQ ID NO:84)	100.000
2	386	KTCQRKFSR (SEQ ID NO:128)	50.000
3	368	DQLKRHQRR (SEQ ID NO:60)	30.000
4	312	RSASETSEK (SEQ ID NO:190)	18.000
5	337	LSHLQMHSR (SEQ ID NO:141)	15.000

6	364	FSRSDQLKR (SEQ ID NO:83)	15.000
7	409	TSEKPFSCR (SEQ ID NO:232)	15.000
8	299	DVRRVPGVA (SEQ ID NO:63)	12.000
9	4	DVRDLNALL (SEQ ID NO:62)	12.000
10	118	SQASSGQAR (SEQ ID NO:216)	10.000
11	343	HSRKHTGEK (SEQ ID NO:111)	9.000
12	169	AQFPNHSFK (SEQ ID NO:36)	9.000
13	292	GVFRGIQDV (SEQ ID NO:103)	8.000
14	325	CAYPGCNKR (SEQ ID NO:44)	7.500
15	425	FARSDLVLR (SEQ ID NO:75)	7.500
16	354	QCDFKDCER (SEQ ID NO:162)	7.500
17	324	MCAYPGCNK (SEQ ID NO:142)	6.000
18	251	AAGSSSSVK (SEQ ID NO:28)	6.000
19	379	GVKPFQCKT (SEQ ID NO:104)	6.000
20	137	CLESQPAIR (SEQ ID NO:47)	5.000

Table VIII
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA A 1101

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	386	KTCQRKFSR (SEQ ID NO:128)	1.800

2	169	AQFPNHSFK (SEQ ID NO:36)	1.200
3	436	NMHQRNMTK (SEQ ID NO:148)	0.800
4	391	KFSRSDHLK (SEQ ID NO:120)	0.600
5	373	HQRRHTGVK (SEQ ID NO:109)	0.600
6	383	FQCKTCQRK (SEQ ID NO:80)	0.600
7	363	RFSRSDQLK (SEQ ID NO:178)	0.600
8	240	QMNLGATLK (SEQ ID NO:168)	0.400
9	287	RIHTHGVFR (SEQ ID NO:182)	0.240
10	100	FTGTAGACR (SEQ ID NO:84)	0.200
11	324	MCAYPGCNK (SEQ ID NO:142)	0.200
12	251	AAGSSSSVK (SEQ ID NO:28)	0.200
13	415	SCRWPSCQK (SEQ ID NO:201)	0.200
14	118	SQASSGQAR (SEQ ID NO:216)	0.120
15	292	GVFRGIQDV (SEQ ID NO:103)	0.120
16	137	CLESQPAIR (SEQ ID NO:47)	0.080
17	425	FARSDLVLR (SEQ ID NO:75)	0.080
18	325	CAYPGCNKR (SEQ ID NO:44)	0.080
19	312	RSASETSEK (SEQ ID NO:190)	0.060
20	65	PPPPHSFI (SEQ ID NO:156)K	0.060

Table IX
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA A 3101

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	386	KTCQRKFSR (SEQ ID NO:128)	9.000
2	287	RIHTHGVFR (SEQ ID NO:182)	6.000
3	137	CLESQPAIR (SEQ ID NO:47)	2.000
4	118	SQASSGQAR (SEQ ID NO:216)	2.000
5	368	DQLKRHQRR (SEQ ID NO:60)	1.200
6	100	FTGTAGACR (SEQ ID NO:84)	1.000
7	293	VFRGIQDVR (SEQ ID NO:238)	0.600
8	325	CAYPGCNKR (SEQ ID NO:44)	0.600
9	169	AQFPNHSFK (SEQ ID NO:36)	0.600
10	279	PILCGAQYR (SEQ ID NO:155)	0.400
11	436	NMHQRNMTK (SEQ ID NO:148)	0.400
12	425	FARSDLVLR (SEQ ID NO:75)	0.400
13	32	AQWAPVLDF (SEQ ID NO:37)	0.240
14	240	QMNLGATLK (SEQ ID NO:168)	0.200
15	354	QCDFKDCER (SEQ ID NO:162)	0.200
16	373	HQRRHTGVK (SEQ ID NO:109)	0.200
17	383	FQCKTCQRK (SEQ ID NO:80)	0.200

18	313	SASETSEKR (SEQ ID NO:197)	0.200
19	358	KDCERRFSR (SEQ ID NO:118)	0.180
20	391	KFSRSDHLK (SEQ ID NO:120)	0.180

Table X
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA A 3302

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	337	LSHLQMHSR (SEQ ID NO:141)	15.000
2	409	TSEKPFSCR (SEQ ID NO:232)	15.000
3	364	FSRSDQLKR (SEQ ID NO:83)	15.000
4	137	CLESQPAIR (SEQ ID NO:47)	9.000
5	368	DQLKRHQRR (SEQ ID NO:60)	9.000
6	287	RIHTHGVFR (SEQ ID NO:182)	4.500
7	210	TGSQALLLR (SEQ ID NO:223)	3.000
8	425	FARSDLVLR (SEQ ID NO:75)	3.000
9	313	SASETSEKR (SEQ ID NO:197)	3.000
10	293	VFRGIQDVR (SEQ ID NO:238)	3.000
11	354	QCDFKDCER (SEQ ID NO:162)	3.000
12	100	FTGTAGACR (SEQ ID NO:84)	3.000
13	118	SQASSGQAR (SEQ ID NO:216)	3.000

14	325	CAYPGCNKR (SEQ ID NO:44)	3.000
15	207	DSCTGSQAL (SEQ ID NO:61)	1.500
16	139	ESQPAIRNQ (SEQ ID NO:72)	1.500
17	299	DVRRVPGVA (SEQ ID NO:63)	1.500
18	419	PSCQKKFAR (SEQ ID NO:159)	1.500
19	272	ESDNHTTPI (SEQ ID NO:71)	1.500
20	4	DVRDLNALL (SEQ ID NO:62)	1.500

Table XI
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B14

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	362	RRFSRSDQL (SEQ ID NO:187)	1000.000
2	332	KRYFKLSHL (SEQ ID NO:127)	300.000
3	423	KKFARSDDEL (SEQ ID NO:122)	150.000
4	390	RKFSRSDHL (SEQ ID NO:183)	150.000
5	439	QRNMTKLQL (SEQ ID NO:173)	20.000
6	329	GCNKRYFKL (SEQ ID NO:90)	10.000
7	10	ALLPAVPSL (SEQ ID NO:34)	10.000
8	180	DPMGQQGSL (SEQ ID NO:59)	9.000
9	301	RRVPGVAPT (SEQ ID NO:189)	6.000

10	126	RMFPNAPYL (SEQ ID NO:185)	5.000
11	371	KRHQRRHTG (SEQ ID NO:126)	5.000
12	225	NLYQMTSQL (SEQ ID NO:147)	5.000
13	144	IRNQGYSTV (SEQ ID NO:117)	4.000
14	429	DELVRHHNM (SEQ ID NO:53)	3.000
15	437	MHQRNMTKL (SEQ ID NO:143)	3.000
16	125	ARMFPNAPY (SEQ ID NO:38)	3.000
17	239	NQMNLGATL (SEQ ID NO:151)	3.000
18	286	YRIHTHGVF (SEQ ID NO:252)	3.000
19	174	HSFKHEDPM (SEQ ID NO:110)	3.000
20	372	RHQRRHTGV (SEQ ID NO:181)	3.000

Table XII
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B40

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	81	AEPHEEQCL (SEQ ID NO:30)	40.000
2	429	DELVRHHNM (SEQ ID NO:53)	24.000
3	410	SEKPFSCRW (SEQ ID NO:207)	20.000
4	318	SEKRPFMCA (SEQ ID NO:208)	15.000
5	233	LECMTWNQM (SEQ ID NO:131)	12.000

6	3	SDVRDLNAL (SEQ ID NO:206)	10.000
7	349	GEKPYQCDF (SEQ ID NO:91)	8.000
8	6	RDLNALLPA (SEQ ID NO:177)	5.000
9	85	EEQCLSAFT (SEQ ID NO:65)	4.000
10	315	SETSEKRPF (SEQ ID NO:209)	4.000
11	261	TEGQSNHST (SEQ ID NO:221)	4.000
12	23	GCALPVSGA (SEQ ID NO:89)	3.000
13	38	LDFAPPGAS (SEQ ID NO:130)	3.000
14	273	SDNHTTPIL (SEQ ID NO:204)	2.500
15	206	TDSCTGSQA (SEQ ID NO:220)	2.500
16	24	CALPVSGAA (SEQ ID NO:43)	2.000
17	98	GQFTGTAGA (SEQ ID NO:99)	2.000
18	30	GAAQWAPVL (SEQ ID NO:86)	2.000
19	84	HEEQCLSAF (SEQ ID NO:107)	2.000
20	26	LPVSGAAQW (SEQ ID NO:138)	2.000

Table XIII
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B60

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	81	AEPHEEQCL (SEQ ID NO:30)	160.000

2	3	SDVRDLNAL (SEQ ID NO:206)	40.000
3	429	DELVRHHNM (SEQ ID NO:53)	40.000
4	233	LECMTWNQM (SEQ ID NO:131)	22.000
5	273	SDNHTTPIL (SEQ ID NO:204)	20.000
6	209	CTGSQALLL (SEQ ID NO:52)	8.000
7	30	GAAQWAPVL (SEQ ID NO:86)	8.000
8	318	SEKRPFMCA (SEQ ID NO:208)	8.000
9	180	DPMGQQGSL (SEQ ID NO:59)	8.000
10	138	LESQPAIRN (SEQ ID NO:132)	5.280
11	239	NQMNLGATL (SEQ ID NO:151)	4.400
12	329	GCNKRYFKL (SEQ ID NO:90)	4.400
13	130	NAPYLPSC (SEQ ID NO:144)	4.400
14	85	EEQCLSAFT (SEQ ID NO:65)	4.400
15	208	SCTGSQALL (SEQ ID NO:202)	4.000
16	207	DSC TGSQAL (SEQ ID NO:61)	4.000
17	218	RTPYSSDNL (SEQ ID NO:194)	4.000
18	261	TEGQSNHST (SEQ ID NO:221)	4.000
19	18	LGGGGGCAL (SEQ ID NO:134)	4.000
20	221	YSSDNL YQM (SEQ ID NO:253)	2.200

Table XIV
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B61

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	318	SEKRPFMCA (SEQ ID NO:208)	20.000
2	429	DELVRHHNM (SEQ ID NO:53)	16.000
3	298	QDVRRVPGV (SEQ ID NO:164)	10.000
4	81	AEPHEEQCL (SEQ ID NO:30)	8.000
5	233	LECMTWNQM (SEQ ID NO:131)	8.000
6	6	RDLNALLPA (SEQ ID NO:177)	5.500
7	85	EEQCLSAFT (SEQ ID NO:65)	4.000
8	261	TEGQSNHST (SEQ ID NO:221)	4.000
9	206	TDSCGTGSQA (SEQ ID NO:220)	2.500
10	295	RGIQDVRRV (SEQ ID NO:179)	2.200
11	3	SDVRDLNAL (SEQ ID NO:206)	2.000
12	250	VAAGSSSSV (SEQ ID NO:236)	2.000
13	29	SGAAQWAPV (SEQ ID NO:211)	2.000
14	315	SETSEKRPF (SEQ ID NO:209)	1.600
15	138	LESQPAIRN (SEQ ID NO:132)	1.200
16	244	GATLKGVAA (SEQ ID NO:88)	1.100
17	20	GGGGCALPV (SEQ ID NO:92)	1.100

18	440	RNMTKLQLA (SEQ ID NO:186)	1.100
19	23	GCALPVSGA (SEQ ID NO:89)	1.100
20	191	QQYSVPPPV (SEQ ID NO:171)	1.000

Table XV
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B62

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	146	NQGYSTVTF (SEQ ID NO:150)	211.200
2	32	AQWAPVLDF (SEQ ID NO:37)	96.000
3	263	GQSNHSTGY (SEQ ID NO:100)	96.000
4	88	CLSAFTVHF (SEQ ID NO:48)	96.000
5	17	SLGGGGGCA (SEQ ID NO:215)	9.600
6	239	NQMNLGATL (SEQ ID NO:151)	8.800
7	191	QQYSVPPPV (SEQ ID NO:171)	8.000
8	98	GQFTGTAGA (SEQ ID NO:99)	8.000
9	384	QCKTCQRKF (SEQ ID NO:163)	6.000
10	40	FAPPGASAY (SEQ ID NO:74)	4.800
11	227	YQMTSQLEC (SEQ ID NO:251)	4.800
12	187	SLGEQQYSV (SEQ ID NO:214)	4.400
13	86	EQCLSAFTV (SEQ ID NO:69)	4.400

14	152	VTFDGTPSY (SEQ ID NO:244)	4.400
15	101	TGTAGACRY (SEQ ID NO:224)	4.000
16	242	NLGATLKGV (SEQ ID NO:146)	4.000
17	92	FTVHFSGQF (SEQ ID NO:85)	4.000
18	7	DLNALLPAV (SEQ ID NO:58)	4.000
19	123	GQARMFPNA (SEQ ID NO:98)	4.000
20	280	ILCGAQYRI (SEQ ID NO:116)	3.120

Table XVI
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B7

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	180	DPMGQQGSL (SEQ ID NO:59)	240.000
2	4	DVRDLNALL (SEQ ID NO:62)	200.000
3	302	RVPGVAPTL (SEQ ID NO:195)	20.000
4	30	GAAQWAPVL (SEQ ID NO:86)	12.000
5	239	NQMNLGATL (SEQ ID NO:151)	12.000
6	130	NAPYLPSCL (SEQ ID NO:144)	12.000
7	10	ALLPAVPSL (SEQ ID NO:34)	12.000
8	299	DVRRVPGVA (SEQ ID NO:63)	5.000
9	208	SCTGSQALL (SEQ ID NO:202)	4.000

10	303	VPGVAPTLV (SEQ ID NO:242)	4.000
11	18	LGGGGGCAL (SEQ ID NO:134)	4.000
12	218	RTPYSSDNL (SEQ ID NO:194)	4.000
13	207	DSCTGSQAL (SEQ ID NO:61)	4.000
14	209	CTGSQALLL (SEQ ID NO:52)	4.000
15	329	GCNKRYFKL (SEQ ID NO:90)	4.000
16	235	CMTWNQMNL (SEQ ID NO:49)	4.000
17	441	NMTKLQLAL (SEQ ID NO:149)	4.000
18	126	RMFPNAPYL (SEQ ID NO:185)	4.000
19	225	NLYQMTSQL (SEQ ID NO:147)	4.000
20	143	AIRNQGYST (SEQ ID NO:33)	3.000

Table XVII
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B8

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	329	GCNKRYFKL (SEQ ID NO:90)	16.000
2	4	DVRDLNALL (SEQ ID NO:62)	12.000
3	316	ETSEKRPFM (SEQ ID NO:73)	3.000
4	180	DPMGQQGSL (SEQ ID NO:59)	1.600
5	208	SCTGSQALL (SEQ ID NO:202)	0.800

6	130	NAPYLPSCS (SEQ ID NO:144)	0.800
7	244	GATLKGVA (SEQ ID NO:88)	0.800
8	30	GAAQWAPVL (SEQ ID NO:86)	0.800
9	299	DVRRVPGVA (SEQ ID NO:63)	0.400
10	420	SCQKKFARS (SEQ ID NO:200)	0.400
11	387	TCQRKFSRS (SEQ ID NO:219)	0.400
12	225	NLYQMTSQL (SEQ ID NO:147)	0.400
13	141	QPAIRNQGY (SEQ ID NO:170)	0.400
14	10	ALLPAVPSL (SEQ ID NO:34)	0.400
15	207	DSCTGSQAL (SEQ ID NO:61)	0.400
16	384	QCKTCQRKF (SEQ ID NO:163)	0.400
17	136	SCLESQPAI (SEQ ID NO:198)	0.300
18	347	HTGEKPYQC (SEQ ID NO:112)	0.300
19	401	HTRTHTGKT (SEQ ID NO:114)	0.200
20	332	KRYFKLSHL (SEQ ID NO:127)	0.200

Table XVIII
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B 2702

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	332	KRYFKLSHL (SEQ ID NO:127)	900.000

2	362	RRFSRSDQL (SEQ ID NO:187)	900.000
3	286	YRIHTHGVF (SEQ ID NO:252)	200.000
4	125	ARMFPNAPY (SEQ ID NO:38)	200.000
5	375	RRHTGVKPF (SEQ ID NO:188)	180.000
6	32	AQWAPVLDF (SEQ ID NO:37)	100.000
7	301	RRVPGVAPT (SEQ ID NO:189)	60.000
8	439	QRNMTKLQL (SEQ ID NO:173)	60.000
9	126	RMFPNAPYL (SEQ ID NO:185)	22.500
10	426	ARSDLVVRH (SEQ ID NO:39)	20.000
11	146	NQGYSTVTF (SEQ ID NO:150)	20.000
12	144	IRNQGYSTV (SEQ ID NO:117)	20.000
13	389	QRKFSRSDH (SEQ ID NO:172)	20.000
14	263	GQSNHSTGY (SEQ ID NO:100)	20.000
15	416	CRWPSCQKK (SEQ ID NO:50)	20.000
16	191	QQYSVPPP (SEQ ID NO:171)	10.000
17	217	LRTPYSSDN (SEQ ID NO:140)	10.000
18	107	CRYGPFGRP (SEQ ID NO:51)	10.000
19	98	GQFTGTAGA (SEQ ID NO:99)	10.000
20	239	NQMNLGATL (SEQ ID NO:151)	6.000

Table XIX
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B 2705

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	332	KRYFKLSHL (SEQ ID NO:127)	30000.000
2	362	RRFSRSDQL (SEQ ID NO:187)	30000.000
3	416	CRWPSCQKK (SEQ ID NO:50)	10000.000
4	439	QRNMTKLQL (SEQ ID NO:173)	2000.000
5	286	YRIHTHGVF (SEQ ID NO:252)	1000.000
6	125	ARMFPNAPY (SEQ ID NO:38)	1000.000
7	294	FRGIQDVRR (SEQ ID NO:81)	1000.000
8	432	VRHHNMHQR (SEQ ID NO:243)	1000.000
9	169	AQFPNHSFK (SEQ ID NO:36)	1000.000
10	375	RRHTGVKPF (SEQ ID NO:188)	900.000
11	126	RMFPNAPYL (SEQ ID NO:185)	750.000
12	144	IRNQGYSTV (SEQ ID NO:117)	600.000
13	301	RRVPGVAPT (SEQ ID NO:189)	600.000
14	32	AQWAPVLDF (SEQ ID NO:37)	500.000
15	191	QQYSVPPPV (SEQ ID NO:171)	300.000
16	373	HQRRHTGVK (SEQ ID NO:109)	200.000
17	426	ARSDELVRH (SEQ ID NO:39)	200.000

18	383	FQCKTCQRK (SEQ ID NO:80)	200.000
19	239	NQMNLGATL (SEQ ID NO:151)	200.000
20	389	QRKFSRSDH (SEQ ID NO:172)	200.000

Table XX
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B 3501

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	278	TPILCGAQY (SEQ ID NO:227)	40.000
2	141	QPAIRNQGY (SEQ ID NO:170)	40.000
3	219	TPYSSDNLY (SEQ ID NO:231)	40.000
4	327	YPGCNKRYF (SEQ ID NO:250)	20.000
5	163	TPSHHAAQF (SEQ ID NO:228)	20.000
6	180	DPMGQQGSL (SEQ ID NO:59)	20.000
7	221	YSSDNLYQM (SEQ ID NO:253)	20.000
8	26	LPVSGAAQW (SEQ ID NO:138)	10.000
9	174	HSFKHEDPM (SEQ ID NO:110)	10.000
10	82	EPHEEQCLS (SEQ ID NO:68)	6.000
11	213	QALLLRTPY (SEQ ID NO:160)	6.000
12	119	QASSGQARM (SEQ ID NO:161)	6.000
13	4	DVRDLNALL (SEQ ID NO:62)	6.000

14	40	FAPPGASAY (SEQ ID NO:74)	6.000
15	120	ASSGQARMF (SEQ ID NO:40)	5.000
16	207	DSCTGSQAL (SEQ ID NO:61)	5.000
17	303	VPGVAPTLV (SEQ ID NO:242)	4.000
18	316	ETSEKRPFM (SEQ ID NO:73)	4.000
19	152	VTFDGTPSY (SEQ ID NO:244)	4.000
20	412	KPFSCRWPS (SEQ ID NO:123)	4.000

Table XXI
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B 3701

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	3	SDVRDLNAL (SEQ ID NO:206)	40.000
2	273	SDNHTTPIL (SEQ ID NO:204)	40.000
3	81	AEPHEEQCL (SEQ ID NO:30)	10.000
4	298	QDVRRVPGV (SEQ ID NO:164)	8.000
5	428	SDELVRHHN (SEQ ID NO:203)	6.000
6	85	EEQCLSAFT (SEQ ID NO:65)	5.000
7	208	SCTGSQALL (SEQ ID NO:202)	5.000
8	4	DVRDLNALL (SEQ ID NO:62)	5.000
9	209	CTGSQALLL (SEQ ID NO:52)	5.000

10	38	LDFAPPGAS (SEQ ID NO:130)	4.000
11	223	SDNLYQMTS (SEQ ID NO:205)	4.000
12	179	EDPMGQQGS (SEQ ID NO:64)	4.000
13	206	TDSCTGSQA (SEQ ID NO:220)	4.000
14	6	RDLNALLPA (SEQ ID NO:177)	4.000
15	84	HEEQCLSAF (SEQ ID NO:107)	2.000
16	233	LECMTWNQM (SEQ ID NO:131)	2.000
17	429	DELVRHHNM (SEQ ID NO:53)	2.000
18	315	SETSEKRPF (SEQ ID NO:209)	2.000
19	349	GEKPYQCDF (SEQ ID NO:91)	2.000
20	302	RVPGVAPTL (SEQ ID NO:195)	1.500

Table XXII
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B 3801

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	437	MHQRNMTKL (SEQ ID NO:143)	36.000
2	434	HHNMHQ RN M (SEQ ID NO:108)	6.000
3	372	RHQRRHTGV (SEQ ID NO:181)	6.000
4	180	DPMGQQGSL (SEQ ID NO:59)	4.000
5	433	RHHNMHQ RN (SEQ ID NO:180)	3.900

6	165	SHHAAQFPN (SEQ ID NO:213)	3.900
7	202	CHTPDSCCT (SEQ ID NO:45)	3.000
8	396	DHLKTHTRT (SEQ ID NO:57)	3.000
9	161	GHTPSHHAA (SEQ ID NO:94)	3.000
10	302	RVPGVAPTL (SEQ ID NO:195)	2.600
11	417	RWPSCQKKF (SEQ ID NO:196)	2.400
12	327	YPGCNKRYF (SEQ ID NO:250)	2.400
13	208	SCTGSQALL (SEQ ID NO:202)	2.000
14	163	TPSHHAAQF (SEQ ID NO:228)	2.000
15	120	ASSGQARMF (SEQ ID NO:40)	2.000
16	18	LGGGGGCAL (SEQ ID NO:134)	2.000
17	177	KHEDPMGQQ (SEQ ID NO:121)	1.800
18	83	PHEEQCLSA (SEQ ID NO:154)	1.800
19	10	ALLPAVPSL (SEQ ID NO:34)	1.300
20	225	NLYQMTSQL (SEQ ID NO:147)	1.300

Table XXIII
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B 3901

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	437	MHQRNMTKL (SEQ ID NO:143)	135.000

2	332	KRYFKLSHL (SEQ ID NO:127)	45.000
3	434	HHNMHQQRNM (SEQ ID NO:108)	30.000
4	362	RRFSRSDQL (SEQ ID NO:187)	30.000
5	372	RHQRRHTGV (SEQ ID NO:181)	30.000
6	10	ALLPAVPSL (SEQ ID NO:34)	9.000
7	439	QRNMTKLQL (SEQ ID NO:173)	7.500
8	390	RKFSRSDHL (SEQ ID NO:183)	6.000
9	396	DHLKTHTRT (SEQ ID NO:57)	6.000
10	239	NQMNLGATL (SEQ ID NO:151)	6.000
11	423	KKFARSDDEL (SEQ ID NO:122)	6.000
12	126	RMFPNAPYL (SEQ ID NO:185)	6.000
13	225	NLYQMTSQL (SEQ ID NO:147)	6.000
14	180	DPMGQQGSL (SEQ ID NO:59)	6.000
15	144	IRNQGYSTV (SEQ ID NO:117)	5.000
16	136	SCLESQPAI (SEQ ID NO:198)	4.000
17	292	GVFRGIQDV (SEQ ID NO:103)	3.000
18	302	RVPGVAPTL (SEQ ID NO:195)	3.000
19	208	SCTGSQALL (SEQ ID NO:202)	3.000
20	207	DSCTGSQAL (SEQ ID NO:61)	3.000

Table XXIV
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B 3902

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	239	NQMNLGATL (SEQ ID NO:151)	24.000
2	390	RKFSRSDHL (SEQ ID NO:183)	20.000
3	423	KKFARSDEL (SEQ ID NO:122)	20.000
4	32	AQWAPVLDF (SEQ ID NO:37)	5.000
5	146	NQGYSTVTF (SEQ ID NO:150)	5.000
6	130	NAPYLPSCCL (SEQ ID NO:144)	2.400
7	225	NLYQMTSQL (SEQ ID NO:147)	2.400
8	30	GAAQWAPVL (SEQ ID NO:86)	2.400
9	441	NMTKLQLAL (SEQ ID NO:149)	2.400
10	302	RVPGVAPTL (SEQ ID NO:195)	2.400
11	126	RMFPNAPYL (SEQ ID NO:185)	2.000
12	218	RTPYSSDNL (SEQ ID NO:194)	2.000
13	209	CTGSQALLL (SEQ ID NO:52)	2.000
14	332	KRYFKLSHL (SEQ ID NO:127)	2.000
15	180	DPMGQQGSL (SEQ ID NO:59)	2.000
16	437	MHQRNMTKL (SEQ ID NO:143)	2.000
17	207	DSCTGSQAL (SEQ ID NO:61)	2.000

18	208	SCTGSQALL (SEQ ID NO:202)	2.000
19	329	GCKNRYFKL (SEQ ID NO:90)	2.000
20	10	ALLPAVPSL (SEQ ID NO:34)	2.000

Table XXV
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B 4403

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	315	SETSEKRPF (SEQ ID NO:209)	80.000
2	349	GEKPYQCDF (SEQ ID NO:91)	80.000
3	84	HEEQCLSAF (SEQ ID NO:107)	60.000
4	410	SEKPFSCRW (SEQ ID NO:207)	48.000
5	429	DELVRHHNM (SEQ ID NO:53)	24.000
6	278	TPILCGAQY (SEQ ID NO:227)	15.000
7	141	QPAIRNQGY (SEQ ID NO:170)	9.000
8	40	FAPPGASAY (SEQ ID NO:74)	9.000
9	213	QALLLRTPY (SEQ ID NO:160)	9.000
10	318	SEKRPFMCA (SEQ ID NO:208)	8.000
11	81	AEPHEEQCL (SEQ ID NO:30)	8.000
12	152	VTFDGTPSY (SEQ ID NO:244)	4.500
13	101	TGTAGACRY (SEQ ID NO:224)	4.500

14	120	ASSGQARMF (SEQ ID NO:40)	4.500
15	261	TEGQSNHST (SEQ ID NO:221)	4.000
16	85	EEQCLSAFT (SEQ ID NO:65)	4.000
17	233	LECMTWNQM (SEQ ID NO:131)	4.000
18	104	AGACRYGPF (SEQ ID NO:31)	4.000
19	3	SDVRDLNAL (SEQ ID NO:206)	3.000
20	185	QGSLGEQQY (SEQ ID NO:166)	3.000

Table XXVI
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B 5101

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	303	VPGVAPTLV (SEQ ID NO:242)	314.600
2	180	DPMGQQGSL (SEQ ID NO:59)	242.000
3	250	VAAGSSSSV (SEQ ID NO:236)	157.300
4	130	NAPYLPSC (SEQ ID NO:144)	50.000
5	30	GAAQWAPVL (SEQ ID NO:86)	50.000
6	20	GGGGCALPV (SEQ ID NO:92)	44.000
7	64	PPPPHSFI (SEQ ID NO:157)	40.000
8	29	SGAAQWAPV (SEQ ID NO:211)	40.000
9	18	LGGGGGCGAL (SEQ ID NO:134)	31.460

10	295	RGIQDVRRV (SEQ ID NO:179)	22.000
11	119	QASSGQARM (SEQ ID NO:161)	18.150
12	418	WPSCQKKFA (SEQ ID NO:246)	12.100
13	82	EPHEEQCLS (SEQ ID NO:68)	12.100
14	110	GPFGPPPPS (SEQ ID NO:96)	11.000
15	272	ESDNHTTPI (SEQ ID NO:71)	8.000
16	306	VAPTLVRS (SEQ ID NO:237)	7.150
17	280	ILCGAQYRI (SEQ ID NO:116)	6.921
18	219	TPYSSDNLY (SEQ ID NO:231)	6.600
19	128	FPNAPYLPS (SEQ ID NO:79)	6.500
20	204	TPTDSCTGS (SEQ ID NO:230)	6.050

Table XXVII
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B 5102

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	295	RGIQDVRRV (SEQ ID NO:179)	290.400
2	303	VPGVAPTLV (SEQ ID NO:242)	200.000
3	180	DPMGQQGSL (SEQ ID NO:59)	133.100
4	250	VAAGSSSSV (SEQ ID NO:236)	110.000
5	30	GAAQWAPVL (SEQ ID NO:86)	55.000

6	130	NAPYLPSCCL (SEQ ID NO:144)	50.000
7	20	GGGGCALPV (SEQ ID NO:92)	44.000
8	29	SGAAQWAPV (SEQ ID NO:211)	44.000
9	64	PPPPHSFI (SEQ ID NO:157)	40.000
10	119	QASSGQARM (SEQ ID NO:161)	36.300
11	110	GPFGPPPPS (SEQ ID NO:96)	27.500
12	412	KPFSCRWPS (SEQ ID NO:123)	25.000
13	18	LGGGGGCAL (SEQ ID NO:134)	24.200
14	24	CALPVSGAA (SEQ ID NO:43)	16.500
15	219	TPYSSDNLY (SEQ ID NO:231)	15.000
16	292	GVFRGIQDV (SEQ ID NO:103)	14.641
17	136	SCLESQPAI (SEQ ID NO:198)	14.520
18	418	WPSCQKKFA (SEQ ID NO:246)	12.100
19	269	TGYESDNHT (SEQ ID NO:225)	11.000
20	351	KPYQCDFKD (SEQ ID NO:124)	11.000

Table XXVIII
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B 5201

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	191	QQYSVPPPV (SEQ ID NO:171)	100.000

2	32	AQWAPVLDF (SEQ ID NO:37)	30.000
3	243	LGATLKGVA (SEQ ID NO:133)	16.500
4	303	VPGVAPTLV (SEQ ID NO:242)	13.500
5	86	EQCLSAFTV (SEQ ID NO:69)	12.000
6	295	RGIQDVRRV (SEQ ID NO:179)	10.000
7	98	GQFTGTAGA (SEQ ID NO:99)	8.250
8	292	GVFRGIQDV (SEQ ID NO:103)	8.250
9	29	SGAAQWAPV (SEQ ID NO:211)	6.000
10	146	NQGYSTVTF (SEQ ID NO:150)	5.500
11	20	GGGGCALPV (SEQ ID NO:92)	5.000
12	239	NQMNLGATL (SEQ ID NO:151)	4.000
13	64	PPPPHSFI (SEQ ID NO:157)	3.600
14	273	SDNHTTPIL (SEQ ID NO:204)	3.300
15	286	YRIHTHGVF (SEQ ID NO:252)	3.000
16	269	TGYESDNHT (SEQ ID NO:225)	3.000
17	406	TGKTSEKPF (SEQ ID NO:222)	2.750
18	327	YPGCNKRYF (SEQ ID NO:250)	2.750
19	7	DLNALLPAV (SEQ ID NO:58)	2.640
20	104	AGACRYGPF (SEQ ID NO:31)	2.500

Table XXIX
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA B 5801

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	230	TSQLECMTW (SEQ ID NO:234)	96.800
2	92	FTVHFSGQF (SEQ ID NO:85)	60.000
3	120	ASSGQARMF (SEQ ID NO:40)	40.000
4	168	AAQFPNHSF (SEQ ID NO:29)	20.000
5	408	KTSEKPFSC (SEQ ID NO:129)	12.000
6	394	RSDHLKTHT (SEQ ID NO:192)	9.900
7	276	HTTPILCGA (SEQ ID NO:115)	7.200
8	218	RTPYSSDNL (SEQ ID NO:194)	6.600
9	152	VTFDGTPSY (SEQ ID NO:244)	6.000
10	40	FAPPGASAY (SEQ ID NO:74)	6.000
11	213	QALLLRTPY (SEQ ID NO:160)	4.500
12	347	HTGEKPYQC (SEQ ID NO:112)	4.400
13	252	AGSSSSVKW (SEQ ID NO:32)	4.400
14	211	GSQALLLRT (SEQ ID NO:102)	4.356
15	174	HSFKHEDPM (SEQ ID NO:110)	4.000
16	317	TSEKRPFMC (SEQ ID NO:233)	4.000
17	26	LPVSGAAQW (SEQ ID NO:138)	4.000

18	289	HTHGVFRGI (SEQ ID NO:113)	3.600
19	222	SSDNLYQMT (SEQ ID NO:217)	3.300
20	96	FSGQFTGTA (SEQ ID NO:82)	3.300

Table XXX
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA CW0301

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	10	ALLPAVPSL (SEQ ID NO:34)	100.000
2	332	KRYFKLSHL (SEQ ID NO:127)	48.000
3	126	RMFPNAPYL (SEQ ID NO:185)	36.000
4	3	SDVRDLNAL (SEQ ID NO:206)	30.000
5	239	NQMNLGATL (SEQ ID NO:151)	24.000
6	225	NLYQMTSQL (SEQ ID NO:147)	24.000
7	180	DPMGQQGSL (SEQ ID NO:59)	20.000
8	362	RRFSRSDQL (SEQ ID NO:187)	12.000
9	329	GCKRYFKL (SEQ ID NO:90)	10.000
10	286	YRIHTHGVF (SEQ ID NO:252)	10.000
11	301	RRVPGVAPT (SEQ ID NO:189)	10.000
12	24	CALPVSGAA (SEQ ID NO:43)	10.000
13	136	SCLESQPAI (SEQ ID NO:198)	7.500
14	437	MHQRNMTKL (SEQ	7.200

		ID NO:143)	
15	390	RKFSRSDHL (SEQ ID NO:183)	6.000
16	423	KKFARSDEL (SEQ ID NO:122)	6.000
17	92	FTVHFSGQF (SEQ ID NO:85)	5.000
18	429	DELVRHHNM (SEQ ID NO:53)	5.000
19	130	NAPYLPSC (SEQ ID NO:144)	4.800
20	30	GAAQWAPVL (SEQ ID NO:86)	4.000

Table XXXI

Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA CW0401

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	356	DFKDCERRF (SEQ ID NO:55)	120.000
2	334	YFKLSHLQM (SEQ ID NO:248)	100.000
3	180	DPMGQQGSL (SEQ ID NO:59)	88.000
4	163	TPSHHAAQF (SEQ ID NO:228)	52.800
5	327	YPGCNKRYF (SEQ ID NO:250)	40.000
6	285	QYRIHTHGV (SEQ ID NO:175)	27.500
7	424	KFARSDELV (SEQ ID NO:119)	25.000
8	326	AYPGCNKRY (SEQ ID NO:42)	25.000
9	192	QYSVPPPVY (SEQ ID NO:176)	25.000
10	417	RWPSCQKKF (SEQ ID NO:196)	22.000

11	278	TPILCGAQY (SEQ ID NO:227)	12.000
12	10	ALLPAVPSL (SEQ ID NO:34)	11.616
13	141	QPAIRNQGY (SEQ ID NO:170)	11.000
14	303	VPGVAPTLV (SEQ ID NO:242)	11.000
15	219	TPYSSDNLY (SEQ ID NO:231)	10.000
16	39	DFAPPGASA (SEQ ID NO:54)	7.920
17	99	QFTGTAGAC (SEQ ID NO:165)	6.000
18	4	DVRDLNALL (SEQ ID NO:62)	5.760
19	70	SFIKQEPSW (SEQ ID NO:210)	5.500
20	63	PPPPPHSF (SEQ ID NO:158)	5.280

Table XXXII

Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA CW0602

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	332	KRYFKLSHL (SEQ ID NO:127)	9.680
2	239	NQMNLGATL (SEQ ID NO:151)	6.600
3	130	NAPYLPSCS (SEQ ID NO:144)	6.600
4	7	DLNALLPAV (SEQ ID NO:58)	6.000
5	441	NMTKLQLAL (SEQ ID NO:149)	6.000
6	225	NLYQMTSQL (SEQ ID NO:147)	6.000

7	4	DVRDLNALL (SEQ ID NO:62)	6.000
8	3	SDVRDLNAL (SEQ ID NO:206)	4.400
9	10	ALLPAVPSL (SEQ ID NO:34)	4.000
10	213	QALLLRTPY (SEQ ID NO:160)	3.300
11	319	EKRPFMCAY (SEQ ID NO:67)	3.000
12	30	GAAQWAPVL (SEQ ID NO:86)	2.200
13	242	NLGATLKGV (SEQ ID NO:146)	2.200
14	292	GVFRGIQDV (SEQ ID NO:103)	2.200
15	207	DSCTGSQAL (SEQ ID NO:61)	2.200
16	362	RRFSRSDQL (SEQ ID NO:187)	2.200
17	439	QRNMTKLQL (SEQ ID NO:173)	2.200
18	295	RGIQDVRRV (SEQ ID NO:179)	2.200
19	423	KKFARSDDEL (SEQ ID NO:122)	2.200
20	180	DPMGQQGSL (SEQ ID NO:59)	2.200

Table XXXIII
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Human HLA CW0702

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	319	EKRPFMCAY (SEQ ID NO:67)	26.880
2	326	AYPGCNKRY (SEQ ID NO:42)	24.000

3	40	FAPPGASAY (SEQ ID NO:74)	14.784
4	192	QYSVPPPVY (SEQ ID NO:176)	12.000
5	278	TPILCGAQY (SEQ ID NO:227)	12.000
6	219	TPYSSDNLY (SEQ ID NO:231)	12.000
7	213	QALLLRTPY (SEQ ID NO:160)	8.800
8	125	ARMFPNAPY (SEQ ID NO:38)	8.000
9	327	YPGCNKRYF (SEQ ID NO:250)	6.600
10	152	VTFDGTPSY (SEQ ID NO:244)	5.600
11	141	QPAIRNQGY (SEQ ID NO:170)	4.800
12	345	RKHTGEKPY (SEQ ID NO:184)	4.000
13	185	QGSLGEQQY (SEQ ID NO:166)	4.000
14	101	TGTAGACRY (SEQ ID NO:224)	4.000
15	375	RRHTGVKPF (SEQ ID NO:188)	4.000
16	263	GQSNHSTGY (SEQ ID NO:100)	4.000
17	163	TPSHHAAQF (SEQ ID NO:228)	3.000
18	33	QWAPVLDFA (SEQ ID NO:174)	2.688
19	130	NAPYLPSCS (SEQ ID NO:144)	2.640
20	84	HEEQCLSAF (SEQ ID NO:107)	2.400

Table XXXIV

Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Mouse MHC Class I Db

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	235	CMTWNQMNL (SEQ ID NO:49)	5255.712
2	126	RMFPNAPYL (SEQ ID NO:185)	1990.800
3	221	YSSDNLYQM (SEQ ID NO:253)	930.000
4	228	QMTSQLECM (SEQ ID NO:169)	33.701
5	239	NQMNLGATL (SEQ ID NO:151)	21.470
6	441	NMTKLQAL (SEQ ID NO:149)	19.908
7	437	MHQRNMTKL (SEQ ID NO:143)	19.837
8	136	SCLESQPAI (SEQ ID NO:198)	11.177
9	174	HSFKHEDPM (SEQ ID NO:110)	10.800
10	302	RVPGVAPTL (SEQ ID NO:195)	10.088
11	130	NAPYLPSC (SEQ ID NO:144)	8.400
12	10	ALLPAVPSL (SEQ ID NO:34)	5.988
13	208	SCTGSQALL (SEQ ID NO:202)	4.435
14	209	CTGSQALLL (SEQ ID NO:52)	3.548
15	238	WNQMNLGAT (SEQ ID NO:245)	3.300
16	218	RTPYSSDNL (SEQ ID NO:194)	3.185
17	24	CALPVSGAA (SEQ	2.851

		ID NO:43)	
18	18	LGGGGGCAL (SEQ ID NO:134).	2.177
19	142	PAIRNQGY (SEQ ID NO:152)	2.160
20	30	GAAQWAPVL (SEQ ID NO:86)	1.680

Table XXXV

Results of BIMAS HLA Peptide Binding Prediction Analysis for Binding of Human WT1 Peptides to Mouse MHC Class I Dd

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	112	FGPPPSQA (SEQ ID NO:76)	48.000
2	122	SGQARMFPN (SEQ ID NO:212)	36.000
3	104	AGACRYGPF (SEQ ID NO:31)	30.000
4	218	RTPYSSDNL (SEQ ID NO:194)	28.800
5	130	NAPYLPSC (SEQ ID NO:144)	20.000
6	302	RVPGVAPTL (SEQ ID NO:195)	20.000
7	18	LGGGGGCAL (SEQ ID NO:134)	20.000
8	81	AEPHEEQCL (SEQ ID NO:30)	10.000
9	29	SGAAQWAPV (SEQ ID NO:211)	7.200
10	423	KKFARSD (SEQ ID NO:122)	7.200
11	295	RGIQDVRRV (SEQ ID NO:179)	7.200
12	390	RKFSRSDHL (SEQ ID NO:183)	6.000
13	332	KRYFKLSHL (SEQ	6.000

		ID NO:127)	
14	362	RRFSRSDQL (SEQ ID NO:187)	6.000
15	417	RWPSCQKKF (SEQ ID NO:196)	6.000
16	160	YGHTPSHHA (SEQ ID NO:249)	6.000
17	20	GGGGCALPV (SEQ ID NO:92)	6.000
18	329	GCNKRYFKL (SEQ ID NO:90)	5.000
19	372	RHQRRHTGV (SEQ ID NO:181)	4.500
20	52	GGPAPPPAP (SEQ ID NO:93)	4.000

Table XXXVI

Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Mouse MHC Class I Kb

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	329	GCNKRYFKL (SEQ ID NO:90)	24.000
2	225	NLYQMTSQL (SEQ ID NO:147)	10.000
3	420	SCQKKFARS (SEQ ID NO:200)	3.960
4	218	RTPYSSDNL (SEQ ID NO:194)	3.630
5	437	MHQARNMTKL (SEQ ID NO:143)	3.600
6	387	TCQRKFSRS (SEQ ID NO:219)	3.600
7	302	RVPGVAPTL (SEQ ID NO:195)	3.300
8	130	NAPYLPSCSL (SEQ ID NO:144)	3.000
9	289	HTHGVFRGI (SEQ ID	3.000

		NO:113)	
10	43	PGASAYGSL (SEQ ID NO:153)	2.400
11	155	DGTPSYGHT (SEQ ID NO:56)	2.400
12	273	SDNHTTPIL (SEQ ID NO:204)	2.200
13	126	RMFPNAPYL (SEQ ID NO:185)	2.200
14	128	FPNAPYLPS (SEQ ID NO:79)	2.000
15	3	SDVRDLNAL (SEQ ID NO:206)	1.584
16	207	DSCTGSQAL (SEQ ID NO:61)	1.584
17	332	KRYFKLSHL (SEQ ID NO:127)	1.500
18	18	LGGGGGCAL (SEQ ID NO:134)	1.320
19	233	LECMTWNQM (SEQ ID NO:131)	1.320
20	441	NMTKLQLAL (SEQ ID NO:149)	1.200

Table XXXVII
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Mouse MHC Class I Kd

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	285	QYRIHTGV (SEQ ID NO:175)	600.000
2	424	KFARSDELV (SEQ ID NO:119)	288.000
3	334	YFKLSHLQM (SEQ ID NO:248)	120.000
4	136	SCLESQPTI (SEQ ID NO:199)	115.200
5	239	NQMNLGATL (SEQ ID NO:151)	115.200
6	10	ALLPAVSSL (SEQ ID NO:35)	115.200
7	47	AYGSLGGPA (SEQ ID NO:41)	86.400
8	180	DPMGQQGSL (SEQ ID NO:59)	80.000
9	270	GYESDNHTA (SEQ ID NO:105)	72.000
10	326	AYPGCNKRY (SEQ ID NO:42)	60.000
11	192	QYSVPPPVY (SEQ ID NO:176)	60.000
12	272	ESDNHTAPI (SEQ ID NO:70)	57.600
13	289	HTHGVFRGI (SEQ ID NO:113)	57.600
14	126	DVRDLNALL (SEQ ID NO:62)	57.600
15	4	CTGSQALLL (SEQ ID NO:52)	57.600
16	208	SCTGSQALL (SEQ ID NO:202)	48.000
17	441	NMTKLQLAL (SEQ ID NO:149)	48.000

18	207	DSCTGSQAL (SEQ ID NO:61)	48.000
19	130	NAPYLPSC (SEQ ID NO:144)	48.000
20	235	CMTWNQMNL (SEQ ID NO:49)	48.000

Table XXXVIII
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Mouse MHC Class I Kk

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	81	AEPHEEQCL (SEQ ID NO:30)	40.000
2	85	EEQCLSAFT (SEQ ID NO:65)	40.000
3	429	DELVRHHNM (SEQ ID NO:53)	20.000
4	315	SETSEKRPF (SEQ ID NO:209)	20.000
5	261	TEGQSNHST (SEQ ID NO:221)	20.000
6	410	SEKPFSCRW (SEQ ID NO:207)	10.000
7	272	ESDNHTTPI (SEQ ID NO:71)	10.000
8	318	SEKRPFMCA (SEQ ID NO:208)	10.000
9	138	LESQPAIRN (SEQ ID NO:132)	10.000
10	233	LECMTWNQM (SEQ ID NO:131)	10.000
11	298	QDVRRVPGV (SEQ ID NO:164)	10.000
12	84	HEEQCLSAF (SEQ ID NO:107)	10.000
13	349	GEKPYQCDF (SEQ ID NO:91)	10.000

14	289	HTHGVFRGI (SEQ ID NO:113)	10.000
15	179	EDPMGQQGS (SEQ ID NO:64)	8.000
16	136	SCLESQPAI (SEQ ID NO:198)	5.000
17	280	ILCGAQYRI (SEQ ID NO:116)	5.000
18	273	SDNHTTPIL (SEQ ID NO:204)	4.000
19	428	SDELVRHHN (SEQ ID NO:203)	4.000
20	3	SDVRDLNAL (SEQ ID NO:206)	4.000

Table XXXIX
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Mouse MHC Class I Ld

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	163	TPSHHAAQF (SEQ ID NO:228)	360.000
2	327	YPGCNKRYF (SEQ ID NO:250)	300.000
3	180	DPMGQQGSL (SEQ ID NO:59)	150.000
4	26	LPVSGAAQW (SEQ ID NO:138)	93.600
5	278	TPILCGAQY (SEQ ID NO:227)	72.000
6	141	QPAIRNQGY (SEQ ID NO:170)	60.000
7	219	TPYSSDNLY (SEQ ID NO:231)	60.000
8	303	VPGVAPTLV (SEQ ID NO:242)	60.000
9	120	ASSGQARMF (SEQ ID NO:40)	50.000

10	63	PPPPPHSF (SEQ ID NO:158)	45.000
11	113	GPPPPSQAS (SEQ ID NO:97)	45.000
12	157	TPSYGHTPS (SEQ ID NO:229)	39.000
13	207	DSCTGSQAL (SEQ ID NO:61)	32.500
14	110	GPFGPPPPS (SEQ ID NO:96)	30.000
15	82	EPHEEQCLS (SEQ ID NO:68)	30.000
16	412	KPFSCRWPS (SEQ ID NO:123)	30.000
17	418	WPSCQKKFA (SEQ ID NO:246)	30.000
18	221	YSSDNLYQM (SEQ ID NO:253)	30.000
19	204	TPTDSCTGS (SEQ ID NO:230)	30.000
20	128	FPNAPYLPS (SEQ ID NO:79)	30.000

Table XL
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Human WT1 Peptides to Cattle HLA A20

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	350	EKPYQCDFK (SEQ ID NO:66)	1000.00
2	319	EKRPFMCAY (SEQ ID NO:67)	500.000
3	423	KKFARSDDEL (SEQ ID NO:122)	500.000
4	345	RKHTGEKPY (SEQ ID NO:184)	500.000
5	390	RKFSRSDHL (SEQ ID NO:183)	500.000
6	137	CLESQPAIR (SEQ ID	120.000

		NO:47)	
7	380	VKPFQCKTC (SEQ ID NO:239)	100.000
8	407	GKTSEKPFs (SEQ ID NO:95)	100.000
9	335	FKLSHLQMH (SEQ ID NO:78)	100.000
10	247	LKGVAAGSS (SEQ ID NO:135)	100.000
11	370	LKRHQRRHT (SEQ ID NO:136)	100.000
12	258	VKWTEGQSN (SEQ ID NO:240)	100.000
13	398	LKTHTRTHT (SEQ ID NO:137)	100.000
14	331	NKRYFKLSH (SEQ ID NO:145)	100.000
15	357	FKDCERRFS (SEQ ID NO:77)	100.000
16	385	CKTCQRKFS (SEQ ID NO:46)	100.000
17	294	FRGIQDVRR (SEQ ID NO:81)	80.000
18	368	DQLKRHQRR (SEQ ID NO:60)	80.000
19	432	VRHHNMHQR (SEQ ID NO:243)	80.000
20	118	SQASSGQAR (SEQ ID NO:216)	80.000

Table XLI
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Mouse WT1 Peptides to Mouse MHC Class I A₀₂₀₁

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	126	RMFPNAPYL (SEQ ID NO:293)	313.968
2	187	SLGEQQYSV (SEQ ID NO:299)	285.163

3	10	ALLPAVSSL (SEQ ID NO:255)	181.794
4	225	NLYQMTSQL (SEQ ID NO:284)	68.360
5	292	GVFRGIQDV (SEQ ID NO:270)	51.790
6	93	TLHFSGQFT (SEQ ID NO:302)	40.986
7	191	QQYSVPPPV (SEQ ID NO:290)	22.566
8	280	ILCGAQYRI (SEQ ID NO:274)	17.736
9	441	NMTKLHVAL (SEQ ID NO:285)	15.428
10	235	CMTWNQMNL (SEQ ID NO:258)	15.428
11	7	DLNALLPAV (SEQ ID NO:261)	11.998
12	242	NLGATLKGM (SEQ ID NO:283)	11.426
13	227	YQMTSQLEC (SEQ ID NO:307)	8.573
14	239	NQMNLGATL (SEQ ID NO:286)	8.014
15	309	TLVRSASET (SEQ ID NO:303)	7.452
16	408	KTSEKPFSC (SEQ ID NO:277)	5.743
17	340	LQMHSRKHT (SEQ ID NO:280)	4.752
18	228	QMTSQLECM (SEQ ID NO:289)	4.044
19	37	VLDFAPPGA (SEQ ID NO:304)	3.378
20	302	RVSGVAPTL (SEQ ID NO:295)	1.869

Table XLII
Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Mouse WT1 Peptides to Mouse MHC Class I Db

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	221	YSSDNLYQM (SEQ ID NO:308)	312.000
2	126	RMFPNAPYL (SEQ ID NO:293)	260.000
3	235	CMTWNQMNL (SEQ ID NO:258)	260.000
4	437	MHQRNMTKL (SEQ ID NO:281)	200.000
5	238	WNQMNLGAT (SEQ ID NO:305)	12.000
6	130	NAPYLPSC (SEQ ID NO:282)	8.580
7	3	SDVRDLNAL (SEQ ID NO:298)	7.920
8	136	SCLESQPTI (SEQ ID NO:296)	7.920
9	81	AEPHEEQCL (SEQ ID NO:254)	6.600
10	10	ALLPAVSSL (SEQ ID NO:255)	6.600
11	218	RTPYSSDNL (SEQ ID NO:294)	6.000
12	441	NMTKLHVAL (SEQ ID NO:285)	3.432
13	228	QMTSQLECM (SEQ ID NO:289)	3.120
14	174	HSFKHEDPM (SEQ ID NO:272)	3.120
15	242	NLGATLKG (SEQ ID NO:283)	2.640
16	261	TEGQSNHGI (SEQ ID NO:301)	2.640

17	225	NLYQMTSQL (SEQ ID NO:284)	2.640
18	207	DSCTGSQAL (SEQ ID NO:263)	2.600
19	119	QASSGQARM (SEQ ID NO:288)	2.600
20	18	LGGGGGCGL (SEQ ID NO:279)	2.600

Table XLIII

Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Mouse WT1 Peptides to Mouse MHC Class I Kb

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	329	GCNKRYFKL (SEQ ID NO:268)	24.000
2	225	NLYQMTSQL (SEQ ID NO:284)	10.000
3	420	SCQKKFARS (SEQ ID NO:297)	3.960
4	218	RTPYSSDNL (SEQ ID NO:294)	3.630
5	437	MHQRNMTKL (SEQ ID NO:281)	3.600
6	387	TCQRKFSRS (SEQ ID NO:300)	3.600
7	289	HTHGVFRGI (SEQ ID NO:273)	3.000
8	130	NAPYLPSC (SEQ ID NO:282)	3.000
9	43	PGASAYGSL (SEQ ID NO:287)	2.400
10	155	DGAPSYGHT (SEQ ID NO:260)	2.400
11	126	RMFPNAPYL (SEQ ID NO:293)	2.200
12	128	FPNAPYLP (SEQ ID NO:267)	2.000

13	207	DSCTGSQAL (SEQ ID NO:263)	1.584
14	3	SDVRDLNAL (SEQ ID NO:298)	1.584
15	332	KRYFKLSHL (SEQ ID NO:276)	1.500
16	233	LECMTWNQM (SEQ ID NO:278)	1.320
17	18	LGGGGGCGL (SEQ ID NO:279)	1.320
18	242	NLGATLKGGM (SEQ ID NO:283)	1.200
19	123	GQARMFPN (SEQ ID NO:269)A	1.200
20	441	NMTKLHVAL (SEQ ID NO:285)	1.200

Table XLIV

Results of BIMAS HLA Peptide Binding Prediction Analysis for
Binding of Mouse WT1 Peptides to Mouse MHC Class I Kd

5

Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation of a Molecule Containing This Subsequence)
1	285	QYRIHTHGV (SEQ ID NO:291)	600.000
2	424	KFARSDELV (SEQ ID NO:275)	288.000
3	334	YFKLSHLQM (SEQ ID NO:306)	120.000
4	136	SCLESQPTI (SEQ ID NO:296)	115.200
5	239	NQMNLGATL (SEQ ID NO:286)	115.200
6	10	ALLPAVSSL (SEQ ID NO:255)	115.200
7	47	AYGSLGGPA (SEQ ID NO:256)	86.400
8	180	DPMGQQGSL (SEQ ID NO:262)	80.000

9	270	GYESDNHTA (SEQ ID NO:271)	72.000
10	192	QYSVPPPVY (SEQ ID NO:292)	60.000
11	326	AYPGCNKRY (SEQ ID NO:257)	60.000
12	289	HTHGVFRGI (SEQ ID NO:273)	57.600
13	4	DVRDLNALL (SEQ ID NO:264)	57.600
14	126	RMFPNAPYL (SEQ ID NO:293)	57.600
15	209	CTGSQALLL (SEQ ID NO:259)	48.000
16	86	EQCLSAFTL (SEQ ID NO:265)	48.000
17	302	RVSGVAPTL (SEQ ID NO:295)	48.000
18	218	RTPYSSDNL (SEQ ID NO:294)	48.000
19	272	ESDNHTAPI (SEQ ID NO:266)	48.000
20	225	NLYQMTSQL (SEQ ID NO:284)	48.000

Table XLV

Results of TSites Peptide Binding Prediction Analysis for
Human WT1 Peptides Capable of Eliciting a Helper T cell Response

5

Peptide	Sequence
p6-23	RDLNALLPAVPSLGGGG (SEQ ID NO:1)
p30-35	GAAQWA (SEQ ID NO:309)
p45-56	ASAYGSLGGPAP (SEQ ID NO:310)
p91-105	AFTVHFSGQFTGTAG (SEQ ID NO:311)
p117-139	PSQASSGQARMFPNAPYLPSCLE (SEQ ID NO:2)
p167-171	HAAQF (SEQ ID NO:312)
p202-233	CHTPTDSCCTGSQALLLRTPYSSDNLNLYQMTSQL (SEQ ID NO:313)
p244-262	GATLKGVAAGSSSSSVKWTE (SEQ ID NO:4)
p287-318	RIHTHGVFRGIQDVRRVPGVAPTLVRSASETS (SEQ ID NO:314)

p333-336	RYFK (SEQ ID NO:315)
p361-374	ERRFSRSDQLKRHQ (SEQ ID NO:316)
p389-410	QRKFSRSDHLKTHTRTHTGKTS (SEQ ID NO:317)
p421-441	CQKKFARSDELVRHHNMHQRN (SEQ ID NO:318)

Certain CTL peptides (shown in Table XLVI) were selected for further study. For each peptide in Table XLVI, scores obtained using BIMAS HLA peptide binding prediction analysis are provided.

5

Table XLVI
WT1 Peptide Sequences and HLA Peptide Binding Predictions

Peptide	Sequence	Comments
p329-337	GCNKRYFKL (SEQ ID NOs: 90 and 268)	Score 24,000
p225-233	NLYQMTSQL (SEQ ID NOs: 147 and 284)	binds also to class II and HLA A2, Kd, score 10,000
p235-243	CMTWNQMNL (SEQ ID NOs: 49 and 258)	binds also to HLA A2, score 5,255,712
p126-134	RMFPNAPYL (SEQ ID NOs: 185 and 293)	binds also to Kd, class II and HLA A2, score 1,990,800
p221-229	YSSDNLYQM (SEQ ID NOs: 253 and 308)	binds also to Ld, score 312,000
p228-236	QMTSQLECM (SEQ ID NOs: 169 and 289)	score 3,120
p239-247	NQMNLGATL (SEQ ID NOs: 151 and 286)	binds also to HLA A 0201, Kd, score 8,015
mouse p136-144	SCLESQPTI (SEQ ID NO:296)	binds also to Kd, 1 mismatch to human
human p136-144	SCLESQPAI (SEQ ID NO:198)	score 7,920
mouse p10-18	ALLPAVSSL	binds also to Kd, HLA A2, 1 mismatch

	(SEQ ID NO:255)	to human
human p10-18	ALLPAVPSL (SEQ ID NO:34)	score 6,600

Peptide binding to C57Bl/6 murine MHC was confirmed using the leukemia cell line RMA-S, as described by Ljunggren et al., *Nature* 346:476-480, 1990. In brief, RMA-S cells were cultured for 7 hours at 26°C in complete medium supplemented with 1% FCS. A total of 10⁶ RMA-S cells were added into each well of a 24-well plate and incubated either alone or with the designated peptide (25ug/ml) for 16 hours at 26°C and additional 3 hours at 37°C in complete medium. Cells were then washed three times and stained with fluorescein isothiocyanate-conjugated anti D^b or anti-K^b antibody (PharMingen, San Diego, CA). Labeled cells were washed twice, resuspended and fixed in 500ul of PBS with 1% paraformaldehyde and analyzed for fluorescence intensity in a flow cytometer (Becton-Dickinson FACSCalibur®). The percentage of increase of D^b or K^b molecules on the surface of the RMA-S cells was measured by increased mean fluorescent intensity of cells incubated with peptide compared with that of cells incubated in medium alone.

Mice were immunized with the peptides capable of binding to murine class I MHC. Following immunization, spleen cells were stimulated *in vitro* and tested for the ability to lyse targets incubated with WT1 peptides. CTL were evaluated with a standard chromium release assay (Chen et al., *Cancer Res.* 54:1065-1070, 1994). 10⁶ target cells were incubated at 37°C with 150μCi of sodium ⁵¹Cr for 90 minutes, in the presence or absence of specific peptides. Cells were washed three times and resuspended in RPMI with 5% fetal bovine serum. For the assay, 10⁴ ⁵¹Cr-labeled target cells were incubated with different concentrations of effector cells in a final volume of 200μl in U-bottomed 96-well plates. Supernatants were removed after 4 to 7 hours at 37°C, and the percentage specific lysis was determined by the formula:

% specific lysis = 100 x (experimental release - spontaneous release)/(maximum release-spontaneous release).

The results, presented in Table XLVII, show that some WT1 peptides can bind to class I MHC molecules, which is essential for generating CTL. Moreover, several of the peptides were able to elicit peptide specific CTL (Figures 9A and 9B), as determined using chromium release assays. Following immunization to CTL peptides p10-18 human, p136-144 human, p136-144 mouse and p235-243, peptide specific CTL lines were generated and clones were established. These results indicate that peptide specific CTL can kill malignant cells expressing WT1.

Table XLVII
Binding of WT1 CTL Peptides to mouse B6 class I antigens

Peptide	Binding Affinity to Mouse MHC Class I
Positive control	91%
negative control	0.5.-1.3%
p235-243	33.6%
p136-144 mouse	27.9%
p136-144 human	52%
p10-18: human	2.2%
p225-233	5.8%
p329-337	1.2%
p126-134	0.9%
p221-229	0.8%
p228-236	1.2%
p239-247	1%

Example 5

Use of a WT1 Polypeptide to Elicit WT1 Specific CTL in Mice

This Example illustrates the ability of a representative WT1 polypeptide to elicit CTL immunity capable of killing WT1 positive tumor cell lines.

P117-139, a peptide with motifs appropriate for binding to class I and class II MHC, was identified as described above using TSITES and BIMAS HLA peptide binding prediction analyses. Mice were immunized as described in Example 3. Following

immunization, spleen cells were stimulated *in vitro* and tested for the ability to lyse targets incubated with WT1 peptides, as well as WT1 positive and negative tumor cells. CTL were evaluated with a standard chromium release assay. The results, presented in Figures 10A-10D, show that P117 can elicit WT1 specific CTL capable of killing WT1 positive tumor cells, whereas no killing of WT1 negative cells was observed. These results demonstrate that peptide specific CTL in fact kill malignant cells expressing WT1 and that vaccine and T cell therapy are effective against malignancies that express WT1.

Similar immunizations were performed using the 9-mer class I MHC binding peptides p136-144, p225-233, p235-243 as well as the 23-mer peptide p117-139. Following immunization, spleen cells were stimulated *in vitro* with each of the 4 peptides and tested for ability to lyse targets incubated with WT1 peptides. CTL were generated specific for p136-144, p235-243 and p117-139, but not for p225-233. CTL data for p235-243 and p117-139 are presented in Figures 11A and 11B. Data for peptides p136-144 and p225-233 are not depicted.

CTL lysis demands that the target WT1 peptides are endogenously processed and presented in association with tumor cell class I MHC molecules. The above WT1 peptide specific CTL were tested for ability to lyse WT1 positive versus negative tumor cell lines. CTL specific for p235-243 lysed targets incubated with the p235-243 peptides, but failed to lyse cell lines that expressed WT1 proteins (Figure 11A). By marked contrast, CTL specific for p117-139 lysed targets incubated with p117-139 peptides and also lysed malignant cells expressing WT1 (Figure 11B). As a negative control, CTL specific for p117-139 did not lyse WT1 negative EL-4 (also referred to herein as E10).

Specificity of WT1 specific lysis was confirmed by cold target inhibition (Figures 12A-12B). Effector cells were plated for various effector: target ratios in 96-well U-bottom plates. A ten-fold excess (compared to hot target) of the indicated peptide-coated target without ^{51}Cr labeling was added. Finally, 10^4 ^{51}Cr -labeled target cells per well were added and the plates incubated at 37°C for 4 hours. The total volume per well was 200 μl .

Lysis of TRAMP-C by p117-139 specific CTL was blocked from 58% to 36% by EL-4 incubated with the relevant peptide p117-139, but not with EL-4 incubated with an irrelevant peptide (Figure 12A). Similarly, lysis of BLK-SV40 was blocked from 18% to 0% by EL-4 incubated with the relevant peptide p117-139 (Figure 12B). Results
 5 validate that WT1 peptide specific CTL specifically kill malignant cells by recognition of processed WT1.

Several segments with putative CTL motifs are contained within p117-139. To determine the precise sequence of the CTL epitope all potential 9-mer peptides within p117-139 were synthesized (Table XLVIII). Two of these peptides (p126-134 and p130-
 10 138) were shown to bind to H-2^b class I molecules (Table XLVIII). CTL generated by immunization with p117-139 lysed targets incubated with p126-134 and p130-138, but not the other 9-mer peptides within p117-139 (Figure 13A).

The p117-139 specific CTL line was restimulated with either p126-134 or p130-138. Following restimulation with p126-134 or p130-138, both T cell lines
 15 demonstrated peptide specific lysis, but only p130-138 specific CTL showed lysis of a WT1 positive tumor cell line (Figures 13B and 13C). Thus, p130-138 appears to be the naturally processed epitope.

20 Table XLVIII
Binding of WT1 CTL 9mer Peptides within p117-139 to mouse B6 class I antigens

Peptide	Binding Affinity to Mouse MHC Class I
P117-125 PSQASSGQA (SEQ ID NO:221)	2%
P118-126 SQASSGQAR (SEQ ID NO:216)	2%
P119-127 QASSGQARM (SEQ ID NOs: 161 and 288)	2%
P120-128 ASSGQARMF (SEQ ID NO:40)	1%
P121-129 SSGQARMFP (SEQ ID NO:222)	1%
P122-130 SGQARMFPN (SEQ ID NO:223)	1%

NO:212)	
P123-131 GQARMFPNA (SEQ ID NOs: 98 and 269)	1%
P124-132 QARMFPNAP (SEQ ID NO:223)	1%
P125-133 ARMFPNAPY (SEQ ID NO:38)	1%
P126-134 RMFPNAPYL (SEQ ID NOs: 185 and 293)	79%
P127-135 MFPNAPYLP (SEQ ID NO:224)	2%
P128-136 FPNAPYLPS (SEQ ID NOs: 79 and 267)	1%
P129-137 PNAPYLPSC (SEQ ID NO:225)	1%
P130-138 NAPYLPSCLE (SEQ ID NOs: 144 and 282)	79%
P131-139 APYLPSCLE (SEQ ID NO:226)	1%

Example 6

5 Identification of WT1 Specific mRNA in Mouse Tumor Cell Lines

This Example illustrates the use of RT-PCR to detect WT1 specific mRNA in cells and cell lines.

10 Mononuclear cells were isolated by density gradient centrifugation, and were immediately frozen and stored at -80°C until analyzed by RT-PCR for the presence of WT1 specific mRNA. RT-PCR was generally performed as described by Fraizer et al., *Blood* 86:4704-4706, 1995. Total RNA was extracted from 10⁷ cells according to standard procedures. RNA pellets were resuspended in 25 µL diethylpyrocarbonate treated water
15 and used directly for reverse transcription. The zinc-finger region (exons 7 to 10) was amplified by PCR as a 330 bp mouse cDNA. Amplification was performed in a thermocycler during one or, when necessary, two sequential rounds of PCR. AmpliTaq

DNA Polymerase (Perkin Elmer Cetus, Norwalk, CT), 2.5 mM MgCl₂ and 20 pmol of each primer in a total reaction volume of 50µl were used. Twenty µL aliquots of the PCR products were electrophoresed on 2% agarose gels stained with ethidium bromide. The gels were photographed with Polaroid film (Polaroid 667, Polaroid Ltd., Hertfordshire, England). Precautions against cross contamination were taken following the recommendations of Kwok and Higuchi, *Nature* 339:237-238, 1989. Negative controls included the cDNA- and PCR-reagent mixes with water instead of cDNA in each experiment. To avoid false negatives, the presence of intact RNA and adequate cDNA generation was evaluated for each sample by a control PCR using β-actin primers. Samples that did not amplify with these primers were excluded from analysis.

Primers for amplification of WT1 in mouse cell lines were: P115: 1458-1478: 5' CCC AGG CTG CAA TAA GAG ATA 3' (forward primer; SEQ ID NO:21); and P116: 1767-1787: 5' ATG TTG TGA TGG CGG ACC AAT 3' (reverse primer; SEQ ID NO:22) (*see* Inoue et al, *Blood* 88:2267-2278, 1996; Fraizer et al., *Blood* 86:4704-4706, 1995).

Beta Actin primers used in the control reactions were: 5' GTG GGG CGC CCC AGG CAC CA 3' (sense primer; SEQ ID NO:23); and 5' GTC CTT AAT GTC ACG CAC GAT TTC 3' (antisense primer; SEQ ID NO:24)

Primers for use in amplifying human WT1 include: P117: 954-974: 5' GGC ATC TGA GAC CAG TGA GAA 3' (SEQ ID NO:25); and P118: 1434-1414: 5' GAG AGT CAG ACT TGA AAG CAGT 3' (SEQ ID NO:5). For nested RT-PCR, primers may be: P119: 1023-1043: 5' GCT GTC CCA CTT ACA GAT GCA 3' (SEQ ID NO:26); and P120: 1345-1365: 5' TCA AAG CGC CAG CTG GAG TTT 3' (SEQ ID NO:27).

Table XLVIII shows the results of WT1 PCR analysis of mouse tumor cell lines. Within Table IV, (++++) indicates a strong WT1 PCR amplification product in the first step RT PCR, (++) indicates a WT1 amplification product that is detectable by first step WT1 RT PCR, (+) indicates a product that is detectable only in the second step of WT1 RT PCR, and (-) indicates WT1 PCR negative.

Table XLIX
Detection of WT1 mRNA in Mouse Tumor Cell Lines

Cell Line	WT1 mRNA
K562 (human leukemia; ATCC): Positive control; (Lozzio and Lozzio, <i>Blood</i> 45:321-334, 1975)	+++
TRAMPC (SV40 transformed prostate, B6); Foster et al., <i>Cancer Res.</i> 57:3325-3330, 1997	+++
BLK-SV40 HD2 (SV40-transf. fibroblast, B6; ATCC); <i>Nature</i> 276:510-511, 1978	++
CTLL (T-cell, B6; ATCC); Gillis, <i>Nature</i> 268:154-156, 1977)	+
FM (FBL-3 subline, leukemia, B6); Glynn and Fefer, <i>Cancer Res.</i> 28:434-439, 1968	+
BALB 3T3 (ATCC); Aaroston and Todaro, <i>J. Cell. Physiol.</i> 72:141-148, 1968	+
S49.1 (Lymphoma, T-cell like, B/C; ATCC); Horibata and Harris, <i>Exp. Cell. Res.</i> 60:61, 1970	+
BNL CL.2 (embryonic liver, B/C; ATCC); <i>Nature</i> 276:510-511, 1978	+
MethA (sarcoma, B/C); Old et al., <i>Ann. NY Acad. Sci.</i> 101:80-106, 1962	-
P3.6.2.8.1 (myeloma, B/C; ATCC); <i>Proc. Natl. Acad. Sci. USA</i> 66:344, 1970	-
P2N (leukemia, DBA/2; ATCC); Melling et al., <i>J. Immunol.</i> 117:1267-1274, 1976	-
BCL1 (lymphoma, B/C; ATCC); Slavin and Strober, <i>Nature</i> 272:624-626, 1977	-
LSTRA (lymphoma, B/C); Glynn et al., <i>Cancer Res.</i> 28:434-439, 1968	-
E10/EL-4 (lymphoma, B6); Glynn et al., <i>Cancer Res.</i> 28:434-439, 1968	-

5

From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

CLAIMS

1. A polypeptide comprising an immunogenic portion of a native WT1, or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with WT1-specific antisera and/or T-cell lines or clones is not substantially diminished, wherein the polypeptide comprises no more than 16 consecutive amino acid residues present within a native WT1 polypeptide.

2. A polypeptide according to claim 1, wherein the immunogenic portion binds to an MHC class I molecule.

3. A polypeptide according to claim 1, wherein the immunogenic portion binds to an MHC class II molecule.

4. A polypeptide according to claim 1, wherein the polypeptide comprises a sequence selected from the group consisting of:

- (a) sequences recited in one or more of Tables II - XLVI;
- (b) variants of the foregoing sequences that differ in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antisera and/or T-cell lines or clones is not substantially diminished; and
- (c) mimetics of the foregoing sequences, wherein the ability of the mimetic to react with antigen-specific antisera and/or T-cell lines or clones is not substantially diminished.

5. A polypeptide according to claim 1, wherein the polypeptide comprises a sequence selected from the group consisting of:

- (a) ALLPAVPSL (SEQ ID NO:34), GATLKGVA A (SEQ ID NO:88), CMTWNQMNL (SEQ ID NOs: 49 and 258), SCLESQPTI (SEQ ID NOs: 199 and 296), SCLESQPAI (SEQ ID NO:198), NLYQMTSQL (SEQ ID NOs: 147 and

284); ALLPAVSSL (SEQ ID NOs: 35 and 255), RMFPNAPYL (SEQ ID NOs: 185 and 293);

(b) variants of the foregoing sequences that differ in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antisera and/or T-cell lines or clones is not substantially diminished; and

(c) mimetics of the foregoing sequences, wherein the ability of the mimetic to react with antigen-specific antisera and/or T-cell lines or clones is not substantially diminished.

6. A polypeptide according to claim 1, wherein the polypeptide comprises 4-16 consecutive amino acids of a native WT1 polypeptide.

7. A polypeptide according to claim 1, wherein the polypeptide comprises 8-10 consecutive amino acids of a native WT1 polypeptide.

8. A polypeptide comprising an immunogenic portion of amino acid residues 1 - 174 of a native WT1 polypeptide, or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with WT1-specific T-cell lines or clones is not substantially diminished, wherein the polypeptide comprises no more than 16 consecutive amino acid residues present within amino acids 175 to 449 of the native WT1 polypeptide.

9. A polypeptide comprising a variant of an immunogenic portion of WT1 that differs in substitutions at between 1 and 3 amino acid positions within the immunogenic portion, such that the ability of the variant to react with WT1-specific antisera and/or T-cell lines or clones is enhanced relative to a native WT1.

10. A mimetic of an immunogenic portion of a WT1 polypeptide, wherein at least one amino acid residue is replaced by a compound that is not an amino

acid, such that the ability of the mimetic to react with antigen-specific antisera and/or T-cell lines or clones is not diminished.

11. A pharmaceutical composition comprising a polypeptide according to claim 1, in combination with a pharmaceutically acceptable carrier or excipient.

12. A pharmaceutical composition according to claim 11, wherein the polypeptide comprises 4-16 consecutive amino acids of a native WT1 polypeptide.

13. A pharmaceutical composition according to claim 11, wherein the polypeptide comprises 8-16 consecutive amino acids of a native WT1 polypeptide.

14. A pharmaceutical composition comprising a polypeptide according to claim 8, in combination with a pharmaceutically acceptable carrier or excipient.

15. A vaccine comprising a polypeptide according to claim 1, in combination with a non-specific immune response enhancer.

16. A vaccine according to claim 15, wherein the polypeptide comprises 4-16 consecutive amino acids of a native WT1 polypeptide.

17. A vaccine according to claim 15, wherein the polypeptide comprises 8-10 consecutive amino acids of a native WT1 polypeptide.

18. A vaccine according to claim 15, wherein the immune response enhancer is an adjuvant.

19. A vaccine comprising a polypeptide according to claim 8, in combination with a non-specific immune response enhancer.

20. A vaccine according to claim 19, wherein the immune response enhancer is an adjuvant.

21. A vaccine comprising:

(a) a WT1 polypeptide, wherein the polypeptide comprises an immunogenic portion of a native WT1 or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific T cell lines or clones is not substantially diminished; and

(b) a non-specific immune response enhancer that preferentially enhances a T cell response in a patient.

22. A vaccine according to claim 21, wherein the immune response enhancer is selected from the group consisting of Montanide ISA50, Seppic MONTANIDE ISA 720, cytokines (*e.g.*, GM-CSF, Flat3-ligand), microspheres, dimethyl dioctadecyl ammoniumbromide (DDA) based adjuvants, AS-1, AS-2, Ribi Adjuvant system based adjuvants, QS21, saponin based adjuvants, Syntex adjuvant in its microfluidized form, MV, ddMV, immune stimulating complex (iscom) based adjuvants and inactivated toxins.

23. A pharmaceutical composition comprising a mimetic according to claim 10, in combination with a pharmaceutically acceptable carrier or excipient.

24. A vaccine comprising a mimetic according to claim 10, in combination with a non-specific immune response enhancer.

25. A polynucleotide encoding a polypeptide according to claim 1 or claim 8.

26. A pharmaceutical composition, comprising:

(a) a polynucleotide encoding a WT1 polypeptide, wherein the polypeptide comprises an immunogenic portion of a native WT1 or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antibodies and/or T cell lines or clones is not substantially diminished; and

(b) a pharmaceutically acceptable carrier or excipient.

27. A pharmaceutical composition, comprising:

(a) an antibody or antigen-binding fragment thereof that specifically binds to a WT1 polypeptide; and

(b) a pharmaceutically acceptable carrier or excipient.

28. A pharmaceutical composition, comprising:

(a) a T cell that specifically reacts with a WT1 polypeptide; and

(b) a pharmaceutically acceptable carrier or excipient.

29. A pharmaceutical composition, comprising:

(a) an antigen presenting cell that expresses

(i) a WT1 polypeptide that comprises an immunogenic portion of a native WT1 or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antibodies and/or T cell lines or clones is not substantially diminished; and

(b) a pharmaceutically acceptable carrier or excipient.

30. A vaccine, comprising:

(a) a polynucleotide encoding a WT1 polypeptide, wherein the polypeptide comprises an immunogenic portion of a native WT1 or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antibodies and/or T cell lines or clones is not substantially diminished; and

(b) a non-specific immune response enhancer.

31. A vaccine, comprising:

(a) an antigen presenting cell that expresses:

(i) a WT1 polypeptide that comprises an immunogenic portion of a native WT1 or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antibodies and/or T cell lines or clones is not substantially diminished; and

(b) a non-specific immune response enhancer.

32. A vaccine comprising:

(a) an anti-idiotypic antibody or antigen-binding fragment thereof that is specifically bound by an antibody that specifically binds to an immunogenic portion of WT1; and

(b) non-specific immune response enhancer.

33. A vaccine according to any one of claims 30-32, wherein the immune response enhancer is an adjuvant.

34. A vaccine according to any one of claims 30-32, wherein the immune response enhancer preferentially enhances a T cell response in a patient.

35. A method for enhancing or inducing an immune response in a human patient, comprising administering to a patient a pharmaceutical composition comprising:

(a) a WT1 polypeptide that comprises an immunogenic portion of a native WT1 or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antibodies and/or T cell lines or clones is not substantially diminished; and

(b) a physiologically acceptable carrier or excipient;

and thereby enhancing or inducing an immune response specific for WT1 or a cell expressing WT1 in the human patient.

36. A method for enhancing or inducing an immune response in a patient, comprising administering to a patient a pharmaceutical composition according to any one of claims 11, 14, 23 or 26-29.

37. A method for enhancing or inducing an immune response in a human patient, comprising administering to a patient a vaccine comprising:

(a) a WT1 polypeptide that comprises an immunogenic portion of a native WT1 or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antibodies and/or T cell lines or clones is not substantially diminished; and

(b) a non-specific immune response enhancer;

and thereby enhancing or inducing an immune response specific for WT1 or a cell expressing WT1 in the human patient.

38. A method for enhancing or inducing an immune response in a patient, comprising administering to a patient a vaccine according to any one of claims 15, 19, 21, 24 or 30-32, and thereby enhancing or inducing an immune response specific for WT1 or a cell expressing WT1 in the patient.

39. A method for inhibiting the development of a malignant disease associated with WT1 expression in a human patient, comprising administering to a human patient a pharmaceutical composition comprising:

(a) a WT1 polypeptide that comprises an immunogenic portion of a native WT1 or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antibodies and/or T cell lines or clones is not substantially diminished; and

(b) a physiologically acceptable carrier or excipient;

and thereby inhibiting the development of a malignant disease associated with WT1 expression in the human patient.

40. A method for inhibiting the development of a malignant disease associated with WT1 expression in a patient, comprising administering to a patient a pharmaceutical composition according to any one of claims 11, 14, 23 or 26-29, and thereby inhibiting the development of a malignant disease in the patient.

41. A method for inhibiting the development of a malignant disease associated with WT1 expression in a human patient, comprising administering to a patient a vaccine comprising:

(a) a WT1 polypeptide that comprises an immunogenic portion of a native WT1 or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antibodies and/or T cell lines or clones is not substantially diminished; and

(b) a non-specific immune response enhancer;
and thereby inhibiting the development of a malignant disease in the patient.

42. A method for inhibiting the development of a malignant disease associated with WT1 expression in a patient, comprising administering to a patient a vaccine according to any one of claims 15, 19, 21, 24 or 30-32, and thereby inhibiting the development of a malignant disease in the patient.

43. A method according to claim 39 or claim 41, wherein the malignant disease is a leukemia.

44. A method according to claim 43, wherein the leukemia is acute myeloid leukemia, acute lymphocytic leukemia or chronic myeloid leukemia.

45. A method according to claim 39 or claim 41, wherein the malignant disease is a cancer.

46. A method according to claim 45, wherein the cancer is breast, lung, thyroid or gastrointestinal cancer or a melanoma.

47. A method according to claim 40, wherein the malignant disease is a leukemia.

48. A method according to claim 47, wherein the leukemia is acute myeloid leukemia, acute lymphocytic leukemia or chronic myeloid leukemia.

49. A method according to claim 40, wherein the malignant disease is a cancer.

50. A method according to claim 49, wherein the cancer is breast, lung, thyroid or gastrointestinal cancer or a melanoma.

51. A method according to claim 42, wherein the malignant disease is a leukemia.

52. A method according to claim 51, wherein the leukemia is acute myeloid leukemia, acute lymphocytic leukemia or chronic myeloid leukemia.

53. A method according to claim 42, wherein the malignant disease is a cancer.

54. A method according to claim 53, wherein the cancer is breast, lung, thyroid or gastrointestinal cancer or a melanoma.

55. A method according to claim 39, wherein the pharmaceutical composition comprises a WT1 polypeptide that comprises a sequence selected from the

group consisting of sequences recited in one or more of Tables II - XLVI and variants of the foregoing sequences that differ in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antisera and/or T-cell lines or clones is not diminished.

56. A method according to claim 39, wherein the pharmaceutical composition comprises a WT1 polypeptide that comprises a sequence selected from the group consisting of ALLPAVPSL (SEQ ID NO:34), GATLKGVAA (SEQ ID NO:88), CMTWNQMNL (SEQ ID NOS: 49 and 258), SCLESQPTI (SEQ ID NOS: 199 and 296), SCLESQPAI (SEQ ID NO:198), NLYQMTSQL (SEQ ID NOS: 147 and 284), ALLPAVSSL (SEQ ID NOS: 35 and 255); RMFPNAPYL (SEQ ID NOS: 185 and 293) and variants of the foregoing sequences that differ in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antisera and/or T-cell lines or clones is not diminished.

57. A method according to claim 41, wherein the vaccine comprises a WT1 polypeptide that comprises a sequence selected from the group consisting of sequences recited in one or more of Tables II - XLVI and variants of the foregoing sequences that differ in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antisera and/or T-cell lines or clones is not diminished.

58. A method according to claim 41, wherein the vaccine comprises a WT1 polypeptide that comprises a sequence selected from the group consisting of ALLPAVPSL (SEQ ID NO:34), GATLKGVAA (SEQ ID NO:88), CMTWNQMNL (SEQ ID NOS: 49 and 258), SCLESQPTI (SEQ ID NOS: 199 and 296), SCLESQPAI (SEQ ID NO:198), NLYQMTSQL (SEQ ID NOS: 147 and 284), ALLPAVSSL (SEQ ID NOS: 35 and 255), RMFPNAPYL (SEQ ID NOS: 185 and 293) and variants of the foregoing sequences that differ in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antisera and/or T-cell lines or clones is not diminished.

59. A method for removing cells expressing WT1 from bone marrow, peripheral blood, or a fraction of bone marrow or peripheral blood, comprising contacting bone marrow, peripheral blood or a fraction of bone marrow or peripheral blood with T cells that specifically react with a WT1 polypeptide, wherein the step of contacting is performed under conditions and for a time sufficient to permit the removal of WT1 positive cells to less than 10% of the number of myeloid or lymphatic cells in the bone marrow, peripheral blood or a fraction of bone marrow or peripheral blood.

60. A method for inhibiting the development of a malignant disease associated with WT1 expression in a patient, comprising administering to a patient bone marrow, peripheral blood or a fraction of bone marrow or peripheral blood prepared according to the method of claim 59.

61. A method according to claim 60, wherein the bone marrow, peripheral blood or fraction is autologous.

62. A method according to claim 60, wherein the bone marrow, peripheral blood or fraction is syngeneic or allogeneic.

63. A method for stimulating and/or expanding T cells, comprising contacting T cells with a WT1 polypeptide, a polynucleotide encoding a WT1 polypeptide and/or an antigen presenting cell that expresses a WT1 polypeptide under conditions and for a time sufficient to permit the stimulation and/or expansion of T cells.

64. A method according to claim 63, wherein the T cells are present within bone marrow, peripheral blood or a fraction of bone marrow or peripheral blood.

65. A method according to claim 63, wherein the bone marrow, peripheral blood or fraction is obtained from a patient afflicted with a malignant disease associated with WT1 expression.

66. A method according to claim 63, wherein the bone marrow, peripheral blood or fraction is obtained from a mammal that is not afflicted with a malignant disease associated with WT1 expression.

67. A method according to claim 63, wherein the T cells are cloned prior to expansion.

68. A method for stimulating and/or expanding T cells in a mammal, comprising administering to a mammal a pharmaceutical composition comprising:

- (a) one or more of:
 - (i) a WT1 polypeptide;
 - (ii) a polynucleotide encoding a WT1 polypeptide; or
 - (iii) an antigen-presenting cell that expresses a WT1 polypeptide; and
 - (b) a physiologically acceptable carrier or excipient;
- and thereby stimulating and/or expanding T cells in a mammal.

69. A method for stimulating and/or expanding T cells in a mammal, comprising administering to a mammal a vaccine comprising:

- (a) one or more of:
 - (i) a WT1 polypeptide;
 - (ii) a polynucleotide encoding a WT1 polypeptide; or
 - (iii) an antigen-presenting cell that expresses a WT1 polypeptide; and
 - (b) a non-specific immune response enhancer;
- and thereby stimulating and/or expanding T cells in a mammal.

70. A method for inhibiting the development of a malignant disease associated with WT1 expression in a patient, comprising administering to a patient T cells prepared according to the method of claim 63.

71. A method according to claim 70, wherein the bone marrow, peripheral blood or fraction is obtained from a patient afflicted with a malignant disease associated with WT1 expression.

72. A method according to claim 70, wherein the bone marrow, peripheral blood or fraction is obtained from a mammal that is not afflicted with a malignant disease associated with WT1 expression.

73. A method for monitoring the effectiveness of an immunization or therapy for a malignant disease associated with WT1 expression in a patient, comprising the steps of:

(a) incubating a first biological sample with one or more of:

- (i) a WT1 polypeptide;
- (ii) a polynucleotide encoding a WT1 polypeptide; or
- (iii) an antigen-presenting cell that expresses a WT1 polypeptide

wherein the first biological sample is obtained from a patient prior to a therapy or immunization, and wherein the incubation is performed under conditions and for a time sufficient to allow immunocomplexes to form;

(b) detecting immunocomplexes formed between the WT1 polypeptide and antibodies in the biological sample that specifically bind to the WT1 polypeptide;

(c) repeating steps (a) and (b) using a second biological sample obtained from the patient following therapy or immunization; and

(d) comparing the number of immunocomplexes detected in the first and second biological samples, and therefrom monitoring the effectiveness of the therapy or immunization in the patient.

74. A method according to claim 73, wherein the step of detecting comprises (a) incubating the immunocomplexes with a detection reagent that is capable of binding to the immunocomplexes, wherein the detection reagent comprises a reporter group, (b) removing unbound detection reagent, and (c) detecting the presence or absence of the reporter group.

75. A method according to claim 74, wherein the detection reagent comprises a second antibody, or antigen-binding fragment thereof, capable of binding to the antibodies that specifically bind to the WT1 polypeptide.

76. A method according to claim 74, wherein the detection reagent comprises Protein A.

77. A method according to claim 74, wherein the reporter group is selected from the group consisting of radioisotopes, fluorescent groups, luminescent groups, enzymes, biotin and dye particles.

78. A method according to claim 73 wherein a reporter group is bound to the WT1 polypeptide, and wherein the step of detecting comprises removing unbound WT1 polypeptide and subsequently detecting the presence or absence of the reporter group.

79. A method for monitoring the effectiveness of an immunization or therapy for a malignant disease associated with WT1 expression in a patient, comprising the steps of:

(a) incubating a first biological sample with one or more of:

- (i) a WT1 polypeptide;
- (ii) a WT1 polynucleotide encoding a WT1 polypeptide; or
- (iii) an antigen-presenting cell that expresses a WT1 polypeptide;

wherein the biological sample comprises CD4+ and/or CD8+ T cells and is obtained from a patient prior to a therapy or immunization, and wherein the incubation is performed under conditions and for a time sufficient to allow specific activation, proliferation and/or lysis of T cells;

- (b) detecting an amount of activation, proliferation and/or lysis of the T cells;

- (c) repeating steps (a) and (b) using a second biological sample comprising CD4+ and/or CD8+ T cells, wherein the second biological sample is obtained from the same patient following therapy or immunization; and

- (d) comparing the amount of activation, proliferation and/or lysis of T cells in the first and second biological samples, and therefrom monitoring the effectiveness of the therapy or immunization in the patient.

80. A method according to claim 73 or claim 79, wherein the malignant disease is a cancer or a leukemia.

81. A method for inhibiting the development of a malignant disease associated with WT1 expression in a patient, comprising the steps of:

- (a) incubating CD4⁺ T cells isolated from a patient with one or more of:

- (i) a WT1 polypeptide;
- (ii) a polynucleotide encoding a WT1 polypeptide; or
- (iii) an antigen presenting cell that expresses a WT1 polypeptide;

such that the T cells proliferate; and

(b) administering to the patient an effective amount of the proliferated T cells, and therefrom inhibiting the development of a malignant disease in the patient.

82. A method according to claim 81, wherein the malignant disease is a cancer or a leukemia.

83. A method according to claim 81, wherein the step of incubating the T cells is repeated one or more times.

84. A method for inhibiting the development of a malignant disease associated with WT1 expression in a patient, comprising the steps of:

(a) incubating CD4⁺ T cells isolated from a patient with one or more of:

- (i) a WT1 polypeptide;
- (ii) a polynucleotide encoding a WT1 polypeptide; or
- (iii) an antigen presenting cell that expresses a WT1 polypeptide;

such that the T cells proliferate;

(b) cloning one or more cells that proliferated in the presence of WT1 polypeptide; and

(c) administering to the patient an effective amount of the cloned T cells.

85. A method according to claim 84, wherein the malignant disease is a cancer or a leukemia.

86. A method for inhibiting the development of a malignant disease associated with WT1 expression in a patient, comprising the steps of:

- (a) incubating CD8⁺ T cells isolated from a patient with one or more of:
- (i) a WT1 polypeptide;
 - (ii) a polynucleotide encoding a WT1 polypeptide; or
 - (iii) an antigen presenting cell expressing a WT1 polypeptide;
- such that the T cells proliferate; and
- (b) administering to the patient an effective amount of the proliferated T cells, and therefrom inhibiting the development of a malignant disease in the patient.

87. A method according to claim 86, wherein the malignant disease is a cancer or a leukemia.

88. A method according to claim 86, wherein the step of incubating the T cells is repeated one or more times.

89. A method for inhibiting the development of a malignant disease associated with WT1 expression in a patient, comprising the steps of:

- (a) incubating CD8⁺ T cells isolated from a patient with one or more of:
- (i) a WT1 polypeptide
 - (ii) a polynucleotide encoding a WT1 polypeptide; or
 - (iii) an antigen presenting cell that expresses a WT1 polypeptide;
- such that the T cells proliferate;
- (b) cloning one or more cells that proliferated in the presence of WT1 polypeptide; and
- (c) administering to the patient an effective amount of the cloned T cells.

90. A method according to claim 89, wherein the malignant disease is a cancer or a leukemia.

91. A method for determining the presence or absence of a malignant disease associated with WT1 expression in a patient, comprising the steps of:

(a) incubating CD4⁺ T cells isolated from a patient with one or more of:

(i) a WT1 polypeptide;
(ii) a polynucleotide encoding a WT1 polypeptide; or
(iii) an antigen presenting cell that expresses a WT1 polypeptide; and

(b) detecting the presence or absence of specific activation of the T cells, therefrom determining the presence or absence of a malignant disease associated with WT1 expression.

92. A method according to claim 91, wherein the malignant disease is a cancer or a leukemia.

93. A method according to claim 91, wherein the step of detecting comprises detecting the presence or absence of proliferation of the T cells.

94. A method for determining the presence or absence of a malignant disease associated with WT1 expression in a patient, comprising the steps of:

(a) incubating CD8⁺ T cells isolated from a patient with a one or more of:

(i) a WT1 polypeptide;
(ii) a polynucleotide encoding a WT1 polypeptide; or
(iii) an antigen presenting cell that expresses a WT1 polypeptide; and

(b) detecting the presence or absence of specific activation of the T cells, thereby determining the presence or absence of a malignant disease associated with WT1 expression.

95. A method according to claim 94, wherein the malignant disease is a cancer or a leukemia.

96. A method according to claim 94 wherein the step of detecting comprises detecting the presence or absence of generation of cytolytic activity.

97. A method for determining the presence or absence of a malignant disease associated with WT1 expression in a patient, comprising the steps of:

(a) incubating a biological sample obtained from a patient with one or more of:

- (i) a WT1 polypeptide;
- (ii) a polynucleotide encoding a WT1 polypeptide; or
- (iii) an antigen presenting cell that expresses a WT1 polypeptide;

wherein the incubation is performed under conditions and for a time sufficient to allow immunocomplexes to form; and

(b) detecting immunocomplexes formed between the WT1 polypeptide and antibodies in the biological sample that specifically bind to the WT1 polypeptide; and therefrom determining the presence or absence of a malignant disease associated with WT1 expression.

98. A method according to claim 97, wherein the malignant disease is a cancer or a leukemia.

99. A method according to claim 97, wherein the step of detecting comprises (a) incubating the immunocomplexes with a detection reagent that is capable

of binding to the immunocomplexes, wherein the detection reagent comprises a reporter group, (b) removing unbound detection reagent, and (c) detecting the presence or absence of the reporter group.

100. A method according to claim 99, wherein the detection reagent comprises a second antibody, or antigen-binding fragment thereof, capable of binding to the antibodies that specifically bind to the WT1 polypeptide.

101. A method according to claim 99, wherein the detection reagent comprises Protein A.

102. A method according to claim 99, wherein the reporter group is selected from the group consisting of radioisotopes, fluorescent groups, luminescent groups, enzymes, biotin and dye particles.

103. A method according to claim 97 wherein a reporter group is bound to the WT1 polypeptide, and wherein the step of detecting comprises removing unbound WT1 polypeptide and subsequently detecting the presence or absence of the reporter group.






FIG. 2

**FIG. 3**

**FIG. 4**






FIG. 5A-5C

*FIG. 6A and 6B*



**FIG. 7A-7D**

5 10 15 20 25 30 35 40 45 50 55 60 65 70 75
MGSDVRLNALLPAVPSTLGGGGCCALPVSGAAGWAFVLDFAFPGASAYGSLGGZAPPPAPPPPPPPPPPPHSEFIRGE
.....AAAAAAAAAAAAAAAAAAAAA.....AAAAA.....AAAAAAAAAAAAA.....
.....RRRR.....
.....
.....
80 85 90 95 100 105 110 115 120 125 130 135 140 145 150
PSWGGAEFHEEQCLSAFTVHFSGQFTGTAGACRYGTFGPPPPSQASSGCARMF7NAPYL2SCLESQPAIRNQGYS
.....AAA.....AAAA.....AAA.....AAAAA.....
.....RRRR.....RRRRR.....
.....DDDDDDDDDD.....
.....
155 160 165 170 175 180 185 190 195 200 205 210 215 220 225
TVTTEGTFSYGHFSSHAAQFFNHSTFHEDFMGQQGSLGZCQYSVPPPVYGCFTPTDSCTGSCALLRTPYSSON
.....AAAAA.....AAAAA.....AA
.....RRRR.....
.....DDDDDDDDDDDDDDDDDD.....
.....
230 235 240 245 250 255 260 265 270 275 280 285 290 295 300
LYQMTSCLECMWNCNHLGATLKGVAAGSSSSVHTTGGQSNHSTOYESONHTT3ELCGAQYRINTHGVTRGIGQOV
AAAAAAAA.....AAA.AAA.....AAAAAAAAAAAAA.....
.....RRRRRRRRRR.....RRRR.....RRRR.....
DDDDDD.....DDDDDDDDDDDD.....
.....ddddd.....
305 310 315 320 325 330 335 340 345 350 355 360 365 370 375
RRVPGVAFPLVRSASSETSEYRPFMCAYFGGKRYFTLSHLQNHRSRYKGTGKPYQCDFMDCERAFERSGQQLKRWQR
AAAAA.....AAAAAAAAAAAAA.....AAAA.AAAAAA.....
.....RRRR.....RRRR.....
.....DDDDDD.....
.....
380 385 390 395 400 405 410 415 420 425 430 435 440 445 450
RHTGVKFFQCKTCQKTFERSCHLKHTRCHTGTSTSEKPFSCRWPSQKKTARSDELVRHHNMHCQNMTKLQLAL
.....AAAA.AAAA.AA.....AAAA.....AAA.....AAAAA.....AAA.....
.....RRRR.RRRR.....
.....
.....ddddd.....

FIG. 8A

5 10 15 20 25 30 35 40 45 50 55 60 65 70 75
MGSDVROLNALLPVSLSLGGGGGCLPVSQAACNAZVLDERPQGASAYGSLGGZAPPPAPPPPPPPPPHSTFKQE
.....AAAAAAAAAAAAAAAA.....AAAAA.....AAAAAAAAAAAAA.....
.....RRRR.....
.....
.....
80 85 90 95 100 105 110 115 120 125 130 135 140 145 150
PSWGGAEZHEEQCLSAFTLFTSGQFTGTAGACRYGZFGZPPPSQASSGQARMFNAPYLPSCLSQPTERNQGYT
.....AAAA.....AAA.....AAAAA.....
.....RRRR.....RRRR.....
.....DDDDDDDDDD.....
.....
155 160 165 170 175 180 185 190 195 200 205 210 215 220 225
TVTFDGAZSYGHTPSHBAQFPNHSFTHEDFMGQQGSLJZQQYSV999VYGCHTPTDSCGTGSAZLLRTFYSSCN
.....AAAA.....AAAAA.....AA
.....RRRR.....
.....DDDDDDDDDDDDDDDDDDDD.....
.....
230 235 240 245 250 255 260 265 270 275 280 285 290 295 300
LYQMTSGLQCTNQNLCATLKGMFAGSSSSVHTTQSNHRTGYSSNHTAPILGTAQVRIHTHGVTRGIQOV
AAAAAAAA.....AAA.AAA.....AAAAAAAAAAAA
.....RRRRRRRRRR.....RRRR.....RRRR.....
DDDDDD.....DDDDDDDDDDDD.....
.....
305 310 315 320 325 330 335 340 345 350 355 360 365 370 375
RRVSGVAPTQVRSASZTSEVRPFMGAYPGCNKATYTLHLQNHSHKHTGKPYCCDTHDCERRFSRSQQLKRWQR
AAAAA.AAAAAAAAAA.....AAAA.AAAAAAAA.
.....RRRR.....RRRR.....
.....DDDDDDDDDDDD.....
.....
380 385 390 395 400 405 410 415 420 425 430 435 440 445 450
RHTGVKZFFQKTCQRKTFRSCHLKTHTTCHTCKTSEKFTSCRWHSQCQKHTARSDELVRHNNMHQRNMTXLEVAL
.....AAAA.AAAA.AA.....AAAA.....AA.....AAAAA.....AAAA.....
.....RRRR.RRRR.....
.....
.....DDDDDDDDDDDD.....

FIG. 8B

A**B****FIG. 9A and 9B**





FIG. 10A

*FIG. 10B*




FIG. 10C






FIG. 10D

A**B****FIG. 11A and 11B**

16/17




FIG. 12A and 12B




FIG. 13A-13C

SEQUENCE LISTING

<110> Corixa Corporation et al.

<120> COMPOSITIONS AND METHODS FOR WT1
SPECIFIC IMMUNOTHERAPY

<130> 210121.533PC

<140> PCT

<141> 2000-10-04

<160> 357

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 17

<212> PRT

<213> Homo sapien

<400> 1

Arg	Asp	Leu	Asn	Ala	Leu	Leu	Pro	Ala	Val	Pro	Ser	Leu	Gly	Gly	Gly
1				5					10					15	
Gly															

<210> 2

<211> 23

<212> PRT

<213> Homo sapien

<400> 2

Pro	Ser	Gln	Ala	Ser	Ser	Gly	Gln	Ala	Arg	Met	Phe	Pro	Asn	Ala	Pro
1				5					10					15	
Tyr	Leu	Pro	Ser	Cys	Leu	Glu									
			20												

<210> 3

<211> 23

<212> PRT

<213> Mus musculus

<400> 3

Pro	Ser	Gln	Ala	Ser	Ser	Gly	Gln	Ala	Arg	Met	Phe	Pro	Asn	Ala	Pro
1				5					10					15	
Tyr	Leu	Pro	Ser	Cys	Leu	Glu									
			20												

<210> 4

<211> 19

<212> PRT

<213> Homo sapien

<400> 4

Gly	Ala	Thr	Leu	Lys	Gly	Val	Ala	Ala	Gly	Ser	Ser	Ser	Ser	Val	Lys
1				5					10					15	

Trp Thr Glu

<210> 5
<211> 22
<212> DNA
<213> Homo sapien

<400> 5
gagagtcaga cttgaaagca gt 22

<210> 6
<211> 20
<212> DNA
<213> Homo sapien

<400> 6
ctgagcctca gcaaattgggc 20

<210> 7
<211> 27
<212> DNA
<213> Homo sapien

<400> 7
gagcatgcat gggctccgac gtgcggg 27

<210> 8
<211> 25
<212> DNA
<213> Homo sapien

<400> 8
ggggtaccca ctgaacggtc cccga 25

<210> 9
<211> 18
<212> DNA
<213> Mus musculus

<400> 9
tccgagccgc acctcatg 18

<210> 10
<211> 18
<212> DNA
<213> Mus musculus

<400> 10
gcctgggatg ctggactg 18

<210> 11
<211> 27
<212> DNA
<213> Mus musculus

<400> 11
gagcatgcga tgggttccga cgtgcgg 27

<210> 12
 <211> 29
 <212> DNA
 <213> Mus musculus

<400> 12
 ggggtacctc aaagcgccac gtggagttt

29

<210> 13
 <211> 17
 <212> PRT
 <213> Mus musculus

<400> 13
 Arg Asp Leu Asn Ala Leu Leu Pro Ala Val Ser Ser Leu Gly Gly Gly
 1 5 10 15
 Gly

<210> 14
 <211> 19
 <212> PRT
 <213> Mus musculus

<400> 14
 Gly Ala Thr Leu Lys Gly Met Ala Ala Gly Ser Ser Ser Ser Val Lys
 1 5 10 15
 Trp Thr Glu

<210> 15
 <211> 15
 <212> PRT
 <213> Homo sapien

<400> 15
 Arg Ile His Thr His Gly Val Phe Arg Gly Ile Gln Asp Val Arg
 1 5 10 15

<210> 16
 <211> 15
 <212> PRT
 <213> Mus musculus

<400> 16
 Arg Ile His Thr His Gly Val Phe Arg Gly Ile Gln Asp Val Arg
 1 5 10 15

<210> 17
 <211> 14
 <212> PRT
 <213> Mus musculus

<400> 17
 Val Arg Arg Val Ser Gly Val Ala Pro Thr Leu Val Arg Ser
 1 5 10

<210> 18
 <211> 14
 <212> PRT
 <213> Homo sapien

<400> 18
 Val Arg Arg Val Pro Gly Val Ala Pro Thr Leu Val Arg Ser
 1 5 10

<210> 19
 <211> 15
 <212> PRT
 <213> Homo sapien

<400> 19
 Cys Gln Lys Lys Phe Ala Arg Ser Asp Glu Leu Val Arg His His
 1 5 10 15

<210> 20
 <211> 15
 <212> PRT
 <213> Mus musculus

<400> 20
 Cys Gln Lys Lys Phe Ala Arg Ser Asp Glu Leu Val Arg His His
 1 5 10 15

<210> 21
 <211> 21
 <212> DNA
 <213> Mus musculus

<400> 21
 cccaggctgc aataagagat a 21

<210> 22
 <211> 21
 <212> DNA
 <213> Mus musculus

<400> 22
 atgttgatgac ggcggaccaa t 21

<210> 23
 <211> 20
 <212> DNA
 <213> Homo sapien

<400> 23
 gtggggcgcc ccaggcacca 20

<210> 24
 <211> 24
 <212> DNA
 <213> Homo sapien

<400> 24
 gtccttaatg ctacgcacga ttcc 24

<210> 25
<211> 21
<212> DNA
<213> Homo sapien

<400> 25
ggcatctgag accagtgaga a 21

<210> 26
<211> 21
<212> DNA
<213> Homo sapien

<400> 26
gctgtccac ttacagatgc a 21

<210> 27
<211> 21
<212> DNA
<213> Homo sapien

<400> 27
tcaaagcgcc agctggagtt t 21

<210> 28
<211> 9
<212> PRT
<213> Homo sapien

<400> 28
Ala Ala Gly Ser Ser Ser Val Lys
1 5

<210> 29
<211> 9
<212> PRT
<213> Homo sapien

<400> 29
Ala Ala Gln Phe Pro Asn His Ser Phe
1 5

<210> 30
<211> 9
<212> PRT
<213> Homo sapien

<400> 30
Ala Glu Pro His Glu Glu Gln Cys Leu
1 5

<210> 31
<211> 9
<212> PRT
<213> Homo sapien

<400> 31

Ala Gly Ala Cys Arg Tyr Gly Pro Phe
1 5

<210> 32
<211> 9
<212> PRT
<213> Homo sapien

<400> 32
Ala Gly Ser Ser Ser Ser Val Lys Trp
1 5

<210> 33
<211> 9
<212> PRT
<213> Homo sapien

<400> 33
Ala Ile Arg Asn Gln Gly Tyr Ser Thr
1 5

<210> 34
<211> 9
<212> PRT
<213> Homo sapien

<400> 34
Ala Leu Leu Pro Ala Val Pro Ser Leu
1 5

<210> 35
<211> 9
<212> PRT
<213> Homo sapien

<400> 35
Ala Leu Leu Pro Ala Val Ser Ser Leu
1 5

<210> 36
<211> 9
<212> PRT
<213> Homo sapien

<400> 36
Ala Gln Phe Pro Asn His Ser Phe Lys
1 5

<210> 37
<211> 9
<212> PRT
<213> Homo sapien

<400> 37
Ala Gln Trp Ala Pro Val Leu Asp Phe
1 5

<210> 38

<211> 9
<212> PRT
<213> Homo sapien

<400> 38
Ala Arg Met Phe Pro Asn Ala Pro Tyr
1 5

<210> 39
<211> 9
<212> PRT
<213> Homo sapien

<400> 39
Ala Arg Ser Asp Glu Leu Val Arg His
1 5

<210> 40
<211> 9
<212> PRT
<213> Homo sapien

<400> 40
Ala Ser Ser Gly Gln Ala Arg Met Phe
1 5

<210> 41
<211> 9
<212> PRT
<213> Homo sapien

<400> 41
Ala Tyr Gly Ser Leu Gly Gly Pro Ala
1 5

<210> 42
<211> 9
<212> PRT
<213> Homo sapien

<400> 42
Ala Tyr Pro Gly Cys Asn Lys Arg Tyr
1 5

<210> 43
<211> 9
<212> PRT
<213> Homo sapien

<400> 43
Cys Ala Leu Pro Val Ser Gly Ala Ala
1 5

<210> 44
<211> 9
<212> PRT
<213> Homo sapien

<400> 44
Cys Ala Tyr Pro Gly Cys Asn Lys Arg
1 5

<210> 45
<211> 9
<212> PRT
<213> Homo sapien

<400> 45
Cys His Thr Pro Thr Asp Ser Cys Thr
1 5

<210> 46
<211> 9
<212> PRT
<213> Homo sapien

<400> 46
Cys Lys Thr Cys Gln Arg Lys Phe Ser
1 5

<210> 47
<211> 9
<212> PRT
<213> Homo sapien

<400> 47
Cys Leu Glu Ser Gln Pro Ala Ile Arg
1 5

<210> 48
<211> 9
<212> PRT
<213> Homo sapien

<400> 48
Cys Leu Ser Ala Phe Thr Val His Phe
1 5

<210> 49
<211> 9
<212> PRT
<213> Homo sapien

<400> 49
Cys Met Thr Trp Asn Gln Met Asn Leu
1 5

<210> 50
<211> 9
<212> PRT
<213> Homo sapien

<400> 50
Cys Arg Trp Pro Ser Cys Gln Lys Lys
1 5

<210> 51
<211> 9
<212> PRT
<213> Homo sapien

<400> 51
Cys Arg Tyr Gly Pro Phe Gly Pro Pro
1 5

<210> 52
<211> 9
<212> PRT
<213> Homo sapien

<400> 52
Cys Thr Gly Ser Gln Ala Leu Leu Leu
1 5

<210> 53
<211> 9
<212> PRT
<213> Homo sapien

<400> 53
Asp Glu Leu Val Arg His His Asn Met
1 5

<210> 54
<211> 9
<212> PRT
<213> Homo sapien

<400> 54
Asp Phe Ala Pro Pro Gly Ala Ser Ala
1 5

<210> 55
<211> 9
<212> PRT
<213> Homo sapien

<400> 55
Asp Phe Lys Asp Cys Glu Arg Arg Phe
1 5

<210> 56
<211> 9
<212> PRT
<213> Homo sapien

<400> 56
Asp Gly Thr Pro Ser Tyr Gly His Thr
1 5

<210> 57
<211> 9
<212> PRT
<213> Homo sapien

<400> 57
Asp His Leu Lys Thr His Thr Arg Thr
1 5

<210> 58
<211> 9
<212> PRT
<213> Homo sapien

<400> 58
Asp Leu Asn Ala Leu Leu Pro Ala Val
1 5

<210> 59
<211> 9
<212> PRT
<213> Homo sapien

<400> 59
Asp Pro Met Gly Gln Gln Gly Ser Leu
1 5

<210> 60
<211> 9
<212> PRT
<213> Homo sapien

<400> 60
Asp Gln Leu Lys Arg His Gln Arg Arg
1 5

<210> 61
<211> 9
<212> PRT
<213> Homo sapien

<400> 61
Asp Ser Cys Thr Gly Ser Gln Ala Leu
1 5

<210> 62
<211> 9
<212> PRT
<213> Homo sapien

<400> 62
Asp Val Arg Asp Leu Asn Ala Leu Leu
1 5

<210> 63
<211> 9
<212> PRT
<213> Homo sapien

<400> 63
Asp Val Arg Arg Val Pro Gly Val Ala
1 5

<210> 64
<211> 9
<212> PRT
<213> Homo sapien

<400> 64
Glu Asp Pro Met Gly Gln Gln Gly Ser
1 5

<210> 65
<211> 9
<212> PRT
<213> Homo sapien

<400> 65
Glu Glu Gln Cys Leu Ser Ala Phe Thr
1 5

<210> 66
<211> 9
<212> PRT
<213> Homo sapien

<400> 66
Glu Lys Pro Tyr Gln Cys Asp Phe Lys
1 5

<210> 67
<211> 9
<212> PRT
<213> Homo sapien

<400> 67
Glu Lys Arg Pro Phe Met Cys Ala Tyr
1 5

<210> 68
<211> 9
<212> PRT
<213> Homo sapien

<400> 68
Glu Pro His Glu Glu Gln Cys Leu Ser
1 5

<210> 69
<211> 9
<212> PRT
<213> Homo sapien

<400> 69
Glu Gln Cys Leu Ser Ala Phe Thr Val
1 5

<210> 70
<211> 9
<212> PRT

<213> Homo sapien

<400> 70

Glu Ser Asp Asn His Thr Ala Pro Ile
1 5

<210> 71

<211> 9

<212> PRT

<213> Homo sapien

<400> 71

Glu Ser Asp Asn His Thr Thr Pro Ile
1 5

<210> 72

<211> 9

<212> PRT

<213> Homo sapien

<400> 72

Glu Ser Gln Pro Ala Ile Arg Asn Gln
1 5

<210> 73

<211> 9

<212> PRT

<213> Homo sapien

<400> 73

Glu Thr Ser Glu Lys Arg Pro Phe Met
1 5

<210> 74

<211> 9

<212> PRT

<213> Homo sapien

<400> 74

Phe Ala Pro Pro Gly Ala Ser Ala Tyr
1 5

<210> 75

<211> 9

<212> PRT

<213> Homo sapien

<400> 75

Phe Ala Arg Ser Asp Glu Leu Val Arg
1 5

<210> 76

<211> 9

<212> PRT

<213> Homo sapien

<400> 76

Phe Gly Pro Pro Pro Pro Ser Gln Ala

1 5

<210> 77
<211> 9
<212> PRT
<213> Homo sapien

<400> 77
Phe Lys Asp Cys Glu Arg Arg Phe Ser
1 5

<210> 78
<211> 9
<212> PRT
<213> Homo sapien

<400> 78
Phe Lys Leu Ser His Leu Gln Met His
1 5

<210> 79
<211> 9
<212> PRT
<213> Homo sapien

<400> 79
Phe Pro Asn Ala Pro Tyr Leu Pro Ser
1 5

<210> 80
<211> 9
<212> PRT
<213> Homo sapien

<400> 80
Phe Gln Cys Lys Thr Cys Gln Arg Lys
1 5

<210> 81
<211> 9
<212> PRT
<213> Homo sapien

<400> 81
Phe Arg Gly Ile Gln Asp Val Arg Arg
1 5

<210> 82
<211> 9
<212> PRT
<213> Homo sapien

<400> 82
Phe Ser Gly Gln Phe Thr Gly Thr Ala
1 5

<210> 83
<211> 9

<212> PRT
<213> Homo sapien

<400> 83
Phe Ser Arg Ser Asp Gln Leu Lys Arg
1 5

<210> 84
<211> 9
<212> PRT
<213> Homo sapien

<400> 84
Phe Thr Gly Thr Ala Gly Ala Cys Arg
1 5

<210> 85
<211> 9
<212> PRT
<213> Homo sapien

<400> 85
Phe Thr Val His Phe Ser Gly Gln Phe
1 5

<210> 86
<211> 9
<212> PRT
<213> Homo sapien

<400> 86
Gly Ala Ala Gln Trp Ala Pro Val Leu
1 5

<210> 87
<211> 9
<212> PRT
<213> Homo sapien

<400> 87
Gly Ala Glu Pro His Glu Glu Gln Cys
1 5

<210> 88
<211> 9
<212> PRT
<213> Homo sapien

<400> 88
Gly Ala Thr Leu Lys Gly Val Ala Ala
1 5

<210> 89
<211> 9
<212> PRT
<213> Homo sapien

<400> 89

Gly Cys Ala Leu Pro Val Ser Gly Ala
1 5

<210> 90
<211> 9
<212> PRT
<213> Homo sapien

<400> 90
Gly Cys Asn Lys Arg Tyr Phe Lys Leu
1 5

<210> 91
<211> 9
<212> PRT
<213> Homo sapien

<400> 91
Gly Glu Lys Pro Tyr Gln Cys Asp Phe
1 5

<210> 92
<211> 9
<212> PRT
<213> Homo sapien

<400> 92
Gly Gly Gly Gly Cys Ala Leu Pro Val
1 5

<210> 93
<211> 9
<212> PRT
<213> Homo sapien

<400> 93
Gly Gly Pro Ala Pro Pro Pro Ala Pro
1 5

<210> 94
<211> 9
<212> PRT
<213> Homo sapien

<400> 94
Gly His Thr Pro Ser His His Ala Ala
1 5

<210> 95
<211> 9
<212> PRT
<213> Homo sapien

<400> 95
Gly Lys Thr Ser Glu Lys Pro Phe Ser
1 5

<210> 96

<211> 9
<212> PRT
<213> Homo sapien

<400> 96
Gly Pro Phe Gly Pro Pro Pro Pro Ser
1 5

<210> 97
<211> 9
<212> PRT
<213> Homo sapien

<400> 97
Gly Pro Pro Pro Pro Ser Gln Ala Ser
1 5

<210> 98
<211> 9
<212> PRT
<213> Homo sapien

<400> 98
Gly Gln Ala Arg Met Phe Pro Asn Ala
1 5

<210> 99
<211> 9
<212> PRT
<213> Homo sapien

<400> 99
Gly Gln Phe Thr Gly Thr Ala Gly Ala
1 5

<210> 100
<211> 9
<212> PRT
<213> Homo sapien

<400> 100
Gly Gln Ser Asn His Ser Thr Gly Tyr
1 5

<210> 101
<211> 9
<212> PRT
<213> Homo sapien

<400> 101
Gly Ser Asp Val Arg Asp Leu Asn Ala
1 5

<210> 102
<211> 9
<212> PRT
<213> Homo sapien

<400> 102
Gly Ser Gln Ala Leu Leu Leu Arg Thr
1 5

<210> 103
<211> 9
<212> PRT
<213> Homo sapien

<400> 103
Gly Val Phe Arg Gly Ile Gln Asp Val
1 5

<210> 104
<211> 9
<212> PRT
<213> Homo sapien

<400> 104
Gly Val Lys Pro Phe Gln Cys Lys Thr
1 5

<210> 105
<211> 9
<212> PRT
<213> Homo sapien

<400> 105
Gly Tyr Glu Ser Asp Asn His Thr Ala
1 5

<210> 106
<211> 9
<212> PRT
<213> Homo sapien

<400> 106
Gly Tyr Glu Ser Asp Asn His Thr Thr
1 5

<210> 107
<211> 9
<212> PRT
<213> Homo sapien

<400> 107
His Glu Glu Gln Cys Leu Ser Ala Phe
1 5

<210> 108
<211> 9
<212> PRT
<213> Homo sapien

<400> 108
His His Asn Met His Gln Arg Asn Met
1 5

<210> 109
<211> 9
<212> PRT
<213> Homo sapien

<400> 109
His Gln Arg Arg His Thr Gly Val Lys
1 5

<210> 110
<211> 9
<212> PRT
<213> Homo sapien

<400> 110
His Ser Phe Lys His Glu Asp Pro Met
1 5

<210> 111
<211> 9
<212> PRT
<213> Homo sapien

<400> 111
His Ser Arg Lys His Thr Gly Glu Lys
1 5

<210> 112
<211> 9
<212> PRT
<213> Homo sapien

<400> 112
His Thr Gly Glu Lys Pro Tyr Gln Cys
1 5

<210> 113
<211> 9
<212> PRT
<213> Homo sapien

<400> 113
His Thr His Gly Val Phe Arg Gly Ile
1 5

<210> 114
<211> 9
<212> PRT
<213> Homo sapien

<400> 114
His Thr Arg Thr His Thr Gly Lys Thr
1 5

<210> 115
<211> 9
<212> PRT
<213> Homo sapien

<400> 115
His Thr Thr Pro Ile Leu Cys Gly Ala
1 5

<210> 116
<211> 9
<212> PRT
<213> Homo sapien

<400> 116
Ile Leu Cys Gly Ala Gln Tyr Arg Ile
1 5

<210> 117
<211> 9
<212> PRT
<213> Homo sapien

<400> 117
Ile Arg Asn Gln Gly Tyr Ser Thr Val
1 5

<210> 118
<211> 9
<212> PRT
<213> Homo sapien

<400> 118
Lys Asp Cys Glu Arg Arg Phe Ser Arg
1 5

<210> 119
<211> 9
<212> PRT
<213> Homo sapien

<400> 119
Lys Phe Ala Arg Ser Asp Glu Leu Val
1 5

<210> 120
<211> 9
<212> PRT
<213> Homo sapien

<400> 120
Lys Phe Ser Arg Ser Asp His Leu Lys
1 5

<210> 121
<211> 9
<212> PRT
<213> Homo sapien

<400> 121
Lys His Glu Asp Pro Met Gly Gln Gln
1 5

<210> 122
<211> 9
<212> PRT
<213> Homo sapien

<400> 122
Lys Lys Phe Ala Arg Ser Asp Glu Leu
1 5

<210> 123
<211> 9
<212> PRT
<213> Homo sapien

<400> 123
Lys Pro Phe Ser Cys Arg Trp Pro Ser
1 5

<210> 124
<211> 9
<212> PRT
<213> Homo sapien

<400> 124
Lys Pro Tyr Gln Cys Asp Phe Lys Asp
1 5

<210> 125
<211> 9
<212> PRT
<213> Homo sapien

<400> 125
Lys Gln Glu Pro Ser Trp Gly Gly Ala
1 5

<210> 126
<211> 9
<212> PRT
<213> Homo sapien

<400> 126
Lys Arg His Gln Arg Arg His Thr Gly
1 5

<210> 127
<211> 9
<212> PRT
<213> Homo sapien

<400> 127
Lys Arg Tyr Phe Lys Leu Ser His Leu
1 5

<210> 128
<211> 9
<212> PRT

<213> Homo sapien

<400> 128
Lys Thr Cys Gln Arg Lys Phe Ser Arg
1 5

<210> 129
<211> 9
<212> PRT
<213> Homo sapien

<400> 129
Lys Thr Ser Glu Lys Pro Phe Ser Cys
1 5

<210> 130
<211> 9
<212> PRT
<213> Homo sapien

<400> 130
Leu Asp Phe Ala Pro Pro Gly Ala Ser
1 5

<210> 131
<211> 9
<212> PRT
<213> Homo sapien

<400> 131
Leu Glu Cys Met Thr Trp Asn Gln Met
1 5

<210> 132
<211> 9
<212> PRT
<213> Homo sapien

<400> 132
Leu Glu Ser Gln Pro Ala Ile Arg Asn
1 5

<210> 133
<211> 9
<212> PRT
<213> Homo sapien

<400> 133
Leu Gly Ala Thr Leu Lys Gly Val Ala
1 5

<210> 134
<211> 9
<212> PRT
<213> Homo sapien

<400> 134
Leu Gly Gly Gly Gly Gly Cys Ala Leu

1 5

<210> 135
<211> 9
<212> PRT
<213> Homo sapien

<400> 135
Leu Lys Gly Val Ala Ala Gly Ser Ser
1 5

<210> 136
<211> 9
<212> PRT
<213> Homo sapien

<400> 136
Leu Lys Arg His Gln Arg Arg His Thr
1 5

<210> 137
<211> 9
<212> PRT
<213> Homo sapien

<400> 137
Leu Lys Thr His Thr Arg Thr His Thr
1 5

<210> 138
<211> 9
<212> PRT
<213> Homo sapien

<400> 138
Leu Pro Val Ser Gly Ala Ala Gln Trp
1 5

<210> 139
<211> 9
<212> PRT
<213> Homo sapien

<400> 139
Leu Gln Met His Ser Arg Lys His Thr
1 5

<210> 140
<211> 9
<212> PRT
<213> Homo sapien

<400> 140
Leu Arg Thr Pro Tyr Ser Ser Asp Asn
1 5

<210> 141
<211> 9

<212> PRT
<213> Homo sapien

<400> 141
Leu Ser His Leu Gln Met His Ser Arg
1 5

<210> 142
<211> 9
<212> PRT
<213> Homo sapien

<400> 142
Met Cys Ala Tyr Pro Gly Cys Asn Lys
1 5

<210> 143
<211> 9
<212> PRT
<213> Homo sapien

<400> 143
Met His Gln Arg Asn Met Thr Lys Leu
1 5

<210> 144
<211> 9
<212> PRT
<213> Homo sapien

<400> 144
Asn Ala Pro Tyr Leu Pro Ser Cys Leu
1 5

<210> 145
<211> 9
<212> PRT
<213> Homo sapien

<400> 145
Asn Lys Arg Tyr Phe Lys Leu Ser His
1 5

<210> 146
<211> 9
<212> PRT
<213> Homo sapien

<400> 146
Asn Leu Gly Ala Thr Leu Lys Gly Val
1 5

<210> 147
<211> 9
<212> PRT
<213> Homo sapien

<400> 147

Asn Leu Tyr Gln Met Thr Ser Gln Leu
1 5

<210> 148
<211> 9
<212> PRT
<213> Homo sapien

<400> 148
Asn Met His Gln Arg Asn Met Thr Lys
1 5

<210> 149
<211> 9
<212> PRT
<213> Homo sapien

<400> 149
Asn Met Thr Lys Leu Gln Leu Ala Leu
1 5

<210> 150
<211> 9
<212> PRT
<213> Homo sapien

<400> 150
Asn Gln Gly Tyr Ser Thr Val Thr Phe
1 5

<210> 151
<211> 9
<212> PRT
<213> Homo sapien

<400> 151
Asn Gln Met Asn Leu Gly Ala Thr Leu
1 5

<210> 152
<211> 9
<212> PRT
<213> Homo sapien

<400> 152
Pro Ala Ile Arg Asn Gln Gly Tyr Ser
1 5

<210> 153
<211> 9
<212> PRT
<213> Homo sapien

<400> 153
Pro Gly Ala Ser Ala Tyr Gly Ser Leu
1 5

<210> 154

<211> 9
<212> PRT
<213> Homo sapien

<400> 154
Pro His Glu Glu Gln Cys Leu Ser Ala
1 5

<210> 155
<211> 9
<212> PRT
<213> Homo sapien

<400> 155
Pro Ile Leu Cys Gly Ala Gln Tyr Arg
1 5

<210> 156
<211> 9
<212> PRT
<213> Homo sapien

<400> 156
Pro Pro Pro Pro His Ser Phe Ile Lys
1 5

<210> 157
<211> 9
<212> PRT
<213> Homo sapien

<400> 157
Pro Pro Pro Pro Pro His Ser Phe Ile
1 5

<210> 158
<211> 9
<212> PRT
<213> Homo sapien

<400> 158
Pro Pro Pro Pro Pro Pro His Ser Phe
1 5

<210> 159
<211> 9
<212> PRT
<213> Homo sapien

<400> 159
Pro Ser Cys Gln Lys Lys Phe Ala Arg
1 5

<210> 160
<211> 9
<212> PRT
<213> Homo sapien

<400> 160
Gln Ala Leu Leu Leu Arg Thr Pro Tyr
1 5

<210> 161
<211> 9
<212> PRT
<213> Homo sapien

<400> 161
Gln Ala Ser Ser Gly Gln Ala Arg Met
1 5

<210> 162
<211> 9
<212> PRT
<213> Homo sapien

<400> 162
Gln Cys Asp Phe Lys Asp Cys Glu Arg
1 5

<210> 163
<211> 9
<212> PRT
<213> Homo sapien

<400> 163
Gln Cys Lys Thr Cys Gln Arg Lys Phe
1 5

<210> 164
<211> 9
<212> PRT
<213> Homo sapien

<400> 164
Gln Asp Val Arg Arg Val Pro Gly Val
1 5

<210> 165
<211> 9
<212> PRT
<213> Homo sapien

<400> 165
Gln Phe Thr Gly Thr Ala Gly Ala Cys
1 5

<210> 166
<211> 9
<212> PRT
<213> Homo sapien

<400> 166
Gln Gly Ser Leu Gly Glu Gln Gln Tyr
1 5

<210> 167
<211> 9
<212> PRT
<213> Homo sapien

<400> 167
Gln Leu Glu Cys Met Thr Trp Asn Gln
1 5

<210> 168
<211> 9
<212> PRT
<213> Homo sapien

<400> 168
Gln Met Asn Leu Gly Ala Thr Leu Lys
1 5

<210> 169
<211> 9
<212> PRT
<213> Homo sapien

<400> 169
Gln Met Thr Ser Gln Leu Glu Cys Met
1 5

<210> 170
<211> 9
<212> PRT
<213> Homo sapien

<400> 170
Gln Pro Ala Ile Arg Asn Gln Gly Tyr
1 5

<210> 171
<211> 9
<212> PRT
<213> Homo sapien

<400> 171
Gln Gln Tyr Ser Val Pro Pro Pro Val
1 5

<210> 172
<211> 9
<212> PRT
<213> Homo sapien

<400> 172
Gln Arg Lys Phe Ser Arg Ser Asp His
1 5

<210> 173
<211> 9
<212> PRT
<213> Homo sapien

<400> 173
Gln Arg Asn Met Thr Lys Leu Gln Leu
1 5

<210> 174
<211> 9
<212> PRT
<213> Homo sapien

<400> 174
Gln Trp Ala Pro Val Leu Asp Phe Ala
1 5

<210> 175
<211> 9
<212> PRT
<213> Homo sapien

<400> 175
Gln Tyr Arg Ile His Thr His Gly Val
1 5

<210> 176
<211> 9
<212> PRT
<213> Homo sapien

<400> 176
Gln Tyr Ser Val Pro Pro Pro Val Tyr
1 5

<210> 177
<211> 9
<212> PRT
<213> Homo sapien

<400> 177
Arg Asp Leu Asn Ala Leu Leu Pro Ala
1 5

<210> 178
<211> 9
<212> PRT
<213> Homo sapien

<400> 178
Arg Phe Ser Arg Ser Asp Gln Leu Lys
1 5

<210> 179
<211> 9
<212> PRT
<213> Homo sapien

<400> 179
Arg Gly Ile Gln Asp Val Arg Arg Val
1 5

<210> 180
<211> 9
<212> PRT
<213> Homo sapien

<400> 180
Arg His His Asn Met His Gln Arg Asn
1 5

<210> 181
<211> 9
<212> PRT
<213> Homo sapien

<400> 181
Arg His Gln Arg Arg His Thr Gly Val
1 5

<210> 182
<211> 9
<212> PRT
<213> Homo sapien

<400> 182
Arg Ile His Thr His Gly Val Phe Arg
1 5

<210> 183
<211> 9
<212> PRT
<213> Homo sapien

<400> 183
Arg Lys Phe Ser Arg Ser Asp His Leu
1 5

<210> 184
<211> 9
<212> PRT
<213> Homo sapien

<400> 184
Arg Lys His Thr Gly Glu Lys Pro Tyr
1 5

<210> 185
<211> 9
<212> PRT
<213> Homo sapien

<400> 185
Arg Met Phe Pro Asn Ala Pro Tyr Leu
1 5

<210> 186
<211> 9
<212> PRT

<213> Homo sapien

<400> 186

Arg Asn Met Thr Lys Leu Gln Leu Ala
1 5

<210> 187

<211> 9

<212> PRT

<213> Homo sapien

<400> 187

Arg Arg Phe Ser Arg Ser Asp Gln Leu
1 5

<210> 188

<211> 9

<212> PRT

<213> Homo sapien

<400> 188

Arg Arg His Thr Gly Val Lys Pro Phe
1 5

<210> 189

<211> 9

<212> PRT

<213> Homo sapien

<400> 189

Arg Arg Val Pro Gly Val Ala Pro Thr
1 5

<210> 190

<211> 9

<212> PRT

<213> Homo sapien

<400> 190

Arg Ser Ala Ser Glu Thr Ser Glu Lys
1 5

<210> 191

<211> 9

<212> PRT

<213> Homo sapien

<400> 191

Arg Ser Asp Glu Leu Val Arg His His
1 5

<210> 192

<211> 9

<212> PRT

<213> Homo sapien

<400> 192

Arg Ser Asp His Leu Lys Thr His Thr

1 5

<210> 193
<211> 9
<212> PRT
<213> Homo sapien

<400> 193
Arg Ser Asp Gln Leu Lys Arg His Gln
1 5

<210> 194
<211> 9
<212> PRT
<213> Homo sapien

<400> 194
Arg Thr Pro Tyr Ser Ser Asp Asn Leu
1 5

<210> 195
<211> 9
<212> PRT
<213> Homo sapien

<400> 195
Arg Val Pro Gly Val Ala Pro Thr Leu
1 5

<210> 196
<211> 9
<212> PRT
<213> Homo sapien

<400> 196
Arg Trp Pro Ser Cys Gln Lys Lys Phe
1 5

<210> 197
<211> 9
<212> PRT
<213> Homo sapien

<400> 197
Ser Ala Ser Glu Thr Ser Glu Lys Arg
1 5

<210> 198
<211> 9
<212> PRT
<213> Homo sapien

<400> 198
Ser Cys Leu Glu Ser Gln Pro Ala Ile
1 5

<210> 199
<211> 9

<212> PRT
<213> Homo sapien

<400> 199
Ser Cys Leu Glu Ser Gln Pro Thr Ile
1 5

<210> 200
<211> 9
<212> PRT
<213> Homo sapien

<400> 200
Ser Cys Gln Lys Lys Phe Ala Arg Ser
1 5

<210> 201
<211> 9
<212> PRT
<213> Homo sapien

<400> 201
Ser Cys Arg Trp Pro Ser Cys Gln Lys
1 5

<210> 202
<211> 9
<212> PRT
<213> Homo sapien

<400> 202
Ser Cys Thr Gly Ser Gln Ala Leu Leu
1 5

<210> 203
<211> 9
<212> PRT
<213> Homo sapien

<400> 203
Ser Asp Glu Leu Val Arg His His Asn
1 5

<210> 204
<211> 9
<212> PRT
<213> Homo sapien

<400> 204
Ser Asp Asn His Thr Thr Pro Ile Leu
1 5

<210> 205
<211> 9
<212> PRT
<213> Homo sapien

<400> 205

Ser Asp Asn Leu Tyr Gln Met Thr Ser
1 5

<210> 206
<211> 9
<212> PRT
<213> Homo sapien

<400> 206

Ser Asp Val Arg Asp Leu Asn Ala Leu
1 5

<210> 207
<211> 9
<212> PRT
<213> Homo sapien

<400> 207

Ser Glu Lys Pro Phe Ser Cys Arg Trp
1 5

<210> 208
<211> 9
<212> PRT
<213> Homo sapien

<400> 208

Ser Glu Lys Arg Pro Phe Met Cys Ala
1 5

<210> 209
<211> 9
<212> PRT
<213> Homo sapien

<400> 209

Ser Glu Thr Ser Glu Lys Arg Pro Phe
1 5

<210> 210
<211> 9
<212> PRT
<213> Homo sapien

<400> 210

Ser Phe Ile Lys Gln Glu Pro Ser Trp
1 5

<210> 211
<211> 9
<212> PRT
<213> Homo sapien

<400> 211

Ser Gly Ala Ala Gln Trp Ala Pro Val
1 5

<210> 212

<211> 9
<212> PRT
<213> Homo sapien

<400> 212
Ser Gly Gln Ala Arg Met Phe Pro Asn
1 5

<210> 213
<211> 9
<212> PRT
<213> Homo sapien

<400> 213
Ser His His Ala Ala Gln Phe Pro Asn
1 5

<210> 214
<211> 9
<212> PRT
<213> Homo sapien

<400> 214
Ser Leu Gly Glu Gln Gln Tyr Ser Val
1 5

<210> 215
<211> 9
<212> PRT
<213> Homo sapien

<400> 215
Ser Leu Gly Gly Gly Gly Gly Cys Ala
1 5

<210> 216
<211> 9
<212> PRT
<213> Homo sapien

<400> 216
Ser Gln Ala Ser Ser Gly Gln Ala Arg
1 5

<210> 217
<211> 9
<212> PRT
<213> Homo sapien

<400> 217
Ser Ser Asp Asn Leu Tyr Gln Met Thr
1 5

<210> 218
<211> 9
<212> PRT
<213> Homo sapien

<400> 218
Ser Val Pro Pro Pro Val Tyr Gly Cys
1 5

<210> 219
<211> 9
<212> PRT
<213> Homo sapien

<400> 219
Thr Cys Gln Arg Lys Phe Ser Arg Ser
1 5

<210> 220
<211> 9
<212> PRT
<213> Homo sapien

<400> 220
Thr Asp Ser Cys Thr Gly Ser Gln Ala
1 5

<210> 221
<211> 9
<212> PRT
<213> Homo sapien

<400> 221
Thr Glu Gly Gln Ser Asn His Ser Thr
1 5

<210> 222
<211> 9
<212> PRT
<213> Homo sapien

<400> 222
Thr Gly Lys Thr Ser Glu Lys Pro Phe
1 5

<210> 223
<211> 9
<212> PRT
<213> Homo sapien

<400> 223
Thr Gly Ser Gln Ala Leu Leu Leu Arg
1 5

<210> 224
<211> 9
<212> PRT
<213> Homo sapien

<400> 224
Thr Gly Thr Ala Gly Ala Cys Arg Tyr
1 5

<210> 225
<211> 9
<212> PRT
<213> Homo sapien

<400> 225
Thr Gly Tyr Glu Ser Asp Asn His Thr
1 5

<210> 226
<211> 9
<212> PRT
<213> Homo sapien

<400> 226
Thr Leu Val Arg Ser Ala Ser Glu Thr
1 5

<210> 227
<211> 9
<212> PRT
<213> Homo sapien

<400> 227
Thr Pro Ile Leu Cys Gly Ala Gln Tyr
1 5

<210> 228
<211> 9
<212> PRT
<213> Homo sapien

<400> 228
Thr Pro Ser His His Ala Ala Gln Phe
1 5

<210> 229
<211> 9
<212> PRT
<213> Homo sapien

<400> 229
Thr Pro Ser Tyr Gly His Thr Pro Ser
1 5

<210> 230
<211> 9
<212> PRT
<213> Homo sapien

<400> 230
Thr Pro Thr Asp Ser Cys Thr Gly Ser
1 5

<210> 231
<211> 9
<212> PRT
<213> Homo sapien

<400> 231
 Thr Pro Tyr Ser Ser Asp Asn Leu Tyr
 1 5

<210> 232
 <211> 9
 <212> PRT
 <213> Homo sapien

<400> 232
 Thr Ser Glu Lys Pro Phe Ser Cys Arg
 1 5

<210> 233
 <211> 9
 <212> PRT
 <213> Homo sapien

<400> 233
 Thr Ser Glu Lys Arg Pro Phe Met Cys
 1 5

<210> 234
 <211> 9
 <212> PRT
 <213> Homo sapien

<400> 234
 Thr Ser Gln Leu Glu Cys Met Thr Trp
 1 5

<210> 235
 <211> 9
 <212> PRT
 <213> Homo sapien

<400> 235
 Thr Val His Phe Ser Gly Gln Phe Thr
 1 5

<210> 236
 <211> 9
 <212> PRT
 <213> Homo sapien

<400> 236
 Val Ala Ala Gly Ser Ser Ser Ser Val
 1 5

<210> 237
 <211> 9
 <212> PRT
 <213> Homo sapien

<400> 237
 Val Ala Pro Thr Leu Val Arg Ser Ala
 1 5

<210> 238
<211> 9
<212> PRT
<213> Homo sapien

<400> 238
Val Phe Arg Gly Ile Gln Asp Val Arg
1 5

<210> 239
<211> 9
<212> PRT
<213> Homo sapien

<400> 239
Val Lys Pro Phe Gln Cys Lys Thr Cys
1 5

<210> 240
<211> 9
<212> PRT
<213> Homo sapien

<400> 240
Val Lys Trp Thr Glu Gly Gln Ser Asn
1 5

<210> 241
<211> 9
<212> PRT
<213> Homo sapien

<400> 241
Val Leu Asp Phe Ala Pro Pro Gly Ala
1 5

<210> 242
<211> 9
<212> PRT
<213> Homo sapien

<400> 242
Val Pro Gly Val Ala Pro Thr Leu Val
1 5

<210> 243
<211> 9
<212> PRT
<213> Homo sapien

<400> 243
Val Arg His His Asn Met His Gln Arg
1 5

<210> 244
<211> 9
<212> PRT

<213> Homo sapien

<400> 244

Val Thr Phe Asp Gly Thr Pro Ser Tyr
1 5

<210> 245

<211> 9

<212> PRT

<213> Homo sapien

<400> 245

Trp Asn Gln Met Asn Leu Gly Ala Thr
1 5

<210> 246

<211> 9

<212> PRT

<213> Homo sapien

<400> 246

Trp Pro Ser Cys Gln Lys Lys Phe Ala
1 5

<210> 247

<211> 9

<212> PRT

<213> Homo sapien

<400> 247

Trp Thr Glu Gly Gln Ser Asn His Ser
1 5

<210> 248

<211> 9

<212> PRT

<213> Homo sapien

<400> 248

Tyr Phe Lys Leu Ser His Leu Gln Met
1 5

<210> 249

<211> 9

<212> PRT

<213> Homo sapien

<400> 249

Tyr Gly His Thr Pro Ser His His Ala
1 5

<210> 250

<211> 9

<212> PRT

<213> Homo sapien

<400> 250

Tyr Pro Gly Cys Asn Lys Arg Tyr Phe

1 5

<210> 251
<211> 9
<212> PRT
<213> Homo sapien

<400> 251
Tyr Gln Met Thr Ser Gln Leu Glu Cys
1 5

<210> 252
<211> 9
<212> PRT
<213> Homo sapien

<400> 252
Tyr Arg Ile His Thr His Gly Val Phe
1 5

<210> 253
<211> 9
<212> PRT
<213> Homo sapien

<400> 253
Tyr Ser Ser Asp Asn Leu Tyr Gln Met
1 5

<210> 254
<211> 9
<212> PRT
<213> Mus musculus

<400> 254
Ala Glu Pro His Glu Glu Gln Cys Leu
1 5

<210> 255
<211> 9
<212> PRT
<213> Mus musculus

<400> 255
Ala Leu Leu Pro Ala Val Ser Ser Leu
1 5

<210> 256
<211> 9
<212> PRT
<213> Mus musculus

<400> 256
Ala Tyr Gly Ser Leu Gly Gly Pro Ala
1 5

<210> 257
<211> 9

<212> PRT
<213> Mus musculus

<400> 257
Ala Tyr Pro Gly Cys Asn Lys Arg Tyr
1 5

<210> 258
<211> 9
<212> PRT
<213> Mus musculus

<400> 258
Cys Met Thr Trp Asn Gln Met Asn Leu
1 5

<210> 259
<211> 9
<212> PRT
<213> Mus musculus

<400> 259
Cys Thr Gly Ser Gln Ala Leu Leu Leu
1 5

<210> 260
<211> 9
<212> PRT
<213> Mus musculus

<400> 260
Asp Gly Ala Pro Ser Tyr Gly His Thr
1 5

<210> 261
<211> 9
<212> PRT
<213> Mus musculus

<400> 261
Asp Leu Asn Ala Leu Leu Pro Ala Val
1 5

<210> 262
<211> 9
<212> PRT
<213> Mus musculus

<400> 262
Asp Pro Met Gly Gln Gln Gly Ser Leu
1 5

<210> 263
<211> 9
<212> PRT
<213> Mus musculus

<400> 263

Asp Ser Cys Thr Gly Ser Gln Ala Leu
1 5

<210> 264
<211> 9
<212> PRT
<213> Mus musculus

<400> 264

Asp Val Arg Asp Leu Asn Ala Leu Leu
1 5

<210> 265
<211> 9
<212> PRT
<213> Mus musculus

<400> 265

Glu Gln Cys Leu Ser Ala Phe Thr Leu
1 5

<210> 266
<211> 9
<212> PRT
<213> Mus musculus

<400> 266

Glu Ser Asp Asn His Thr Ala Pro Ile
1 5

<210> 267
<211> 9
<212> PRT
<213> Mus musculus

<400> 267

Phe Pro Asn Ala Pro Tyr Leu Pro Ser
1 5

<210> 268
<211> 9
<212> PRT
<213> Mus musculus

<400> 268

Gly Cys Asn Lys Arg Tyr Phe Lys Leu
1 5

<210> 269
<211> 9
<212> PRT
<213> Mus musculus

<400> 269

Gly Gln Ala Arg Met Phe Pro Asn Ala
1 5

<210> 270

<211> 9
<212> PRT
<213> Mus musculus

<400> 270
Gly Val Phe Arg Gly Ile Gln Asp Val
1 5

<210> 271
<211> 9
<212> PRT
<213> Mus musculus

<400> 271
Gly Tyr Glu Ser Asp Asn His Thr Ala
1 5

<210> 272
<211> 9
<212> PRT
<213> Mus musculus

<400> 272
His Ser Phe Lys His Glu Asp Pro Met
1 5

<210> 273
<211> 9
<212> PRT
<213> Mus musculus

<400> 273
His Thr His Gly Val Phe Arg Gly Ile
1 5

<210> 274
<211> 9
<212> PRT
<213> Mus musculus

<400> 274
Ile Leu Cys Gly Ala Gln Tyr Arg Ile
1 5

<210> 275
<211> 9
<212> PRT
<213> Mus musculus

<400> 275
Lys Phe Ala Arg Ser Asp Glu Leu Val
1 5

<210> 276
<211> 9
<212> PRT
<213> Mus musculus

<400> 276
Lys Arg Tyr Phe Lys Leu Ser His Leu
1 5

<210> 277
<211> 9
<212> PRT
<213> Mus musculus

<400> 277
Lys Thr Ser Glu Lys Pro Phe Ser Cys
1 5

<210> 278
<211> 9
<212> PRT
<213> Mus musculus

<400> 278
Leu Glu Cys Met Thr Trp Asn Gln Met
1 5

<210> 279
<211> 9
<212> PRT
<213> Mus musculus

<400> 279
Leu Gly Gly Gly Gly Gly Cys Gly Leu
1 5

<210> 280
<211> 9
<212> PRT
<213> Mus musculus

<400> 280
Leu Gln Met His Ser Arg Lys His Thr
1 5

<210> 281
<211> 9
<212> PRT
<213> Mus musculus

<400> 281
Met His Gln Arg Asn Met Thr Lys Leu
1 5

<210> 282
<211> 9
<212> PRT
<213> Mus musculus

<400> 282
Asn Ala Pro Tyr Leu Pro Ser Cys Leu
1 5

<210> 283
<211> 9
<212> PRT
<213> Mus musculus

<400> 283
Asn Leu Gly Ala Thr Leu Lys Gly Met
1 5

<210> 284
<211> 9
<212> PRT
<213> Mus musculus

<400> 284
Asn Leu Tyr Gln Met Thr Ser Gln Leu
1 5

<210> 285
<211> 9
<212> PRT
<213> Mus musculus

<400> 285
Asn Met Thr Lys Leu His Val Ala Leu
1 5

<210> 286
<211> 9
<212> PRT
<213> Mus musculus

<400> 286
Asn Gln Met Asn Leu Gly Ala Thr Leu
1 5

<210> 287
<211> 9
<212> PRT
<213> Mus musculus

<400> 287
Pro Gly Ala Ser Ala Tyr Gly Ser Leu
1 5

<210> 288
<211> 9
<212> PRT
<213> Mus musculus

<400> 288
Gln Ala Ser Ser Gly Gln Ala Arg Met
1 5

<210> 289
<211> 9
<212> PRT
<213> Mus musculus

<400> 289
Gln Met Thr Ser Gln Leu Glu Cys Met
1 5

<210> 290
<211> 9
<212> PRT
<213> Mus musculus

<400> 290
Gln Gln Tyr Ser Val Pro Pro Pro Val
1 5

<210> 291
<211> 9
<212> PRT
<213> Mus musculus

<400> 291
Gln Tyr Arg Ile His Thr His Gly Val
1 5

<210> 292
<211> 9
<212> PRT
<213> Mus musculus

<400> 292
Gln Tyr Ser Val Pro Pro Pro Val Tyr
1 5

<210> 293
<211> 9
<212> PRT
<213> Mus musculus

<400> 293
Arg Met Phe Pro Asn Ala Pro Tyr Leu
1 5

<210> 294
<211> 9
<212> PRT
<213> Mus musculus

<400> 294
Arg Thr Pro Tyr Ser Ser Asp Asn Leu
1 5

<210> 295
<211> 9
<212> PRT
<213> Mus musculus

<400> 295
Arg Val Ser Gly Val Ala Pro Thr Leu
1 5

<210> 296
<211> 9
<212> PRT
<213> Mus musculus

<400> 296
Ser Cys Leu Glu Ser Gln Pro Thr Ile
1 5

<210> 297
<211> 9
<212> PRT
<213> Mus musculus

<400> 297
Ser Cys Gln Lys Lys Phe Ala Arg Ser
1 5

<210> 298
<211> 9
<212> PRT
<213> Mus musculus

<400> 298
Ser Asp Val Arg Asp Leu Asn Ala Leu
1 5

<210> 299
<211> 9
<212> PRT
<213> Mus musculus

<400> 299
Ser Leu Gly Glu Gln Gln Tyr Ser Val
1 5

<210> 300
<211> 9
<212> PRT
<213> Mus musculus

<400> 300
Thr Cys Gln Arg Lys Phe Ser Arg Ser
1 5

<210> 301
<211> 9
<212> PRT
<213> Mus musculus

<400> 301
Thr Glu Gly Gln Ser Asn His Gly Ile
1 5

<210> 302
<211> 9
<212> PRT

<213> Mus musculus

<400> 302

Thr Leu His Phe Ser Gly Gln Phe Thr
1 5

<210> 303

<211> 9

<212> PRT

<213> Mus musculus

<400> 303

Thr Leu Val Arg Ser Ala Ser Glu Thr
1 5

<210> 304

<211> 9

<212> PRT

<213> Mus musculus

<400> 304

Val Leu Asp Phe Ala Pro Pro Gly Ala
1 5

<210> 305

<211> 9

<212> PRT

<213> Mus musculus

<400> 305

Trp Asn Gln Met Asn Leu Gly Ala Thr
1 5

<210> 306

<211> 9

<212> PRT

<213> Mus musculus

<400> 306

Tyr Phe Lys Leu Ser His Leu Gln Met
1 5

<210> 307

<211> 9

<212> PRT

<213> Mus musculus

<400> 307

Tyr Gln Met Thr Ser Gln Leu Glu Cys
1 5

<210> 308

<211> 9

<212> PRT

<213> Mus musculus

<400> 308

Tyr Ser Ser Asp Asn Leu Tyr Gln Met

```

1              5

<210> 309
<211> 6
<212> PRT
<213> Homo sapien

<400> 309
Gly Ala Ala Gln Trp Ala
1              5

<210> 310
<211> 12
<212> PRT
<213> Homo sapien

<400> 310
Ala Ser Ala Tyr Gly Ser Leu Gly Gly Pro Ala Pro
1              5              10

<210> 311
<211> 15
<212> PRT
<213> Homo sapien

<400> 311
Ala Phe Thr Val His Phe Ser Gly Gln Phe Thr Gly Thr Ala Gly
1              5              10              15

<210> 312
<211> 5
<212> PRT
<213> Homo sapien

<400> 312
His Ala Ala Gln Phe
1              5

<210> 313
<211> 32
<212> PRT
<213> Homo sapien

<400> 313
Cys His Thr Pro Thr Asp Ser Cys Thr Gly Ser Gln Ala Leu Leu Leu
1              5              10              15
Arg Thr Pro Tyr Ser Ser Asp Asn Leu Tyr Gln Met Thr Ser Gln Leu
                20              25              30

<210> 314
<211> 32
<212> PRT
<213> Homo sapien

<400> 314
Arg Ile His Thr His Gly Val Phe Arg Gly Ile Gln Asp Val Arg Arg
1              5              10              15
Val Pro Gly Val Ala Pro Thr Leu Val Arg Ser Ala Ser Glu Thr Ser

```

50

```

                20                25                30

<210> 315
<211> 4
<212> PRT
<213> Homo sapien

<400> 315
Arg Tyr Phe Lys
1

<210> 316
<211> 14
<212> PRT
<213> Homo sapien

<400> 316
Glu Arg Arg Phe Ser Arg Ser Asp Gln Leu Lys Arg His Gln
1          5          10

<210> 317
<211> 22
<212> PRT
<213> Homo sapien

<400> 317
Gln Arg Lys Phe Ser Arg Ser Asp His Leu Lys Thr His Thr Arg Thr
1          5          10          15
His Thr Gly Lys Thr Ser
20

<210> 318
<211> 21
<212> PRT
<213> Homo sapien

<400> 318
Cys Gln Lys Lys Phe Ala Arg Ser Asp Glu Leu Val Arg His His Asn
1          5          10          15
Met His Gln Arg Asn
20

<210> 319
<211> 449
<212> PRT
<213> Homo sapien

<400> 319
Met Gly Ser Asp Val Arg Asp Leu Asn Ala Leu Leu Pro Ala Val Pro
1          5          10          15
Ser Leu Gly Gly Gly Gly Gly Cys Ala Leu Pro Val Ser Gly Ala Ala
20          25          30
Gln Trp Ala Pro Val Leu Asp Phe Ala Pro Pro Gly Ala Ser Ala Tyr
35          40          45
Gly Ser Leu Gly Gly Pro Ala Pro Pro Pro Ala Pro Pro Pro Pro
50          55          60
Pro Pro Pro Pro His Ser Phe Ile Lys Gln Glu Pro Ser Trp Gly Gly
65          70          75          80

```


Ala	Glu	Pro	His	Glu	Glu	Gln	Cys	Leu	Ser	Ala	Phe	Thr	Val	His	Phe
				85					90					95	
Ser	Gly	Gln	Phe	Thr	Gly	Thr	Ala	Gly	Ala	Cys	Arg	Tyr	Gly	Pro	Phe
			100					105					110		
Gly	Pro	Pro	Pro	Pro	Ser	Gln	Ala	Ser	Ser	Gly	Gln	Ala	Arg	Met	Phe
			115				120					125			
Pro	Asn	Ala	Pro	Tyr	Leu	Pro	Ser	Cys	Leu	Glu	Ser	Gln	Pro	Ala	Ile
							135					140			
Arg	Asn	Gln	Gly	Tyr	Ser	Thr	Val	Thr	Phe	Asp	Gly	Thr	Pro	Ser	Tyr
145					150					155					160
Gly	His	Thr	Pro	Ser	His	His	Ala	Ala	Gln	Phe	Pro	Asn	His	Ser	Phe
				165						170					175
Lys	His	Glu	Asp	Pro	Met	Gly	Gln	Gln	Gly	Ser	Leu	Gly	Glu	Gln	Gln
			180					185					190		
Tyr	Ser	Val	Pro	Pro	Pro	Val	Tyr	Gly	Cys	His	Thr	Pro	Thr	Asp	Ser
			195				200					205			
Cys	Thr	Gly	Ser	Gln	Ala	Leu	Leu	Leu	Arg	Thr	Pro	Tyr	Ser	Ser	Asp
						215					220				
Asn	Leu	Tyr	Gln	Met	Thr	Ser	Gln	Leu	Glu	Cys	Met	Thr	Trp	Asn	Gln
225					230					235					240
Met	Asn	Leu	Gly	Ala	Thr	Leu	Lys	Gly	Val	Ala	Ala	Gly	Ser	Ser	Ser
				245					250					255	
Ser	Val	Lys	Trp	Thr	Glu	Gly	Gln	Ser	Asn	His	Ser	Thr	Gly	Tyr	Glu
			260					265					270		
Ser	Asp	Asn	His	Thr	Thr	Pro	Ile	Leu	Cys	Gly	Ala	Gln	Tyr	Arg	Ile
			275				280					285			
His	Thr	His	Gly	Val	Phe	Arg	Gly	Ile	Gln	Asp	Val	Arg	Arg	Val	Pro
			290			295				300					
Gly	Val	Ala	Pro	Thr	Leu	Val	Arg	Ser	Ala	Ser	Glu	Thr	Ser	Glu	Lys
305					310					315				320	
Arg	Pro	Phe	Met	Cys	Ala	Tyr	Pro	Gly	Cys	Asn	Lys	Arg	Tyr	Phe	Lys
				325					330					335	
Leu	Ser	His	Leu	Gln	Met	His	Ser	Arg	Lys	His	Thr	Gly	Glu	Lys	Pro
			340					345					350		
Tyr	Gln	Cys	Asp	Phe	Lys	Asp	Cys	Glu	Arg	Arg	Phe	Ser	Arg	Ser	Asp
			355				360					365			
Gln	Leu	Lys	Arg	His	Gln	Arg	Arg	His	Thr	Gly	Val	Lys	Pro	Phe	Gln
						375					380				
Cys	Lys	Thr	Cys	Gln	Arg	Lys	Phe	Ser	Arg	Ser	Asp	His	Leu	Lys	Thr
385					390					395				400	
His	Thr	Arg	Thr	His	Thr	Gly	Lys	Thr	Ser	Glu	Lys	Pro	Phe	Ser	Cys
				405					410					415	
Arg	Trp	Pro	Ser	Cys	Gln	Lys	Lys	Phe	Ala	Arg	Ser	Asp	Glu	Leu	Val
			420					425					430		
Arg	His	His	Asn	Met	His	Gln	Arg	Asn	Met	Thr	Lys				

```
<210> 320
<211> 449
<212> PRT
<213> Mus musculus
```

<400> 320
Met Gly Ser Asp Val Arg Asp Leu Asn Ala Leu Leu Pro Ala Val Ser
1 5 10 15
Ser Leu Gly Gly Gly Gly Gly Cys Gly Leu Pro Val Ser Gly Ala Ala

			20					25					30		
Gln	Trp	Ala	Pro	Val	Leu	Asp	Phe	Ala	Pro	Pro	Gly	Ala	Ser	Ala	Tyr
		35					40					45			
Gly	Ser	Leu	Gly	Gly	Pro	Ala	Pro	Pro	Pro	Ala	Pro	Pro	Pro	Pro	Pro
	50					55					60				
Pro	Pro	Pro	Pro	His	Ser	Phe	Ile	Lys	Gln	Glu	Pro	Ser	Trp	Gly	Gly
65					70					75					80
Ala	Glu	Pro	His	Glu	Gln	Cys	Leu	Ser	Ser	Ala	Phe	Thr	Leu	His	Phe
				85				90						95	
Ser	Gly	Gln	Phe	Thr	Gly	Thr	Ala	Gly	Ala	Cys	Arg	Tyr	Gly	Pro	Phe
			100					105					110		
Gly	Pro	Pro	Pro	Pro	Ser	Gln	Ala	Ser	Ser	Gly	Gln	Ala	Arg	Met	Phe
		115					120					125			
Pro	Asn	Ala	Pro	Tyr	Leu	Pro	Ser	Cys	Leu	Glu	Ser	Gln	Pro	Thr	Ile
	130					135						140			
Arg	Asn	Gln	Gly	Tyr	Ser	Thr	Val	Thr	Phe	Asp	Gly	Ala	Pro	Ser	Tyr
145					150					155					160
Gly	His	Thr	Pro	Ser	His	His	Ala	Ala	Gln	Phe	Pro	Asn	His	Ser	Phe
				165						170				175	
Lys	His	Glu	Asp	Pro	Met	Gly	Gln	Gln	Gly	Ser	Leu	Gly	Glu	Gln	Gln
			180					185					190		
Tyr	Ser	Val	Pro	Pro	Pro	Val	Tyr	Gly	Cys	His	Thr	Pro	Thr	Asp	Ser
		195					200					205			
Cys	Thr	Gly	Ser	Gln	Ala	Leu	Leu	Leu	Arg	Thr	Pro	Tyr	Ser	Ser	Asp
	210					215					220				
Asn	Leu	Tyr	Gln	Met	Thr	Ser	Gln	Leu	Glu	Cys	Met	Thr	Trp	Asn	Gln
225					230					235					240
Met	Asn	Leu	Gly	Ala	Thr	Leu	Lys	Gly	Met	Ala	Ala	Gly	Ser	Ser	Ser
				245					250					255	
Ser	Val	Lys	Trp	Thr	Glu	Gly	Gln	Ser	Asn	His	Gly	Ile	Gly	Tyr	Glu
			260					265					270		
Ser	Asp	Asn	His	Thr	Ala	Pro	Ile	Leu	Cys	Gly	Ala	Gln	Tyr	Arg	Ile
		275					280					285			
His	Thr	His	Gly	Val	Phe	Arg	Gly	Ile	Gln	Asp	Val	Arg	Arg	Val	Ser
	290					295					300				
Gly	Val	Ala	Pro	Thr	Leu	Val	Arg	Ser	Ala	Ser	Glu	Thr	Ser	Glu	Lys
305					310					315					320
Arg	Pro	Phe	Met	Cys	Ala	Tyr	Pro	Gly	Cys	Asn	Lys	Arg	Tyr	Phe	Lys
				325					330					335	
Leu	Ser	His	Leu	Gln	Met	His	Ser	Arg	Lys	His	Thr	Gly	Glu	Lys	Pro
			340					345					350		
Tyr	Gln	Cys	Asp	Phe	Lys	Asp	Cys	Glu	Arg	Arg	Phe	Ser	Arg	Ser	Asp
		355					360					365			
Gln	Leu	Lys	Arg	His	Gln	Arg	Arg	His	Thr	Gly	Val	Lys	Pro	Phe	Gln
	370					375					380				

$$\begin{aligned} \langle 210 \rangle & 321 \\ \langle 211 \rangle & 9 \end{aligned}$$

<212> PRT
<213> Homo sapien and Mus musculus

<400> 321
Pro Ser Gln Ala Ser Ser Gly Gln Ala
1 5

<210> 322
<211> 9
<212> PRT
<213> Homo sapien and Mus musculus

<400> 322
Ser Ser Gly Gln Ala Arg Met Phe Pro
1 5

<210> 323
<211> 9
<212> PRT
<213> Homo sapien and Mus musculus

<400> 323
Gln Ala Arg Met Phe Pro Asn Ala Pro
1 5

<210> 324
<211> 9
<212> PRT
<213> Homo sapien and Mus musculus

<400> 324
Met Phe Pro Asn Ala Pro Tyr Leu Pro
1 5

<210> 325
<211> 9
<212> PRT
<213> Homo sapien and Mus musculus

<400> 325
Pro Asn Ala Pro Tyr Leu Pro Ser Cys
1 5

<210> 326
<211> 9
<212> PRT
<213> Homo sapien and Mus musculus

<400> 326
Ala Pro Tyr Leu Pro Ser Cys Leu Glu
1 5

<210> 327
<211> 261
<212> PRT
<213> Homo sapiens

<400> 327

```

Met Trp Val Pro Val Val Phe Leu Thr Leu Ser Val Thr Trp Ile Gly
                    5                      10                      15

Ala Ala Pro Leu Ile Leu Ser Arg Ile Val Gly Gly Trp Glu Cys Glu
                20                      25                      30

Lys His Ser Gln Pro Trp Gln Val Leu Val Ala Ser Arg Gly Arg Ala
                35                      40                      45

Val Cys Gly Gly Val Leu Val His Pro Gln Trp Val Leu Thr Ala Ala
                50                      55                      60

His Cys Ile Arg Asn Lys Ser Val Ile Leu Leu Gly Arg His Ser Leu
                65                      70                      75

Phe His Pro Glu Asp Thr Gly Gln Val Phe Gln Val Ser His Ser Phe
                85                      90                      95

Pro His Pro Leu Tyr Asp Met Ser Leu Leu Lys Asn Arg Phe Leu Arg
                100                     105                     110

Pro Gly Asp Asp Ser Ser His Asp Leu Met Leu Leu Arg Leu Ser Glu
                115                     120                     125

Pro Ala Glu Leu Thr Asp Ala Val Lys Val Met Asp Leu Pro Thr Gln
                130                     135                     140

Glu Pro Ala Leu Gly Thr Thr Cys Tyr Ala Ser Gly Trp Gly Ser Ile
                145                     150                     155                     160

Glu Pro Glu Glu Phe Leu Thr Pro Lys Lys Leu Gln Cys Val Asp Leu
                165                     170                     175

His Val Ile Ser Asn Asp Val Cys Ala Gln Val His Pro Gln Lys Val
                180                     185                     190

Thr Lys Phe Met Leu Cys Ala Gly Arg Trp Thr Gly Gly Lys Ser Thr
                195                     200                     205

Cys Ser Gly Asp Ser Gly Gly Pro Leu Val Cys Asn Gly Val Leu Gln
                210                     215                     220

Gly Ile Thr Ser Trp Gly Ser Glu Pro Cys Ala Leu Pro Glu Arg Pro
                225                     230                     235                     240

Ser Leu Tyr Thr Lys Val Val His Tyr Arg Lys Trp Ile Lys Asp Thr
                245                     250                     255

Ile Val Ala Asn Pro
                260

```

<210> 328

<211> 386

<212> PRT

<213> Homo sapiens

<400> 328

```

Met Arg Ala Ala Pro Leu Leu Leu Ala Arg Ala Ala Ser Leu Ser Leu
           5              10              15

Gly Phe Leu Phe Leu Leu Phe Phe Trp Leu Asp Arg Ser Val Leu Ala
           20              25              30

Lys Glu Leu Lys Phe Val Thr Leu Val Phe Arg His Gly Asp Arg Ser
           35              40              45

Pro Ile Asp Thr Phe Pro Thr Asp Pro Ile Lys Glu Ser Ser Trp Pro
           50              55              60

Gln Gly Phe Gly Gln Leu Thr Gln Leu Gly Met Glu Gln His Tyr Glu
           65              70              75              80

Leu Gly Glu Tyr Ile Arg Lys Arg Tyr Arg Lys Phe Leu Asn Glu Ser
           85              90              95

Tyr Lys His Glu Gln Val Tyr Ile Arg Ser Thr Asp Val Asp Arg Thr
           100             105             110

Leu Met Ser Ala Met Thr Asn Leu Ala Ala Leu Phe Pro Pro Glu Gly
           115             120             125

Val Ser Ile Trp Asn Pro Ile Leu Leu Trp Gln Pro Ile Pro Val His
           130             135             140

Thr Val Pro Leu Ser Glu Asp Gln Leu Leu Tyr Leu Pro Phe Arg Asn
           145             150             155             160

Cys Pro Arg Phe Gln Glu Leu Glu Ser Glu Thr Leu Lys Ser Glu Glu
           165             170             175

Phe Gln Lys Arg Leu His Pro Tyr Lys Asp Phe Ile Ala Thr Leu Gly
           180             185             190

Lys Leu Ser Gly Leu His Gly Gln Asp Leu Phe Gly Ile Trp Ser Lys
           195             200             205

Val Tyr Asp Pro Leu Tyr Cys Glu Ser Val His Asn Phe Thr Leu Pro
           210             215             220

Ser Trp Ala Thr Glu Asp Thr Met Thr Lys Leu Arg Glu Leu Ser Glu
           225             230             235             240

Leu Ser Leu Leu Ser Leu Tyr Gly Ile His Lys Gln Lys Glu Lys Ser
           245             250             255

Arg Leu Gln Gly Gly Val Leu Val Asn Glu Ile Leu Asn His Met Lys
           260             265             270

Arg Ala Thr Gln Ile Pro Ser Tyr Lys Lys Leu Ile Met Tyr Ser Ala
           275             280             285

His Asp Thr Thr Val Ser Gly Leu Gln Met Ala Leu Asp Val Tyr Asn

```

290 295 300
 Gly Leu Leu Pro Pro Tyr Ala Ser Cys His Leu Thr Glu Leu Tyr Phe
 305 310 315 320
 Glu Lys Gly Glu Tyr Phe Val Glu Met Tyr Tyr Arg Asn Glu Thr Gln
 325 330 335
 His Glu Pro Tyr Pro Leu Met Leu Pro Gly Cys Ser Pro Ser Cys Pro
 340 345 350
 Leu Glu Arg Phe Ala Glu Leu Val Gly Pro Val Ile Pro Gln Asp Trp
 355 360 365
 Ser Thr Glu Cys Met Thr Thr Asn Ser His Gln Gly Thr Glu Asp Ser
 370 375 380
 Thr Asp
 385

 <210> 329
 <211> 261
 <212> PRT
 <213> Homo sapiens

 <400> 329
 Met Trp Val Pro Val Val Phe Leu Thr Leu Ser Val Thr Trp Ile Gly
 5 10 15
 Ala Ala Pro Leu Ile Leu Ser Arg Ile Val Gly Gly Trp Glu Cys Glu
 20 25 30
 Lys His Ser Gln Pro Trp Gln Val Leu Val Ala Ser Arg Gly Arg Ala
 35 40 45
 Val Cys Gly Gly Val Leu Val His Pro Gln Trp Val Leu Thr Ala Ala
 50 55 60
 His Cys Ile Arg Asn Lys Ser Val Ile Leu Leu Gly Arg His Ser Leu
 65 70 75 80
 Phe His Pro Glu Asp Thr Gly Gln Val Phe Gln Val Ser His Ser Phe
 85 90 95
 Pro His Pro Leu Tyr Asp Met Ser Leu Leu Lys Asn Arg Phe Leu Arg
 100 105 110
 Pro Gly Asp Asp Ser Ser His Asp Leu Met Leu Leu Arg Leu Ser Glu
 115 120 125
 Pro Ala Glu Leu Thr Asp Ala Val Lys Val Met Asp Leu Pro Thr Gln
 130 135 140
 Glu Pro Ala Leu Gly Thr Thr Cys Tyr Ala Ser Gly Trp Gly Ser Ile
 145 150 155 160
 Glu Pro Glu Glu Phe Leu Thr Pro Lys Lys Leu Gln Cys Val Asp Leu

57

	165		170		175
His Val Ile Ser Asn Asp Val Cys Ala Gln Val His Pro Gln Lys Val	180	185	190		
Thr Lys Phe Met Leu Cys Ala Gly Arg Trp Thr Gly Gly Lys Ser Thr	195	200	205		
Cys Ser Gly Asp Ser Gly Gly Pro Leu Val Cys Asn Gly Val Leu Gln	210	215	220		
Gly Ile Thr Ser Trp Gly Ser Glu Pro Cys Ala Leu Pro Glu Arg Pro	225	230	235	240	
Ser Leu Tyr Thr Lys Val Val His Tyr Arg Lys Trp Ile Lys Asp Thr	245	250	255		
Ile Val Ala Asn Pro	260				

<210> 330
 <211> 220
 <212> PRT
 <213> Homo sapien

	<400> 330
Glu Asp Cys Ser Pro His Ser Gln Pro Trp Gln Ala Ala Leu Val Met	1 5 10 15
Glu Asn Glu Leu Phe Cys Ser Gly Val Leu Val His Pro Gln Trp Val	20 25 30
Leu Ser Ala Ala His Cys Phe Gln Asn Ser Tyr Thr Ile Gly Leu Gly	35 40 45
Leu His Ser Leu Glu Ala Asp Gln Glu Pro Gly Ser Gln Met Val Glu	50 55 60
Ala Ser Leu Ser Val Arg His Pro Glu Tyr Asn Arg Pro Leu Leu Ala	65 70 75 80
Asn Asp Leu Met Leu Ile Lys Leu Asp Glu Ser Val Ser Glu Ser Asp	85 90 95
Thr Ile Arg Ser Ile Ser Ile Ala Ser Gln Cys Pro Thr Ala Gly Asn	100 105 110
Ser Cys Leu Val Ser Gly Trp Gly Leu Leu Ala Asn Gly Arg Met Pro	115 120 125
Thr Val Leu Gln Cys Val Asn Val Ser Val Val Ser Glu Glu Val Cys	130 135 140
Ser Lys Leu Tyr Asp Pro Leu Tyr His Pro Ser Met Phe Cys Ala Gly	145 150 155 160
Gly Gly Gln Asp Gln Lys Asp Ser Cys Asn Gly Asp Ser Gly Gly Pro	165 170 175
Leu Ile Cys Asn Gly Tyr Leu Gln Gly Leu Val Ser Phe Gly Lys Ala	180 185 190
Pro Cys Gly Gln Val Gly Val Pro Gly Val Tyr Thr Asn Leu Cys Lys	195 200 205
Phe Thr Glu Trp Ile Glu Lys Thr Val Gln Ala Ser	210 215 220

<210> 331
 <211> 77

<212> PRT

<213> Homo sapien

<400> 331

Leu	Val	Ser	Gly	Ser	Cys	Ser	Gln	Ile	Ile	Asn	Gly	Glu	Asp	Cys	Ser
1				5					10					15	
Pro	His	Ser	Gln	Pro	Trp	Gln	Ala	Ala	Leu	Val	Met	Glu	Asn	Glu	Leu
			20				25						30		
Phe	Cys	Ser	Gly	Val	Leu	Val	His	Pro	Gln	Trp	Val	Leu	Ser	Ala	Thr
		35					40					45			
His	Cys	Phe	Gln	Asn	Ser	Tyr	Thr	Ile	Gly	Leu	Gly	Leu	His	Ser	Leu
	50					55					60				
Glu	Ala	Asp	Gln	Glu	Pro	Gly	Ser	Gln	Met	Val	Glu	Ala			
65					70					75					

<210> 332

<211> 22

<212> PRT

<213> Homo sapien

<400> 332

Gln	His	Asn	Gly	Pro	Ile	Pro	Ser	Leu	Thr	Pro	Pro	Ser	Gly	Ser	Leu
1				5					10					15	
Val	Ser	Gly	Ser	Cys	Ser										
			20												

<210> 333

<211> 553

<212> PRT

<213> Homo sapien

<400> 333

Met	Val	Gln	Arg	Leu	Trp	Val	Ser	Arg	Leu	Leu	Arg	His	Arg	Lys	Ala
1				5					10					15	
Gln	Leu	Leu	Leu	Val	Asn	Leu	Leu	Thr	Phe	Gly	Leu	Glu	Val	Cys	Leu
			20					25					30		
Ala	Ala	Gly	Ile	Thr	Tyr	Val	Pro	Pro	Leu	Leu	Leu	Glu	Val	Gly	Val
		35					40					45			
Glu	Glu	Lys	Phe	Met	Thr	Met	Val	Leu	Gly	Ile	Gly	Pro	Val	Leu	Gly
	50					55					60				
Leu	Val	Cys	Val	Pro	Leu	Leu	Gly	Ser	Ala	Ser	Asp	His	Trp	Arg	Gly
65					70				75					80	
Arg	Tyr	Gly	Arg	Arg	Arg	Pro	Phe	Ile	Trp	Ala	Leu	Ser	Leu	Gly	Ile
			85					90						95	
Leu	Leu	Ser	Leu	Phe	Leu	Ile	Pro	Arg	Ala	Gly	Trp	Leu	Ala	Gly	Leu
		100						105					110		
Leu	Cys	Pro	Asp	Pro	Arg	Pro	Leu	Glu	Leu	Ala	Leu	Leu	Ile	Leu	Gly
		115					120					125			
Val	Gly	Leu	Leu	Asp	Phe	Cys	Gly	Gln	Val	Cys	Phe	Thr	Pro	Leu	Glu
	130					135					140				
Ala	Leu	Leu	Ser	Asp	Leu	Phe	Arg	Asp	Pro	Asp	His	Cys	Arg	Gln	Ala
145					150					155				160	
Tyr	Ser	Val	Tyr	Ala	Phe	Met	Ile	Ser	Leu	Gly	Gly	Cys	Leu	Gly	Tyr
			165					170						175	
Leu	Leu	Pro	Ala	Ile	Asp	Trp	Asp	Thr	Ser	Ala	Leu	Ala	Pro	Tyr	Leu
		180						185					190		
Gly	Thr	Gln	Glu	Glu	Cys	Leu	Phe	Gly	Leu	Leu	Thr	Leu	Ile	Phe	Leu

		195					200					205						
Thr	Cys	Val	Ala	Ala	Thr	Leu	Leu	Val	Ala	Glu	Glu	Ala	Ala	Leu	Gly			
	210					215					220							
Pro	Thr	Glu	Pro	Ala	Glu	Gly	Leu	Ser	Ala	Pro	Ser	Leu	Ser	Pro	His			
225					230					235					240			
Cys	Cys	Pro	Cys	Arg	Ala	Arg	Leu	Ala	Phe	Arg	Asn	Leu	Gly	Ala	Leu			
				245					250					255				
Leu	Pro	Arg	Leu	His	Gln	Leu	Cys	Cys	Arg	Met	Pro	Arg	Thr	Leu	Arg			
			260					265					270					
Arg	Leu	Phe	Val	Ala	Glu	Leu	Cys	Ser	Trp	Met	Ala	Leu	Met	Thr	Phe			
		275					280					285						
Thr	Leu	Phe	Tyr	Thr	Asp	Phe	Val	Gly	Glu	Gly	Leu	Tyr	Gln	Gly	Val			
	290				295						300							
Pro	Arg	Ala	Glu	Pro	Gly	Thr	Glu	Ala	Arg	Arg	His	Tyr	Asp	Glu	Gly			
305					310					315					320			
Val	Arg	Met	Gly	Ser	Leu	Gly	Leu	Phe	Leu	Gln	Cys	Ala	Ile	Ser	Leu			
				325					330					335				
Val	Phe	Ser	Leu	Val	Met	Asp	Arg	Leu	Val	Gln	Arg	Phe	Gly	Thr	Arg			
			340					345					350					
Ala	Val	Tyr	Leu	Ala	Ser	Val	Ala	Ala	Phe	Pro	Val	Ala	Ala	Gly	Ala			
		355				360						365						
Thr	Cys	Leu	Ser	His	Ser	Val	Ala	Val	Val	Thr	Ala	Ser	Ala	Ala	Leu			
	370				375						380							
Thr	Gly	Phe	Thr	Phe	Ser	Ala	Leu	Gln	Ile	Leu	Pro	Tyr	Thr	Leu	Ala			
385					390					395					400			
Ser	Leu	Tyr	His	Arg	Glu	Lys	Gln	Val	Phe	Leu	Pro	Lys	Tyr	Arg	Gly			
			405						410					415				
Asp	Thr	Gly	Gly	Ala	Ser	Ser	Glu	Asp	Ser	Leu	Met	Thr	Ser	Phe	Leu			
			420					425				430						
Pro	Gly	Pro	Lys	Pro	Gly	Ala	Pro	Phe	Pro	Asn	Gly	His	Val	Gly	Ala			
		435				440						445						
Gly	Gly	Ser	Gly	Leu	Leu	Pro	Pro	Pro	Pro	Ala	Leu	Cys	Gly	Ala	Ser			
	450				455					460								
Ala	Cys	Asp	Val	Ser	Val	Arg	Val	Val	Val	Gly	Glu	Pro	Thr	Glu	Ala			
465					470					475					480			
Arg	Val	Val	Pro	Gly	Arg	Gly	Ile	Cys	Leu	Asp	Leu	Ala	Ile	Leu	Asp			
			485						490					495				
Ser	Ala	Phe	Leu	Leu	Ser	Gln	Val	Ala	Pro	Ser	Leu	Phe	Met	Gly	Ser			
			500					505					510					
Ile	Val	Gln	Leu	Ser	Gln	Ser	Val											

```
<210> 334
<211> 385
<212> DNA
<213> Homo sapien
```

<400> 334						
actacacaca	ctccacttgc	ccttgtgaga	cactttgtcc	cagcacttta	ggaatgctga	60
ggtcggacca	gccacatctc	atgtgcaaga	ttgccagca	gacatcaggt	ctgagagttc	120
cccttttaaa	aaaggggact	tgtctaaaaa	agaagtctag	ccacgattgt	gtagagcagc	180
tgtgtgtgc	tggagattca	cttttgagag	agttctctc	tgagacctga	tccttagagg	240
ctgggcagtc	ttgcacatga	gatggggctg	gtctgatctc	agcactcctt	agtctgcttg	300

cctctcccag ggccccagcc tggccacacc tgcttacagg gcactctcag atgcccatatc 360
 catagtttct gtgctagtgg accgt 385

<210> 335
 <211> 647
 <212> DNA
 <213> Homo sapien
 <220>
 <221> misc_feature
 <222> (1)...(647)
 <223> n = A,T,C or G

<400> 335
 acgattttca ttatcatgta aatcgggtca ctcaaggggc caaccacagc tgggagccac 60
 tgctcagggg aaggttcata tgggactttc tactgcccac gggtctatac aggatataaa 120
 ggngcctcac agtatagatc tggtagcaaa gaagaagaaa caaacactga tctctttctg 180
 ccacccctct gacccttttg aactcctctg acccttttaga acaagcctac ctaatatctg 240
 cttagagaaa gaccaacaac ggccctcaaag gatctcttac catgaaggtc tcagctaatt 300
 cttggctaag atgtgggttc cacattaggt tctgaatatg gggggaaggg tcaatttgct 360
 cattttgtgt gtggataaag tcaggatgcc caggggccag agcagggggc tgcttgcttt 420
 gggaacaatg gctgagcata taaccatagg ttatggggaa caaaacaaca tcaaagtcac 480
 tgtatcaatt gccatgaaga cttgagggac ctgaatctac cgattcatct taaggcagca 540
 ggaccagttt gagtggcaac aatgcagcag cagaatcaat ggaaacaaca gaatgattgc 600
 aatgtccttt tttttctcct gcttctgact tgataaaagg ggaccgt 647

<210> 336
 <211> 526
 <212> DNA
 <213> Homo sapien
 <220>
 <221> misc_feature
 <222> (1)...(526)
 <223> n = A,T,C or G

<400> 336
 caaatttgag ccaatgacat agaattttac aaatcaagaa gcttattctg gggccatttc 60
 ttttgacgtt ttctctaaac tactaaagag gcattaatga tccataaatt atattatcta 120
 ctttacagc attttaaagt tgttcagcat gaaatattag ctacagggga agctaaataa 180
 attaaacatg gaataaagat ttgtccttaa atataatcta caagaagact ttgatatttg 240
 tttttcacia gtgaagcatt cttataaagt gtcataacct ttttggggaa actatgggaa 300
 aaaatgggga aactctgaag ggtttttaagt atcttacctg aagctacaga ctccataacc 360
 tctctttaca gggagctcct gcagccccta cagaaatgag tggctgagat tcttgattgc 420
 acagcaagag cttctcatct aaaccctttc ctttttttagt atctgtgtat caagtataaa 480
 agttctataa actgtagtnt acttatttta atccccaaag cacagt 526

<210> 337
 <211> 854
 <212> DNA
 <213> Homo sapien

<400> 337
 ccttttctag ttcaccagtt ttctgcaagg atgctgggta gggagtgtct gcaggaggag 60
 caagtctgaa accaaatcta ggaaacatag gaaacgagcc aggcacaggg ctgggtgggccc 120
 atcaggggacc accctttggg ttgatatttt gcttaatctg catcttttga gtaagatcat 180

ctggcagtag	aagctgttct	ccagggtacat	ttctctagct	catgtacaaa	aacatcctga	240
aggactttgt	caggtgcctt	gctaaaagcc	agatgcgttc	ggcacttcct	tggctctgagg	300
ttaattgcac	acctacaggc	actgggctca	tgctttcaag	tattttgtcc	tcacttttagg	360
gtgagtga	gatccccatt	ataggagcac	ttgggagaga	tcatataaaa	gctgactctt	420
gagtacatgc	agtaatgggg	tagatgtgtg	tgggtgtgtc	tcattcctgc	aaggggtgctt	480
gttagggagt	gtttccagga	ggaacaagtc	tgaaaaccaat	catgaaataa	atggtaggtg	540
tgaactggaa	aactaattca	aaagagagat	cgtgatatca	gtgtggttga	tacaccttgg	600
caatatggaa	ggctctaatt	tgcccatatt	tgaataata	attcagcttt	ttgtaataca	660
aaataacaaa	ggattgagaa	tcattggtgtc	taatgtataa	aagaccagc	aaacataaat	720
atatcaactg	cataaatgta	aaatgcatgt	gacccaagaa	ggccccaag	tggcagacaa	780
cattgtaccc	attttccctt	ccaaaatgtg	agcggcgggc	ctgctgcttt	caaggctgtc	840
acacgggatg	tcag					854

<210> 338

<211> 251

<212> DNA

<213> Homo sapien

<400> 338

ctcaaagcta	atctctcggg	aatcaaacca	gaaaagggca	aggatcttag	gcatgggtga	60
tgtggataag	gccagggtcaa	tggctgcaag	catgcagaga	aagagggtaca	tcggagcgtg	120
caggctgcgt	tccgtcctta	cgatgaagac	cacgatgcag	tttccaaaca	ttgccactac	180
atacatggaa	aggaggggga	agccaacca	gaaatgggct	ttctctaatac	ctgggatacc	240
aataagcaca	a					251

<210> 339

<211> 3279

<212> DNA

<213> Homo sapiens

<400> 339

cttctcgcag	cccccatgct	ggtgaggggc	acgggcagga	acagtggacc	caacatggaa	60
atgctggagg	gtgtcaggaa	gtgatcgggc	tctggggcag	ggaggagggg	tggggagtgt	120
cactgggagg	ggacatcctg	cagaaggtag	gagtgcagaa	acacccgctg	caggggaggg	180
gagagccctg	cggcacctgg	gggagcagag	ggagcagcac	ctgcccaggc	ctgggaggag	240
gggcctggag	ggcgtgagga	ggagcgaggg	ggctgcatgg	ctggagttag	ggatcagggg	300
cagggcgcgga	gatggcctca	cacagggaag	agagggcccc	tctcgcaggg	cctcacctgg	360
gccacaggag	gacactgctt	ttcctctgag	gagtcaggag	ctgtggatgg	tgtcggacag	420
aagaaggaca	gggcctggct	caggtgtcca	gaggctgtcg	ctggcttccc	tttgggatca	480
gactgcaggg	agggagggcg	gcagggttgt	ggggggagtg	acgatgagga	tgacctgggg	540
gtggctccag	gccttgcccc	tgcctggggc	ctcacccagc	ctccctcaca	gtctcctggc	600
cctcagtctc	tccccctcac	tccatcctcc	atctggcctc	agtgggtcat	tctgatcact	660
gaactgacca	taccagccc	tggccacggc	cctccatggc	tccccaatgc	cctggagagg	720
ggacatctag	tcagagagta	gtcctgaaga	ggtggcctct	gcgatgtgcc	tgtgggggca	780
gcacctcgca	gatgggtccc	gccctcatcc	tgtcgcacctg	tctgcaggga	ctgtcctcct	840
ggaccttgcc	ccttgtgcag	gagctggacc	ctgaagtccc	ctcccatag	gccaaactg	900
gagccttggt	ccctctgttg	gactccctgc	ccatattctt	gtgggagtgg	gttctggaga	960
catttctgtc	gtttcttgag	agctgggaat	tgcctcagtc	catctgcctg	cgcggttctg	1020
agagatggag	ttgcctaggc	agttattggg	gccaatcttt	ctcactgtgt	ctctcctcct	1080
ttacccttag	ggtgattctg	ggggctccact	tgtctgtaat	ggtgtgcttc	aaggtatcac	1140
atcatggggc	cctgagccat	gtgccctgcc	tgaaaagcct	gctgtgtaca	ccaaggtggg	1200
gcattaccgg	aagtggatca	aggacaccat	cgcagccaac	ccctgagtgc	ccctgtccca	1260
cccctacctc	tagtaaattt	aagtccacct	cacgttcttg	catcacttgg	cctttctgga	1320
tgtcggacac	ctgaagcttg	gaactcacct	ggccgaagct	cgagcctcct	gagtcctact	1380
gacctgtgct	ttctggtgtg	gagtcagggg	ctgctaggaa	aagggaatggg	cagacacagg	1440
tgtatgcaa	tgtttctgaa	atgggtataa	tttctgcttc	tccttcggaa	cactggctgt	1500
ctctgaagac	ttctcgctca	gtttcagtga	ggacacacac	aaagacgtgg	gtgaccatgt	1560

```

tgtttgtggg gtgcagagat gggaggggtg gggcccaccc tggaagagtg gacagtgaca 1620
caaggtggac actctctaca gatcactgag gataagctgg agccacaatg catgaggcac 1680
acacacagca aggttgacgc tgtaaacata gccacgctg tcctgggggc actgggaagc 1740
ctagataagg cagttagcag aaagaagggg aggatcctcc tatgttgttg aaggagggac 1800
tagggggaga aactgaaagc tgattaatta caggaggttt gttcaggtcc cccaaaccac 1860
cgtcagattt gatgatttcc tagcaggact tacagaaata aagagctatc atgctgtggt 1920
ttattatggg ttgttacatt gataggatac atactgaaat cagcaaacaa aacagatgta 1980
tagattagag tgtggagaaa acagaggaaa acttgacgtt acgaagactg gcaacttggc 2040
tttactaagt tttcagactg gcaggaagtc aaacctatta ggctgaggac cttgtggagt 2100
gtagctgac cagctgatag aggaactagc cagggtgggg cctttccctt tggatggggg 2160
gcatatccga cagttattct ctccaagtgg agacttacgg acagcatata attctccctg 2220
caaggatgta tgataatatg taaaaagtaa ttccaactga ggaagctcac ctgatcctta 2280
gtgtccaggg tttttactgg ggggtctgtg gacgagtatg gactacttga ataattgacc 2340
tgaagtcttc agacctgagg ttccctagag ttcaaacaga tacagcatgg tccagagtcc 2400
cagatgtaca aaaacaggga ttcatcacia atcccatctt tagcatgaag ggtctggcat 2460
ggcccaaggc cccaagtata tcaaggcact tgggcagaac atgccaagga atcaaagtgc 2520
atctcccagg agttattcaa gggtagagccc tttacttggg atgtacaggc tttgagcagt 2580
gcagggtctg tgagtcaacc ttttattgta caggggatga gggaaagga gaggatgagg 2640
aagccccctt ggggatttgg tttggtcttg tgatcaggtg gtctatgggg ctatccctac 2700
aaagaagaat ccagaaatag gggcacattg aggaatgata ctgagcccaa agagcattca 2760
atcattgttt tatttgcctt cttttcacac cattggtgag ggagggatta ccaccctggg 2820
gttatgaaga tggttgaaca cccacacat agcaccggag atatgagatc aacagtttct 2880
tagccataga gattcacagc ccagagcagg aggacgctgc acaccatgca ggatgacatg 2940
ggggatgcgc tcgggattgg tgtgaagaag caaggactgt tagaggcagg ctttatagta 3000
acaagacggt ggggcaaact ctgatttccg tgggggaatg tcatggtctt gctttactaa 3060
gttttgagac tggcaggtag tgaaactcat taggctgaga accttgtgga atgcagctga 3120
cccagctgat agaggaagta gccaggtggg agcctttccc agtgggtgtg ggacatatct 3180
ggcaagattt tgtggcactc ctggttacag atactggggc agcaaataaa actgaatctt 3240
gttttcagac cttaaaaaaa aaaaaaaaaa aaaagtttt 3279

```

<210> 340

<211> 154

<212> PRT

<213> Homo sapiens

<400> 340

```

Met Ala Gly Val Arg Asp Gln Gly Gln Gly Ala Arg Trp Pro His Thr
          5                                10                        15

```

```

Gly Lys Arg Gly Pro Leu Leu Gln Gly Leu Thr Trp Ala Thr Gly Gly
          20                                25                        30

```

```

His Cys Phe Ser Ser Glu Glu Ser Gly Ala Val Asp Gly Ala Gly Gln
          35                                40                        45

```

```

Lys Lys Asp Arg Ala Trp Leu Arg Cys Pro Glu Ala Val Ala Gly Phe
          50                                55                        60

```

```

Pro Leu Gly Ser Asp Cys Arg Glu Gly Gly Arg Gln Gly Cys Gly Gly
          65                                70                        75                        80

```

```

Ser Asp Asp Glu Asp Asp Leu Gly Val Ala Pro Gly Leu Ala Pro Ala
          85                                90                        95

```

```

Trp Ala Leu Thr Gln Pro Pro Ser Gln Ser Pro Gly Pro Gln Ser Leu
          100                               105                        110

```

```

Pro Ser Thr Pro Ser Ser Ile Trp Pro Gln Trp Val Ile Leu Ile Thr

```

115 120 125
 Glu Leu Thr Ile Pro Ser Pro Ala His Gly Pro Pro Trp Leu Pro Asn
 130 135 140
 Ala Leu Glu Arg Gly His Leu Val Arg Glu
 145 150

<210> 341
 <211> 718
 <212> DNA
 <213> Homo sapien

<220>
 <221> misc_feature
 <222> (1)...(718)
 <223> n = A,T,C or G

<400> 341
 ggagatttgt gtggtttgca gccgagggag accaggaaga tctgcatggt gggaaggacc 60
 tgatgataca gaggtgagaa ataagaaagg ctgctgactt taccatctga ggccacacat 120
 ctgctgaaat ggagataatt aacatcacta gaaacagcaa gatgacaata taatgtctaa 180
 gtagtgacat gtttttgcac atttccagcc cttttaaata tccacacaca caggaagcac 240
 aaaaggaagc acagagatcc ctgggagaaa tgcccggccg ccatcttggg tcatcgatga 300
 gcctcgccct gtgcctgntc ccgcttgtga gggaaggaca ttagaaaatg aattgatgtg 360
 ttccctaaag gatggcagga aaacagatcc tgttgtggat atttatttga acgggattac 420
 agatttgaaa tgaagtcaca aagtgagcat taccaatgag aggaaaacag acgagaaaat 480
 cttgatgggt cacaagacat gcaacaaaca aaatggaata ctgtgatgac acgagcagcc 540
 aactggggag gagataccac ggggcagagg tcaggattct ggccctgctg cctaactgtg 600
 cgttatacca atcattttcta tttctaccct caaacaagct gtngaataatc tgacttacgg 660
 ttcttntggc ccacattttc atnatccacc cctcctttt aaannttantic caaantgt 718

<210> 342
 <211> 3112
 <212> DNA
 <213> Homo sapiens

<400> 342
 catttgtgtt ggagaaaaac agaggggaga tttgtgtggc tgcagccgag ggagaccagg 60
 aagatctgca tgggtgggaag gacctgatga tacagatgtt gataggagac aattaaaggc 120
 tgggaaggcag tggatgcctg atgatgaagt ggactttcaa actggggcac tactgaaacg 180
 atgggatggc cagagacaca ggagatgagt tggagcaagc tcaataacaa agtggttcaa 240
 cgaggacttg gaattgcatg gagctggagc tgaagttagt cccaattgtt tactagttag 300
 gtgaatgtgg atgattggat gatcatttct catctctgag cctcaggttc cccatccata 360
 aaatgggata cacagtatga tctataaagt gggatatagt atgatctact tcaactgggtt 420
 atttgaagga tgaattgaga taattttattt cagggtgccta gaacaatgcc cagattagta 480
 catttggttg aactgagaaa tggcataaca ccaaatttaa tatatgtcag atgttactat 540
 gattatcatt caatctcata gttttgtcat ggcccaattt atcctcactt gtgcctcaac 600
 aaattgaact gttaacaaag gaatctctgg tcctgggtaa tggctgagca ccaactgagca 660
 tttccattcc agttggcttc ttgggtttgc tagctgcata actagtcata ttaataaat 720
 gaagttttta cattttctcca gtgatttttt tatctcacct ttgaagatac tatgttatgt 780
 gattaaataa agaacttgag aagaacaggt ttcattaaac ataaaatcaa tgtagacgca 840
 aattttcttg atgggcaata cttatgttca caggaaatgc tttaaaatat gcagaagata 900
 attaaatggc aatggacaaa gtgaaaaact tagacttttt tttttttttt ggaagtatct 960
 ggatgttcct tagtcaacta aaggagaact gaaaaatagc agtgagttcc acataatcca 1020
 acctgtgaga ttaaggctct ttgtggggaa ggacaaagat ctgtaaattt acagtttctt 1080
 tccaaagcca acgtcgaatt ttgaaacata tcaaagctct tcttcaagac aaataatcta 1140
 tagtacatct ttcttatggg atgcacttat gaaaaatggg ggctgtcaac atctagtcac 1200

```

tttagctctc aaaatgggtc attttaagag aaagtttttag aatctcatat ttattcctgt 1260
ggaaggacag cattgtggct tggactttat aaggctcttta ttcaactaaa taggtgagaa 1320
ataagaaagg ctgctgactt taccatctga ggccacacat ctgctgaaat ggagataatt 1380
aacatcacta gaaacagcaa gatgacaata taatgtctaa gtagtgacat gtttttgcac 1440
atttccagcc ccttttaaata tccacacaca caggaagcac aaaaggaagc acagagatcc 1500
ctgggagaaa tgcccgccg ccatcttggtg tcatcgatga gcctcgccct gtgcctggtc 1560
ccgcttgatga gggaaggaca ttagaaaaatg aattgatgtg ttccttaaag gatgggcagg 1620
aaaacagatc ctggttgatga tatttatgtt aacgggatta cagatttgaa atgaagtcac 1680
aaagttagca ttaccaatga gaggaaaaca gacgagaaaa tcttgatggc ttcacaagac 1740
atgcaacaaa caaaatggaa tactgtgatg acatgaggca gccaaagctgg ggaggagata 1800
accacggggc agagggtcag gattctggcc ctgctgccta aactgtgcgt tcataacca 1860
atcatttcat atttctaacc ctcaaaacaa agctgttgta atatctgac tctacggttc 1920
cttctgggccc caacattctc catatatcca gccacactca tttttaatat ttagttccca 1980
gatctgtact gtgacctttc tacactgtag aataacatta ctcattttgt tcaaagacc 2040
ttcgtgttgc tgccataatg gtagctgact gtttttccta aggagtgttc tggcccagg 2100
gatctgtgaa caggctggga agcatctcaa gatctttcca gggttatact tactagcaca 2160
cagcatgac attacggagt gaattatcta atcaacatca tcctcagtgt ctttgcccat 2220
actgaaatto atttcccact tttgtgccc ttctcaagac ctcaaaatgt cattccatta 2280
atatcacagg attaactttt ttttttaacc tggaagaatt caatgttaca tgcagctatg 2340
ggaatttaac tacatatatt gttttccagt gcaaagatga ctaagtcctt tatccctccc 2400
ctttgtttga ttttttttcc agtataaagt taaaatgctt agccttgtag tgaggctgta 2460
tacagccaca gccctcctcc atccctccag ccttatctgt catcaccatc aacctctccc 2520
atgcacctaa acaaaatcta acttgtaatt ccttgaacat gtcaggcata cattattcct 2580
tctgcctgag aagctcttcc ttgtctctta aatctagaat gatgtaaagt tttgaataag 2640
ttgactatct tacttcatgc aaagaaggga cacatatgag attcatcatc acatgagaca 2700
gcaaatacta aaagtgtaat ttgattataa gagttagatg aatatatga aatgcaagag 2760
ccacagaggg aatgtttatg gggcacgttt gtaagcctgg gatgtgaagc aaaggcagg 2820
aacctcatag tatcttatat aatatacttc atttctctat ctctatcaca atatccaaca 2880
agcttttcac agaattcatg cagtgcacaa ccccaaaggt aacctttatc catttcatgg 2940
tgagtgcgct ttagaatttt ggcaaatcat actggtcact tatctcaact ttgagatgtg 3000
tttgtccttg tagttaattg aaagaaatag ggcactcttg tgagccactt tagggttcac 3060
tcctggcaat aaagaattta caaagagcaa aaaaaaaaaa aaaaaaaaaa aa 3112

```

<210> 343

<211> 2229

<212> DNA

<213> Homo sapiens

<400> 343

```

agctctttgt aaattcttta ttgccaggag tgaaccctaa agtggctcac aagagtgcc 60
tatttctttc aattaactac aaggacaaac acatctcaaa gttgagataa gtgaccagta 120
tgatttgcca aaattctaaa gcgcactcac catgaaatgg ataaaggtta cctttgggga 180
tttgactgct atgaattctg tgaaaagctt gttggatatt gtgatagaga tagagaaatg 240
aagtatatta tataagatac tatgagggtc cctgcctttg cttcacatcc caggcttaca 300
aacgtgcccc ataaacatto cctctgtggc tcttgcatth catatattha tctaaactct 360
tataatcaaa tacactttta gtatttgcgt totcatgtga tgatgaatct catatgtgtc 420
ccttctttgc atgaagtaag atagtcaact tattcaaaac tttacatcat tctagattta 480
agagacaagg aagagcttct caggcagaag gaataatgta tgctgacat gttcaaggaa 540
ttacaagtta gattttgttt aggtgcatgg gaggggttga tgggtgatgac agataaggct 600
ggaggggatg ggagaggctg tggctgtata cagcctcagt acaaggctaa gcattttaac 660
tttatactgg aaaaaaaatc aaacaaaggg gagggataaa ggacttagtc atctttgcac 720
tggaaaacaa aatatgtaat taaattccca tagctgcatg taacattgaa ttcttcagg 780
ttaaaaaaaa agttaatcct gtgatattaa tggaatgaca ttttgaggtc ttgagaatgg 840
gcacaaaagt gggaaatgaa tttcagtatg ggcaaagaca ctgaggatga tgttgattag 900
ataattcact ccgtaattgat catgctgtgt gctagtaagt ataaccctgg aaagatcttg 960
agatgcttcc cagcctgttc acagatcccc tggggcagaa cactccttag gaaaaacagt 1020
cagctacata ttaggcagca acacgaaggg tctttgaaca aaatgagtaa tgttattcta 1080
cagtgtagaa aggtcacagt acagatctgg gaactaaata ttaaaaaatga gtgtggctgg 1140

```

atatatggag	aatgttgggc	ccagaaggaa	ccgtagagat	cagatattac	aacagctttg	1200
ttttgagggt	tagaaatatg	aaatgatttg	gttatgaacg	cacagtttag	gcagcagggc	1260
cagaatcctg	accctctgcc	ccgtgggtat	ctcctcccca	gcttggtctg	ctcatgtcat	1320
cacagatttc	cattttgttt	gttgcattgc	ttgtgaagcc	atcaagattt	tctcgtctgt	1380
tttccctctca	ttggtaaatg	tcactttgtg	acttcatttc	aaatctgtaa	tcccgttcaa	1440
ataaatatcc	acaacaggat	ctgttttctt	gccccatcct	taagggaacac	atcaattcat	1500
tttctaattg	ccttccctca	caagcgggac	caggcacagg	gcgaggctca	tcgatgaccc	1560
aagatggcgg	ccgggcattt	ctcccaggga	tctctgtgct	tccttttgtg	cttcctgtgt	1620
gtgtggatat	ttaaaggggc	tggaaatgtg	caaaaacatg	tcactactta	gacattatat	1680
tgtcatcttg	ctgtttctag	tgatgttaat	tatctccatt	tcagcagatg	tgtggcctca	1740
gatggtaaag	tcagcagcct	ttcttatttc	tcacctggaa	atacatacga	ccatttgagg	1800
agacaaatgg	caaggtgtca	gcataccctg	aacttgagtt	gagagctaca	cacaatatta	1860
ttggtttccg	agcatcacia	acaccctctc	tgtttcttca	ctgggcacag	aattttaata	1920
cttatttcag	tgggctgttg	gcaggaacaa	atgaagcaat	ctacataaag	tcactagtgc	1980
agtgcctgac	acacaccatt	ctcttgaggt	ccccctctaga	gatcccacag	gtcatatgac	2040
ttcttgggga	gcagtggctc	acacctgtaa	tcccagcact	ttgggaggct	gaggcaggtg	2100
ggtcacctga	ggtcaggagt	tcaagaccag	cctggccaat	atggtgaaac	cccctctcta	2160
ctaaaaatac	aaaaattagc	tgggcgtgct	ggtgcatgcc	tgtaatccca	gccccaacac	2220
aatggaatt						2229

<210> 344

<211> 2426

<212> DNA

<213> Homo sapiens

<400> 344

gtaaattctt	tattgccagg	agtgaaccct	aaagtggctc	acaagagtgc	cctattttctt	60
tcaattaact	acaaggacaa	acacatctca	aagttgagat	aagtgaccag	tatgatttgc	120
caaaattcta	aagcgcactc	accatgaaat	ggataaagg	tacctttggg	gatttgcact	180
gcatgaattc	tgtgaaaagc	ttgttgata	ttgtgataga	gatagagaaa	tgaagtatat	240
tatataagat	actatgaggt	tccctgcctt	tgcttcacat	cccaggctta	caaacgtgcc	300
ccataaacad	tccctctgtg	gctcttgcct	ttcatatatt	tatctaaact	cttataatca	360
aattacactt	ttagtatttg	ctgtctcatg	tgatgatgaa	tctcatatgt	gtcccttctt	420
tgcatgaagt	aagatagtca	acttattcaa	aactttacat	cattctagat	ttaagagaca	480
aggaagagct	tctcaggcag	aaggaataat	gtatgcctga	catgttcaag	gaattacaag	540
ttagattttg	tttaggtgca	tgggaggggt	tgatggtgat	gacagataag	gctggaggga	600
tggggagagg	ctgtggctgt	atacagcctc	agtacaaggc	taagcatttt	aactttatac	660
tggaaaaaaa	atcaaacaaa	ggggagggat	aaaggactta	gtcatctttg	cactggaaaa	720
caaaatattg	aattaaattc	ccatagctgc	atgtaacatt	gaattcttcc	aggttaaaaa	780
aaaaagttaa	tcctgtgata	ttaatggaat	gactttttga	ggctctgaga	atgggcacaa	840
aagtgggaaa	tgaatttcag	tatgggcaaa	gacactgagg	atgatgttga	ttagataatt	900
cactccgtaa	tgatcatgct	gtgtgctagt	aagtataacc	ctggaaaagat	cttgagatgc	960
ttcccagcct	gttcacagat	cccctgggcc	agaacactcc	ttaggaaaaa	cagtcagcta	1020
catattaggc	agcaacacga	agggctcttg	aacaaaatga	gtaatgttat	tctacagtgt	1080
agaaaggtca	cagtacagat	ctgggaacta	aatattaaaa	atgagtgtgg	ctggatatat	1140
ggagaatggt	gggcccagaa	ggaaccgtag	agatcagata	ttacaacagc	tttgttttga	1200
gggttagaaa	tatgaaatga	tttggttatg	aacgcacagt	ttaggcagca	gggccagagt	1260
cctgaccctc	tgccccgtgg	ttatctcctc	cccagcttgg	ctgcctcatg	tcatacacagt	1320
attccatttt	gtttgttgca	tgtcttgtga	agccatcaag	attttctcgt	ctgttttctt	1380
ctcattggta	atgctcactt	tgtgacttca	tttcaaactc	gtaatccgt	tcaaataaat	1440
atccacaaca	ggatctgttt	tcctgcccct	cctttaaggga	acacatcaat	tcatttttcta	1500
atgtccttcc	ctcacaagcg	ggaccaggca	cagggcgagg	ctcatcgatg	acccaagatg	1560
gcggccgggc	atttctccca	gggatctctg	tgcttccttt	tgtgcttctt	gtgtgtgtgg	1620
atattttaaag	gggctggaaa	tgtgcaaaaa	catgtcacta	cttagacatt	atattgtcat	1680
cttgctgttt	ctagtgtgtg	taattatctc	catttcagca	gatgtgtggc	ctcagatggg	1740
aaagtcaagca	gcctttctta	tttctcacct	ggaaatacat	acgaccattt	gaggagacaa	1800
atggcaaggt	gtcagcatat	cctgaacttg	agttgagagc	tacacacaat	attattgggt	1860
tccgagcatc	acaaacaccc	tctctgtttc	ttcactgggc	acagaatttt	aatacttatt	1920

```

tcagtgggct gttggcagga acaaatgaag caatctacat aaagtcacta gtgcagtgcc 1980
tgacacacac cattctcttg aggtcccctc tagagatccc acaggtcata tgacttcttg 2040
gggagcagtg gctcacacct gtaatcccag cactttggga ggctgaggca ggtgggtcac 2100
ctgaggtcag gaggttcaaga ccagcctggc caatatgggtg aaaccccatc tctactaaaa 2160
atacaaaaaa tagctgggcg tgctggtgca tgcctgtaat ccagctact tgggaggctg 2220
aggcaggaga attgctggaa catgggaggc ggaggttgca gtgagctgta attgtgccat 2280
tgactcga cctgggcgac agagtggaa tctgtttcca aaaaacaaac aaacaaaaaa 2340
ggcatagtca gatacaacgt ggggtgggatg tgtaaataga agcaggatat aaagggcagtg 2400
gggtgacggt tttgccaac acaatg
2426

```

<210> 345

<211> 812

<212> DNA

<213> Homo sapiens

<400> 345

```

gaacaaaatg agtaatgtta ttctacagt tagaaaggct acagtacaga tctgggaact 60
aaatattaaa aatgagtgtg gctggatata tggagaatgt tgggccaga aggaaccgta 120
gagatcagat attacaacag ctttgttttg agggtagaa atatgaaatg atttggttat 180
gaacgcacag tttaggcagc agggccagaa tcctgaccct ctgccccgtg gttatctcct 240
ccccagcttg gctgcctcat gtcacacag tattccattt tgttgttg atgtcttg 300
aagccatcaa gattttctcg tctgttttcc tctcattggg aatgctcact ttgtgacttc 360
atttcaaate tgtaatcccg ttcaaataaa tatccacaac aggatctgtt ttctgcca 420
tcctttaagg aacacatcaa ttcatTTTTt aatgtccttc ctcacaagc gggaccaggc 480
acagggcgag gctcatcgat gacccaagat ggcgccggg catttctccc agggatctct 540
gtgcttcctt ttgtgcttcc tgtgtgtgtg gatatttaaa ggggctggaa atgtgcaaaa 600
acatgtcact acttagacat tatattgtca tcttgctgtt tctagtgtg ttaattatct 660
ccatttcagc agatgtgtgg cctcagatgg taaagtcagc agcctttctt atttctcacc 720
tctgtatcat caggctcttc ccaccatgca gatcttctcg gtctcctcgt gctgcagcca 780
cacaaatctc cctctgttt ttctgatgcc ag
812

```

<210> 346

<211> 2417

<212> DNA

<213> Homo sapien

<400> 346

```

ggcgccgct ctgagctag tgggatcccc cgggctgcac gaattcggca cgagtgagtt 60
ggagttttac ctgtattgtt ttaatttcaa caagcctgag gactagccac aaatgtaccc 120
agtttacaaa tggagaaaca ggtgcaaaaa ggttggtacc tgtcaaaggc cgtatgtggc 180
agagccaaga tttgagccca gttatgtctg atgaacttag cctatgctct ttaaacttct 240
gaatgctgac cattgaggat atctaaactt agatcaattg cattttccct ccaagactat 300
ttacttatca atacaataat accaccttta ccaatctatt gttttgatac gagactcaaa 360
tatgccagat atatgtaaaa gcaacctaca agctctctaa tcatgctcac ctaaaagatt 420
cccggtatct aataggctca aagaaacttc ttctagaaat ataaaagaga aaattggatt 480
atgcaaaaaa tcattattaa tttttttcat ccactcttta attcagcaaa catttatctg 540
ttgttgactt tatgcagtat ggctttttaa ggattggggg acaggtgaag aacggggtgc 600
cagaatgcat cctcctacta atgaggctag tacacatttg cattttaaaa tgccctgtcc 660
agctgggcat ggtggatcat gcctgtaate tcaacattgg aaggccaagg caggaggatt 720
gcttcagccc aggagttcaa gaccagcctg ggcaacatag aaagacccca tctctcaatc 780
aatcaatcaa tgccctgtct ttgaaaataa aactctttta gaaaggttta atgggcaggg 840
tgtggtagct catgcctata atacagcact ttgggaggct gaggcaggag gatcacttta 900
gccagaagt tcaagaccag cctgggcaac aagtgcacc tcatctcaat tttttaataa 960
aatgaatata tacataagga aagataaaaa gaaaagttta atgaaagaat acagtataaa 1020
acaaatctct tggacctaaa agtatttttg ttcaagccaa atattgtgaa tcacctctct 1080
gtgttgagga tacagaatat ctaagcccag gaaactgagc agaaagttca tgtactaact 1140
aattcaaccg aggcaggca aaaatgagc taactaatca atccaggca agggcaaat 1200
tagacggaac ctgactctgg tctattaagc gacaactttc cctctgttgt attttcttt 1260

```


tattcaatgt	aaaaggataa	aaactctcta	aaactaaaaa	caatgtttgt	caggagttac	1320
aaaccatgac	caactaatta	tggggaatca	taaaatatga	ctgtatgaga	tcttgatggt	1380
ttacaaagt	tacccactgt	taatcacttt	aaacattaat	gaacttaaaa	atgaatttac	1440
ggagattgga	atgtttcttt	cctgttgat	tagttggctc	aggctgcat	aacaaaatac	1500
cacagactgg	gaggcttaag	taacagaaat	tcatttctca	cagttctggg	ggctggaagt	1560
ccacgatcaa	ggtgcaggaa	aggcaggctt	cattctgagg	cccctctctt	ggctcacatg	1620
tggccaccct	cccactgctg	gctcacatga	cctctttgtg	ctcctggaaa	gaggggtgtg	1680
gggacagagg	gaaagagaag	gagagggaa	tctctgggtg	ctcgtctttc	aaggacccta	1740
acctgggcca	ctttggccca	ggcactgtgg	ggtggggggg	tgtggctgct	ctgctctgag	1800
tggccaagat	aaagcaacag	aaaaatgtcc	aaagctgtgc	agcaaagaca	agccaccgaa	1860
cagggatctg	ctcatcagt	tggggacctc	caagtcggcc	accctggagg	caagcccca	1920
cagagcccat	gcaaggtggc	agcagcagaa	gaagggaatt	gtccctgtcc	ttggcacatt	1980
cctcaccgac	ctggtgatgc	tggacactgc	gatgaatggt	aatgtggatg	agaatatgat	2040
ggactcccag	aaaaggagac	ccagctgctc	aggtggctgc	aaatcattac	agccttcac	2100
ctggggagga	actgggggcc	tggttctggg	tcagagagca	gcccagttag	ggtgagagct	2160
acagcctgtc	ctgccagctg	gatccccagt	cccgtcaac	cagtaatcaa	ggctgagcag	2220
atcaggcttc	ccggagctgg	tcttgggaag	ccagccctgg	ggtgagttgg	ctcctgctgt	2280
ggtactgaga	caatattgtc	ataaattcaa	tgcgcccttg	tatccctttt	tcttttttat	2340
ctgtctacat	ctataatcac	tatgcatact	agtctttgtt	agtgtttcta	ttcmacttaa	2400
tagagatatg	ttatact					2417

<210> 347

<211> 320

<212> DNA

<213> Homo sapien

<400> 347

cccctgaagg	cttcttggtta	gaaaatagta	cagttacaac	caataggaac	aacaaaaaga	60
aaaagtttgt	gacattgtag	tagggagtgt	gtaccctcta	ctcccatca	aaaaaaaaat	120
ggatacatgg	ttaaaggata	raagggaat	attttatcat	atgttctaaa	agagaaggaa	180
gagaaaatac	tactttctcr	aaatggaagc	ccttaaaggt	gctttgatac	tgaaggacac	240
aaatgtggcc	gtccatcctc	ctttaragtt	gcatgacttg	gacacggtaa	ctgttgagct	300
tttaractcm	gcattgtgac					320

<210> 348

<211> 539

<212> DNA

<213> Homo sapien

<400> 348

acgggactta	tcaaataaag	ataggaaaag	aagaaaactc	aaatattata	ggcagaaatg	60
ctaaagggtt	taaaatatgt	caggattgga	agaaggcatg	gataaagaac	aaagttcagt	120
taggaaagag	aaacacagaa	ggaagagaca	caataaaaag	cattatgtat	tctgtgagaa	180
gtcagacagt	aagattttgt	ggaaatgggt	tggtttggtg	tatggtatgt	attttagcaa	240
taatctttat	ggcagagaaa	gctaaaaatc	tttagcttgc	gtgaatgatc	acttgctgaa	300
ttcctcaagg	taggcatgat	gaaggagggt	ttagaggaga	cacagacaca	atgaactgac	360
ctagatagaa	agccttagta	tactcagcta	ggaatagtga	ttctgagggc	acactgtgac	420
atgattatgt	cttaccatgt	atggtagtga	tggggatgat	aggaaggaa	aacttatggc	480
atattttcac	ccccacaaaa	gtcagttaaa	tattgggaca	ctaaccatcc	aggtcaaga	539

<210> 349

<211> 2984

<212> DNA

<213> Homo sapien

<400> 349

atccctcctt	cccactctc	ctttccagaa	ggcacttggt	gtcttatctg	ttggactctg	60
------------	-----------	------------	------------	------------	------------	----

aaaacacttc	aggcgccctt	ccaaggcttc	cccaaacc	taagcagccg	cagaagcgct	120
cccagagctgc	cttctccac	actcaggtga	tgcagttgga	gaggaagttc	agccatcaga	180
agtacctgtc	ggcccctgaa	cgggccacc	tggccaagaa	cctcaagctc	acggagaccc	240
aagtgaagat	atggttccag	aacagacgct	ataagactaa	gcgaaagcag	ctctctcgg	300
agctgggaga	cttggaag	cactcctctt	tgcggccct	gaaagaggag	gccttctccc	360
gggcctccct	ggtctccgtg	tataacagct	atccttacta	cccatacctg	tactgctgg	420
gcagctggag	cccagctttt	tggtaatgcc	agctcaggtg	acaaccatta	tgatcaaaaa	480
ctgccttccc	cagggtgtct	ctatgaaaag	cacaaggggc	caaggtcagg	gagcaagagg	540
tgtgcacacc	aaagctattg	gagatttgcg	tggaaatctc	asattcttca	ctggtgagac	600
aatgaaacaa	cagagacagt	gaaaagttta	atacctaagt	cattccccc	gtgcatactg	660
taggtcattt	tttttgcctt	tggtacactg	tttgaagggg	agagagggaa	aatcaagtgg	720
tattttccag	cactttgtat	gattttggat	gagctgtaca	cccaaggatt	ctgttctgca	780
actccatcct	cctgtgtcac	tgaatatcaa	ctctgaaaga	gcaaaccata	caggagaaag	840
gacaaccagg	atgagatgt	caccaactga	attaaactta	agtccagaag	cctcctgttg	900
gccttggaa	atggccaagg	ctctctctgt	ccctgtaaaa	gagaggggca	aatagagagt	960
ctccaagaga	acgccctcat	gctcagcaca	tatttgcattg	ggagggggag	atgggtggga	1020
ggagatgaaa	atatcagctt	ttcttattcc	tttttattcc	ttttaaaatg	gtatgccaac	1080
ttaagtattt	acaggggtgg	ccaaatagaa	caagatgcac	tcgctgtgat	tttaagacaa	1140
gctgtataaa	cagaactcca	ctgcaagagg	gggggcccgg	ccaggagaat	ctccgcttgt	1200
ccaagacagg	ggcctaagga	gggtctccac	actgctgcta	ggggctgttg	cattttttta	1260
ttagtagaaa	gtggaaaggc	ctcttctcaa	cttttttccc	ttgggctgga	gaatttagaa	1320
tcagaagttt	ctggagttt	tcaggctatc	atatatactg	tatcctgaaa	ggcaacataa	1380
ttcttccttc	cctcctttta	aaattttgtg	ttcctttttg	cagcaattac	tcactaaagg	1440
gcttcatttt	agtccagatt	tttagtctgg	ctgcacctaa	cttatgcctc	gcttatttag	1500
cccagatct	ggtctttttt	tttttttttt	tttttccgtc	tcccaaagc	tttatctgtc	1560
ttgacttttt	aaaaaagttt	gggggcagat	tctgaattgg	ctaaaagaca	tgcattttta	1620
aaactagcaa	ctcttatttc	tttcctttta	aaatacatag	cattaaatcc	caaatcctat	1680
ttaaagacct	gacagcttga	gaaggtcact	actgcattta	taggaccttc	tggtggttct	1740
gctgttacgt	ttgaagtctg	acaatccttg	agaatccttg	catgcagagg	aggtaagagg	1800
tattggattt	tcacagagga	agaacacagc	gcagaatgaa	gggccaggct	tactgagctg	1860
tccagtggag	ggctcatggg	tgggacatgg	aaaagaaggc	agcctaggcc	ctggggagcc	1920
cagtccactg	agcaagcaag	ggactgagtg	agccttttgc	aggaaaaggc	taagaaaaag	1980
gaaaaccatt	ctaaaacaca	acaagaaact	gtccaaatgc	tttgggaact	gtgtttattg	2040
cctataatgg	gtcccaaaaa	tgggtaacct	agacttcaga	gagaatgagc	agagagcaaa	2100
ggagaaatct	ggctgtcctt	ccattttcat	tctgttatct	caggtgagct	ggtagagggg	2160
agacattaga	aaaaaatgaa	acaacaaaac	aattactaat	gaggtacgct	gaggcctggg	2220
agtctcttga	ctccactact	taattccgtt	tagtgagaaa	cctttcaatt	ttcttttatt	2280
agaagggcca	gcttactgtt	ggtggcaaaa	ttgccaat	aagttaatag	aaagttggcc	2340
aatttcaccc	cttttctgt	ggtttgggct	ccacattgca	atgttcaatg	ccacgtgctg	2400
ctgacaccga	cggagtagt	agccagcaca	aaagcgagg	tagcctgaat	tgctttctgc	2460
tctttacatt	tcttttaaaa	taagcattta	gtgctcagtc	cctactgagt	actctttctc	2520
tcccctcttc	tgaattta	tctttcaact	tgcaatttgc	aaggattaca	catttctactg	2580
tgatgtatat	tgtgttgcaa	aaaaaaaaaa	aagtgtcttt	gtttaaaatt	acttggtttg	2640
tgaatccatc	ttgctttttc	ccatttgga	ctagtcat	acccatctct	gaactggtag	2700
aaaaacatct	gaagagctag	tctatcagca	tctgacaggt	gaattggatg	gttctcagaa	2760
ccatttcacc	cagacagcct	gtttctatcc	tgtttaataa	attagtttgg	gttctctaca	2820
tgcataacaa	accctgctcc	aatctgtcac	ataaaaagtc	gtgacttgaa	gtttagtcag	2880
cacccccacc	aaactttatt	tttctatgtg	ttttttgcaa	catatgagtg	ttttgaaaa	2940
aaagtaccca	tgtcttttatt	agaaaaaaaa	aaaaaaaaaa	aaaa		2984

<210> 350

<211> 147

<212> PRT

<213> Homo sapien

<400> 350

Pro	Ser	Phe	Pro	Thr	Leu	Leu	Ser	Arg	Arg	His	Leu	Gly	Ser	Tyr	Leu
1				5					10					15	

Leu Asp Ser Glu Asn Thr Ser Gly Ala Leu Pro Arg Leu Pro Gln Thr
 20 25 30
 Pro Lys Gln Pro Gln Lys Arg Ser Arg Ala Ala Phe Ser His Thr Gln
 35 40 45
 Val Ile Glu Leu Glu Arg Lys Phe Ser His Gln Lys Tyr Leu Ser Ala
 50 55 60
 Pro Glu Arg Ala His Leu Ala Lys Asn Leu Lys Leu Thr Glu Thr Gln
 65 70 75 80
 Val Lys Ile Trp Phe Gln Asn Arg Arg Tyr Lys Thr Lys Arg Lys Gln
 85 90 95
 Leu Ser Ser Glu Leu Gly Asp Leu Glu Lys His Ser Ser Leu Pro Ala
 100 105 110
 Leu Lys Glu Glu Ala Phe Ser Arg Ala Ser Leu Val Ser Val Tyr Asn
 115 120 125
 Ser Tyr Pro Tyr Tyr Pro Tyr Leu Tyr Cys Val Gly Ser Trp Ser Pro
 130 135 140
 Ala Phe Trp
 145

<210> 351
 <211> 1289
 <212> DNA
 <213> Homo sapien

<400> 351
 agccaggcgt ccctctgcct gccactcag tggcaacacc cgggagctgt tttgtccttt 60
 gtggagcctc agcagttccc tctttcagaa ctactgcca agagccctga acaggagcca 120
 ccatgcagtg cttcagcttc attaagacca tgatgaccc cttcaatttg ctcactcttc 180
 tgtgtggtgc agccctgttg gcagtgaggc tctgggtgtc aatcgatggg gcactccttc 240
 tgaagatctt cgggccactg tgcgtccagt ccatgcagtt tgtcaacgtg ggctacttcc 300
 tcatcgacgc cggcggttggt gtctttgtct ttgggttccct gggctgctat ggtgctaaga 360
 ctgagagcaa gtgtgccctc gtgacgttct tcttcactcct cctcctcatc ttcattgctg 420
 aggttgacgc tgctgtggtc gccttggtgt acaccacaat ggctgagcac ttcctgacgt 480
 tgctggtagt gctgcccac aagaaagatt atgggttccca ggaagacttc actcaagtgt 540
 ggaacaccac catgaaaggc ctcaagtgtc gtggcttcac caactatacg gattttgagg 600
 actcacccta cttcaaagag aacagtgcct tccccccatt ctgttgcaat gacaacgtca 660
 ccaacacagc caatgaaacc tgcaccaagc aaaaggctca cgacaaaaaa gtagagggtt 720
 gcttcaatca gcttttgtat gacatccgaa ctaatgcagt caccgtgggt ggtgtggcag 780
 ctggaattgg gggcctcgag ctggctgcca tgattgtgtc catgtatctg tactgcaatc 840
 tacaataagt ccacttctgc ctctgccact actgctgcca catgggaact gtgaagaggc 900
 accctggcaa gcagcagtgga ttggggggagg ggacaggatc taacaatgtc acttgggcca 960
 gaatggacct gccctttctg ctccagactt ggggctagat agggaccact ccttttagcg 1020
 atgcctgact ttcttccat tgggtgggtgg atgggtgggg ggcattccag agcctctaag 1080
 gtagccagtt ctgttgccca ttccccagc ctattaaacc cttgatatgc cccctaggcc 1140
 tagtgggtgat cccagtgtct tactggggga tgagagaaag gcattttata gcctgggcat 1200
 aagtgaatc agcagagcct ctgggtggat gtgtagaagg cacttcaaaa tgcataaacc 1260
 tgttacaatg ttaaaaaaaaa aaaaaaaaaa 1289

<210> 352
 <211> 241
 <212> PRT
 <213> Homo sapien

<400> 352
 Met Gln Cys Phe Ser Phe Ile Lys Thr Met Met Ile Leu Phe Asn Leu
 1 5 10 15
 Leu Ile Phe Leu Cys Gly Ala Ala Leu Leu Ala Val Gly Ile Trp Val
 20 25 30

Ser Ile Asp Gly Ala Ser Phe Leu Lys Ile Phe Gly Pro Leu Ser Ser
 35 40 45
 Ser Ala Met Gln Phe Val Asn Val Gly Tyr Phe Leu Ile Ala Ala Gly
 50 55 60
 Val Val Val Phe Ala Leu Gly Phe Leu Gly Cys Tyr Gly Ala Lys Thr
 65 70 75 80
 Glu Ser Lys Cys Ala Leu Val Thr Phe Phe Phe Ile Leu Leu Leu Ile
 85 90 95
 Phe Ile Ala Glu Val Ala Ala Ala Val Val Ala Leu Val Tyr Thr Thr
 100 105 110
 Met Ala Glu His Phe Leu Thr Leu Leu Val Val Pro Ala Ile Lys Lys
 115 120 125
 Asp Tyr Gly Ser Gln Glu Asp Phe Thr Gln Val Trp Asn Thr Thr Met
 130 135 140
 Lys Gly Leu Lys Cys Cys Gly Phe Thr Asn Tyr Thr Asp Phe Glu Asp
 145 150 155 160
 Ser Pro Tyr Phe Lys Glu Asn Ser Ala Phe Pro Pro Phe Cys Cys Asn
 165 170 175
 Asp Asn Val Thr Asn Thr Ala Asn Glu Thr Cys Thr Lys Gln Lys Ala
 180 185 190
 His Asp Gln Lys Val Glu Gly Cys Phe Asn Gln Leu Leu Tyr Asp Ile
 195 200 205
 Arg Thr Asn Ala Val Thr Val Gly Gly Val Ala Ala Gly Ile Gly Gly
 210 215 220
 Leu Glu Leu Ala Ala Met Ile Val Ser Met Tyr Leu Tyr Cys Asn Leu
 225 230 235 240
 Gln

<210> 353

<211> 173

<212> DNA

<213> Homo sapien

<220>

<221> misc_feature

<222> (1)...(173)

<223> n = A,T,C or G

<400> 353

acattgtttt	tttgagataa	agcattgana	gagctctcct	taacgtgaca	caatggaagg	60
actggaacac	ataccacat	ctttgtttctg	agggataatt	ttctgataaa	gtcttgctgt	120
atattcaagc	acatatgtta	tatattattc	agttccatgt	ttatagccta	gtt	173

<210> 354

<211> 676

<212> DNA

<213> Homo sapien

<400> 354

gaaattaagt	atgagctaaa	ttccctgtta	aaacctotag	gggtgacaga	tctcttcaac	60
cagggtcaaag	ctgatctttc	tggaatgtca	ccaaccaagg	gcctatatatt	atcaaaagcc	120
atccacaagt	catacctgga	tgtcagcgaa	gagggcacgg	aggcagcagc	agccactggg	180
gacagcatcg	ctgtaaaaag	cctaccaatg	agagctcagt	tcaaggcgaa	ccacccttc	240
ctgttcttta	taaggcacac	tcataccaac	acgatcctat	tctgtggcaa	gcttgccctc	300
ccctaatacag	atgggggttg	gtaagggtca	gagttgcaga	tgaggtgcag	agacaatcct	360
gtgactttcc	cacggccaaa	aagctgttca	caacctcacgc	acctctgtgc	ctcagtttgc	420
tcactgtcaa	aataggtcta	ggatttcttc	caaccatttc	atgagttgtg	aagctaaggc	480
tttgttaatc	atggaaaaag	gtagacttat	gcagaaagcc	tttctggcct	tcttatctgt	540

ggtgtctcat ttgagtgtctg tccagtgaca tgatcaagtc aatgagtaaa attttaaggg 600
 attagatttt cttgacttgt atgtatctgt gagatcttga ataagtgacc tgacatctct 660
 gcttaaagaa aaccag 676

<210> 355
 <211> 463
 <212> DNA
 <213> Homo sapien

<400> 355
 acttcatcag gccataatgg gtgcctcccg tgagaatcca agcacctttg gactgcgcga 60
 tgtagatgag cgggctgaag atcttgcgca tgcgcggctt cagggcgaag ttcttggcgc 120
 ccccggtcac agaaatgacc aggttgggtg ttttcagggtg ccagtgtctg gtcagcagct 180
 cgtaaaggat ttccgcgtcc gtgtgcgagg acagacgtat atacttccct ttcttcccca 240
 gtgtctcaaa ctgaatatcc ccaaaggcgt cggtaggaaa ttcttgggtg tgtttcttgt 300
 agttccattt ctacttttgg ttgatctggg tgccttccat gtgctggctc tgggcatagc 360
 cacacttgca cacattctcc ctgataagca cgatggtgtg gacaggaagg aaggatttca 420
 ttgagcctgc ttatggaaac tgggtattgtt agcttaaata gac 463

<210> 356
 <211> 2106
 <212> DNA
 <213> Homo sapiens

<400> 356
 -atgcatcacc atcaccatca catgggctcc gacgttcgtg acctgaacgc actgctgccg 60
 gcagttccgt ccctgggtgg tgggtgggtg tgcgcactgc cggttagcgg tgcagcacag 120
 tgggctccgg ttctggaact cgcaccgcgg ggtgcatccg catacgggtc cctgggtggg 180
 ccggcaccgc cgcgggcacc gccgcgcggc ccgcgcggc cgcgcactc cttcatcaaa 240
 caggaaccga gctgggggtg tgcagaacc cagcaagaac agtgcctgag cgcattcacc 300
 gttcacttct ccggccagtt cactggcaca gccggagcct gtcgctacgg gcccttcggt 360
 cctcctccgc ccagccaggg gtcacccggc caggccagga tgtttcctaa cgcgccctac 420
 ctgcccagct gccctgagag ccagcccgtt attcgcaatc aggggttacag cacggtcacc 480
 ttcgacggga cgcaccagta cggtcacacg ccctgcgacc atgcggcgca gttccccaac 540
 cactcattca agcatgagga tcccatgggc cagcagggtc cgtggtgga gcagcagtag 600
 tcggtgccgc ccccggtcta tggctgccac acccccaccg acagctgcac cggcagccag 660
 gctttgctgc tgaggacgcc ctacagcagt gacaatttat accaaatgac atcccagctt 720
 gaatgatga cctggaatca gatgaactta ggagccacct taaagggcca cagcacaggg 780
 tacgagagcg ataaccacac aacgcccac cctgcggag cccaatacag aatacacacg 840
 cacggtgtct tcagaggcat tcaggatgtg ctgagtgatg ctggagtagc cccgactctt 900
 gtacggctcg catctgagac cagtgaagaa cgcaccttca tgtgtgctta ccaggctgc 960
 aataagagat attttaagct gtcccactta cagatgcaca gcaggaagca cactggtgag 1020
 aaaccatacc agtgtgactt caaggactgt gaacgaagg tttttcgttc agaccagctc 1080
 aaaagacacc aaaggagaca tacagggtgtg aaaccattcc agtgtaaaac ttgtcagcga 1140
 aagttctccc ggtccgacca cctgaagacc cacaccagga ctcatacagg tgaaaagccc 1200
 ttcagctgtc ggtggccaag ttgtcagaaa aagtttgccc ggtcagatga attagtccgc 1260
 catcacaaca tgcacagag aaacatgacc aaactccagc tggcgcttga attcatgtgg 1320
 gtcccgggtt tcttctcac cctgtccgtg acgtggattg gtgctgcacc cctcatcctg 1380
 tctcggattg tgggaggctg ggagtgcgag aagcattccc aaccctggca ggtgcttgtg 1440
 gcctctcgtg gcagggcagt ctgcggcggt gttctgggtg acccccagtg ggtcctcaca 1500
 gctgcccact gcatcaggaa caaaagcgtg atcttgctgg gtgcggcacag cctgtttcat 1560
 cctgaagaca caggccaggt atttcagggt agccacagct tcccacaccc gctctacgat 1620
 atgagcctcc tgaagaatcg attcctcagg ccagggtgat actccagcca cgacctcatg 1680
 ctgctccgcc tgtcagagcc tgccgagctc acggatgctg tgaaggatcat ggacctgccc 1740
 acccaggagc cagcactggg gaccacctgc tacgcctcag gctggggcag cattgaacca 1800
 gaggagtctc tgaccceaaa gaaacttcag tgtgtggacc tccatgttat ttccaatgac 1860
 gtgtgtgcgc aagttcaccc tcagaaggta accaagttca tgctgtgtgc tggacgctgg 1920
 acagggggca aaagcacctg ctcggtgat tctggggggc cacttgtctg taatggtgtg 1980

cttcaaggta tcacgtcatg gggcagtgaa ccatgtgcc tgcccgaag gccttcctg 2040
 tacaccaagg tgggtgcatta ccggaagtgg atcaaggaca ccatcgtggc caaccctga 2100
 gaattc 2106

<210> 357

<211> 692

<212> PRT

<213> Homo sapiens

<400> 357

Met Gly Ser Asp Val Arg Asp Leu Asn Ala Leu Leu Pro Ala Val Pro
 5 10 15

Ser Leu Gly Gly Gly Gly Gly Cys Ala Leu Pro Val Ser Gly Ala Ala
 20 25 30

Gln Trp Ala Pro Val Leu Asp Phe Ala Pro Pro Gly Ala Ser Ala Tyr
 35 40 45

Gly Ser Leu Gly Gly Pro Ala Pro Pro Pro Ala Pro Pro Pro Pro
 50 55 60

Pro Pro Pro Pro His Ser Phe Ile Lys Gln Glu Pro Ser Trp Gly Gly
 65 70 75 80

Ala Glu Pro His Glu Glu Gln Cys Leu Ser Ala Phe Thr Val His Phe
 85 90 95

Ser Gly Gln Phe Thr Gly Thr Ala Gly Ala Cys Arg Tyr Gly Pro Phe
 100 105 110

Gly Pro Pro Pro Pro Ser Gln Ala Ser Ser Gly Gln Ala Arg Met Phe
 115 120 125

Pro Asn Ala Pro Tyr Leu Pro Ser Cys Leu Glu Ser Gln Pro Ala Ile
 130 135 140

Arg Asn Gln Gly Tyr Ser Thr Val Thr Phe Asp Gly Thr Pro Ser Tyr
 145 150 155 160

Gly His Thr Pro Ser His His Ala Ala Gln Phe Pro Asn His Ser Phe
 165 170 175

Lys His Glu Asp Pro Met Gly Gln Gln Gly Ser Leu Gly Glu Gln Gln
 180 185 190

Tyr Ser Val Pro Pro Pro Val Tyr Gly Cys His Thr Pro Thr Asp Ser
 195 200 205

Cys Thr Gly Ser Gln Ala Leu Leu Leu Arg Thr Pro Tyr Ser Ser Asp
 210 215 220

Asn Leu Tyr Gln Met Thr Ser Gln Leu Glu Cys Met Thr Trp Asn Gln
 225 230 235 240

Met Asn Leu Gly Ala Thr Leu Lys Gly His Ser Thr Gly Tyr Glu Ser
 245 250 255

Asp Asn His Thr Thr Pro Ile Leu Cys Gly Ala Gln Tyr Arg Ile His
 260 265 270
 Thr His Gly Val Phe Arg Gly Ile Gln Asp Val Arg Arg Val Pro Gly
 275 280 285
 Val Ala Pro Thr Leu Val Arg Ser Ala Ser Glu Thr Ser Glu Lys Arg
 290 295 300
 Pro Phe Met Cys Ala Tyr Pro Gly Cys Asn Lys Arg Tyr Phe Lys Leu
 305 310 315 320
 Ser His Leu Gln Met His Ser Arg Lys His Thr Gly Glu Lys Pro Tyr
 325 330 335
 Gln Cys Asp Phe Lys Asp Cys Glu Arg Arg Phe Phe Arg Ser Asp Gln
 340 345 350
 Leu Lys Arg His Gln Arg Arg His Thr Gly Val Lys Pro Phe Gln Cys
 355 360 365
 Lys Thr Cys Gln Arg Lys Phe Ser Arg Ser Asp His Leu Lys Thr His
 370 375 380
 Thr Arg Thr His Thr Gly Glu Lys Pro Phe Ser Cys Arg Trp Pro Ser
 385 390 395 400
 Cys Gln Lys Lys Phe Ala Arg Ser Asp Glu Leu Val Arg His His Asn
 405 410 415
 Met His Gln Arg Asn Met Thr Lys Leu Gln Leu Ala Leu Glu Phe Met
 420 425 430
 Trp Val Pro Val Val Phe Leu Thr Leu Ser Val Thr Trp Ile Gly Ala
 435 440 445
 Ala Pro Leu Ile Leu Ser Arg Ile Val Gly Gly Trp Glu Cys Glu Lys
 450 455 460
 His Ser Gln Pro Trp Gln Val Leu Val Ala Ser Arg Gly Arg Ala Val
 465 470 475 480
 Cys Gly Gly Val Leu Val His Pro Gln Trp Val Leu Thr Ala Ala His
 485 490 495
 Cys Ile Arg Asn Lys Ser Val Ile Leu Leu Gly Arg His Ser Leu Phe
 500 505 510
 His Pro Glu Asp Thr Gly Gln Val Phe Gln Val Ser His Ser Phe Pro
 515 520 525
 His Pro Leu Tyr Asp Met Ser Leu Leu Lys Asn Arg Phe Leu Arg Pro
 530 535 540
 Gly Asp Asp Ser Ser His Asp Leu Met Leu Leu Arg Leu Ser Glu Pro
 545 550 555 560

Ala Glu Leu Thr Asp Ala Val Lys Val Met Asp Leu Pro Thr Gln Glu
 565 570 575
 Pro Ala Leu Gly Thr Thr Cys Tyr Ala Ser Gly Trp Gly Ser Ile Glu
 580 585 590
 Pro Glu Glu Phe Leu Thr Pro Lys Lys Leu Gln Cys Val Asp Leu His
 595 600 605
 Val Ile Ser Asn Asp Val Cys Ala Gln Val His Pro Gln Lys Val Thr
 610 615 620
 Lys Phe Met Leu Cys Ala Gly Arg Trp Thr Gly Gly Lys Ser Thr Cys
 625 630 635 640
 Ser Gly Asp Ser Gly Gly Pro Leu Val Cys Asn Gly Val Leu Gln Gly
 645 650 655
 Ile Thr Ser Trp Gly Ser Glu Pro Cys Ala Leu Pro Glu Arg Pro Ser
 660 665 670
 Leu Tyr Thr Lys Val Val His Tyr Arg Lys Trp Ile Lys Asp Thr Ile
 675 680 685
 Val Ala Asn Pro
 690